arXiv:1008.0249v1 [math.DS] 2 Aug 2010

A proof of the Kuramoto's conjecture for a bifurcation structure of the infinite dimensional Kuramoto model

Faculty of Mathematics, Kyushu University, Fukuoka, 819-0395, Japan
Hayato CHIBA [1]

July 31 , 2010

Abstract

The Kuramoto model is a system of ordinary differential equations for describing synchronization phenomena defined as a coupled phase oscillators. In this paper, a bifurcation structure of the infinite dimensional Kuramoto model is investigated. For a certain non-selfadjoint linear operator, which defines a linear part of the Kuramoto model, the spectral theory on a space of generalized functions is developed with the aid of a rigged Hilbert space to avoid a continuous spectrum on the imaginary axis. Although the linear operator has an unbounded continuous spectrum on a Hilbert space, it is shown that it admits a spectral decomposition consisting of a countable number of eigenfunctions on a space of generalized functions. The semigroup generated by the linear operator is calculated by using the spectral decomposition to prove the linear stability of a steady state of the system. The center manifold theory is also developed on a space of generalized functions. It is proved that there exists a finite dimensional center manifold on a space of generalized functions, while a center manifold on a Hilbert space is of infinite dimensional because of the continuous spectrum on the imaginary axis. The results are applied to the stability and bifurcation theory of the Kuramoto model to obtain a bifurcation diagram conjectured by Kuramoto. If the coupling strength K between oscillators is smaller than some threshold K_c, the de-synchronous state proves to be asymptotically stable, and if K exceeds K_c, a nontrivial solution, which corresponds to the synchronization, bifurcates from the de-synchronous state.

Keywords: infinite dimensional dynamical systems; center manifold theory; continuous spectrum; spectral theory; rigged Hilbert space; coupled oscillators

---

[1] E mail address : chiba@math.kyushu-u.ac.jp

Contents

1 Introduction **3**

2 Continuous model **7**

3 Transition point formula and the linear instability **9**
3.1 Analysis of the operator $\sqrt{-1}\mathcal{M}$ 10
3.2 Eigenvalues of the operator T_1 and the transition point formula 11

4 Linear stability theory **19**
4.1 Resonance poles 19
4.2 Gaussian-type case 23
4.3 Rational case 29

5 Spectral theory **30**
5.1 Rigged Hilbert space 30
5.2 Spectral theory on $\mathrm{Exp}_+ \subset L^2(\mathbf{R}, g(\omega)d\omega) \subset \mathrm{Exp}'_+$ 32
5.3 Spectral theory on $F_p \subset L^2(\mathbf{R}, g(\omega)d\omega) \subset F'_p$ 48
5.4 Spectral theory on $(H_+, L^2(\mathbf{R}, g(\omega)d\omega), H'_+)$ 60

6 Nonlinear stability **61**

7 Bifurcation theory **65**
7.1 Center manifold theorem 66
7.2 Phase space of the perturbed system 68
7.3 Localization of the semiflow 74
7.4 Proof of the center manifold theorem 79
7.5 Reduction to the center manifold 89

A Resonance poles of the Gaussian distribution **95**

B Proof of the existence of solutions of Eq.(7.9) **97**



Fig. 1: The order parameter of the Kuramoto model.

1 Introduction

Collective synchronization phenomena are observed in a variety of areas such as chemical reactions, engineering circuits and biological populations [37]. In order to investigate such phenomena, Kuramoto [25] proposed the system of ordinary differential equations

$$\frac{d\theta_i}{dt} = \omega_i + \frac{K}{N}\sum_{j=1}^{N} \sin(\theta_j - \theta_i),\ \ i = 1, \cdots, N, \tag{1.1}$$

where $\theta_i = \theta_i(t) \in [0, 2\pi)$ is a dependent variable which denotes the phase of an i-th oscillator on a circle, $\omega_i \in \mathbf{R}$ denotes its natural frequency, $K > 0$ is a coupling strength, and where N is the number of oscillators. Eq.(1.1) is derived by means of the averaging method from coupled dynamical systems having limit cycles, and now it is called the *Kuramoto model*.

It is obvious that when $K = 0$, $\theta_i(t)$ and $\theta_j(t)$ rotate on a circle at different velocities unless ω_i is equal to ω_j, and it is true for sufficiently small $K > 0$. On the other hand, if K is sufficiently large, it is numerically observed that some of oscillators or all of them tend to rotate at the same velocity on average, which is called the *synchronization* [37, 42]. If N is small, such a transition from de-synchronization to synchronization may be well revealed by means of the bifurcation theory [11, 27, 28]. However, if N is large, it is difficult to investigate the transition from the view point of the bifurcation theory and it is still far from understood.

In order to evaluate whether synchronization occurs or not, Kuramoto introduced the *order parameter* $r(t)e^{\sqrt{-1}\psi(t)}$ by

$$r(t)e^{\sqrt{-1}\psi(t)} := \frac{1}{N}\sum_{j=1}^{N} e^{\sqrt{-1}\theta_j(t)}, \tag{1.2}$$

where $r, \psi \in \mathbf{R}$. The order parameter gives the centroid of oscillators. It seems that if synchronous state is formed, $r(t)$ takes a positive number, while if de-synchronization is stable, $r(t)$ is zero on time average (see Fig.1). Further, this is true for every t when N is sufficiently large so that statistical-mechanical description is applied. Based on this



Fig. 2: Typical bifurcation diagrams of the order parameter for the cases that (a) $g(\omega)$ is even and unimodal (b) $g(\omega)$ is even and bimodal. Solid lines denote stable solutions and dotted lines denote unstable solutions.

observation and some formal calculations, Kuramoto conjectured a bifurcation diagram of $r(t)$ as follows:

Kuramoto's conjecture

Suppose that $N \to \infty$ and natural frequencies ω_i's are distributed according to a probability density function $g(\omega)$. If $g(\omega)$ is an even and unimodal function such that $g''(0) \neq 0$, then the bifurcation diagram of $r(t)$ is given as Fig.2 (a); that is, if the coupling strength K is smaller than $K_c := 2/(\pi g(0))$, then $r(t) \equiv 0$ is asymptotically stable. On the other hand, if K is larger than K_c, the synchronous state emerges; there exists a positive constant r_c such that $r(t) = r_c$ is asymptotically stable. Near the transition point K_c, r_c is of order $O((K - K_c)^{1/2})$.

A function $g(\omega)$ is called unimodal (at $\omega = 0$) if $g(\omega_1) > g(\omega_2)$ for $0 \leq \omega_1 < \omega_2$ and $g(\omega_1) < g(\omega_2)$ for $\omega_1 < \omega_2 \leq 0$. Now the value $K_c = 2/(\pi g(0))$ is called the *Kuramoto's transition point*. See [26] and [42] for the Kuramoto's discussion.

In the last two decades, many studies to confirm the Kuramoto's conjecture have been done. Significant papers of Strogatz *et al.* [43, 44] investigated the linear stability of the trivial solution, which corresponds to the de-synchronous state $r \equiv 0$. In [43], they introduced the continuous model for the Kuramoto model to describe the situation $N \to \infty$. They derived the Kuramoto's transition point $K_c = 2/(\pi g(0))$ and showed that if $K > K_c$, the de-synchronous state is unstable because of eigenvalues on the right half plane. On the other hand, when $0 < K \leq K_c$, the linear operator T_1, which defines the linearized equation of the continuous model around the de-synchronous state, has no eigenvalues. However, in [44], they found that an analytic continuation of the resolvent $(\lambda - T_1)^{-1}$ may have poles (*resonance poles*) on the left half plane, and with the aid of numerical simulations, they remarked a possibility that resonance poles induce exponential decay of the order parameter r. This claim will be rigorously proved in this paper. In [33], the spectra of linearized systems around other steady states, which correspond to solutions with positive $r = r_c$, are investigated. They found that linear operators, which define linearized systems around synchronous states have continuous spectra on the imaginary axis. Nevertheless, they again remarked that such solutions can be asymptotically stable because of the resonance poles.

Since results of Strogatz *et al.* are based on the linearized analysis, effects of nonlinear terms are neglected. To investigate nonlinear dynamics, the bifurcation theory is often used. However, investigating the bifurcation structure near the transition point K_c involves further difficult problems because the operator T_1 has a continuous spectrum on the imaginary axis, that is, a center manifold in a usual sense is of infinite dimensional. To avoid this difficulty, Bonilla *et al.* [2, 7, 8] and Crawford *et al.* [12, 13, 14] added a perturbation (noise) with the strength $D > 0$ to the Kuramoto model. Then, the continuous spectrum moves to the left side by D, and thus the usual center manifold reduction is applicable. When $g(\omega)$ is an even and unimodal function, they obtained the Kuramoto's bifurcation diagram (Fig.2 (a)), however, obviously their methods are not valid when $D = 0$. For example, in Crawford's method, an eigenfunction of T_1 associated with a center subspace diverges as $D \to 0$ because an eigenvalue on the imaginary axis is embedded in the continuous spectrum as $D \to 0$. Thus the original Kuramoto's conjecture is still open.

Despite the active interest in the case that the distribution function $g(\omega)$ is even and unimodal, bifurcation diagrams of r for $g(\omega)$ other than the even and unimodal case are not understood well. Martens et al. [30] investigated the bifurcation diagram for a bimodal $g(\omega)$ which consists of two Lorentzian distributions. In particular, they found that stable synchronous states can coexist with stable de-synchronous states if K is slightly smaller than K_c (see Fig.2 (b)). Their analysis depends on extensive symmetries of the Kuramoto model found by Ott and Antonsen [35, 36] (see also [31]) and on the special form of $g(\omega)$, however, such a diagram seems to be common for any bimodal distributions.

In this paper, the stability, spectral and bifurcation theory of the continuous model of the Kuramoto model will be developed to prove the Kuramoto's conjecture. In particular, the spectrum and the semigroup of a linear operator T_1 on the weighted Lebesgue space $L^2(\mathbf{R}, g(\omega)d\omega)$, which is obtained by linearizing the continuous model around the de-synchronous state, will be investigated in detail. At first, we derive the transition point (bifurcation point) K_c for any distribution function $g(\omega)$. As a corollary, the Kuramoto's transition point $K_c = 2/(\pi g(0))$ is obtained if $g(\omega)$ is an even and unimodal function. When $K > K_c$, it is proved that the de-synchronous state is unstable because the operator T_1 has eigenvalues on the right half plane.

On the other hand, when $0 < K \leq K_c$, the operator T_1 has no eigenvalues and the continuous spectrum lies on the imaginary axis. Thus the stability of the de-synchronous state is nontrivial. Despite this fact, under appropriate assumptions for $g(\omega)$, the order parameter proves to decay exponentially to zero as $t \to \infty$ because of the existence of resonance poles on the left half plane, as was expected by Strogatz et al. [44]. To prove it, the notion of spectrum is extended. Roughly speaking, the spectrum is the set of singularities of the resolvent $(\lambda - T_1)^{-1}$. However, if $g(\omega)$ has an analytic continuation, the resolvent proves to have an analytic continuation, and it has singularities on the second Riemann sheet, which are called the resonance poles. By using the Laplace inversion formula for a semigroup, we will prove that the resonance poles induce exponential decay of the order parameter. It suggests that in general, linear stability of a trivial solution of a linear equation on an infinite dimensional space is determined by not only the spectrum of the linear operator but also its resonance poles.

Next purpose is to investigate a bifurcation from $K = K_c$. To handle the continuous spectrum on the imaginary axis, a spectral theory of the resonance poles is developed with the aid of a rigged Hilbert space (Gelfand triplet). A rigged Hilbert space consists of three spaces $X \subset H \subset X'$: a space X of test functions, a Hilbert space H (in our problem, this is the weighted Lebesgue space $L^2(\mathbf{R}, g(\omega)d\omega)$) and the dual space X' of X (a space of continuous linear functionals on X called generalized functions). A suitable choice of X depends on $g(\omega)$. In this paper, two cases are considered: (i) $g(\omega)$ is analytic and decays faster than any exponential functions (e.g. Gaussian distribution), (ii) $g(\omega)$ is a rational function (e.g. Lorentzian distribution $g(\omega) = 1/(\pi(1+\omega^2))$). For the case (i), $X := \mathrm{Exp}_+$ is a space of holomorphic functions $\phi(z)$ defined near the real axis and the upper half plane such that $\sup_{\mathrm{Im}(z)\geq -\varepsilon} |\phi(z)|e^{-\beta|z|}$ is finite for some $\varepsilon > 0$ and $\beta \geq 0$. For the case (ii), $X := H_+$ is a space of bounded holomorphic functions on the real axis and the upper half plane. For both cases, it is proved that the operator T_1 admits the spectral decomposition on X' by regarding the resonance poles as eigenvalues. Corresponding eigenvectors in X' are called the *generalized eigenfunctions* associated with resonance poles. Despite the fact that T_1 is *not* a selfadjoint operator and it has the continuous spectrum, the spectral decomposition of T_1 consists of a countable number of generalized eigenfunctions: roughly speaking, any element ϕ in X is decomposed as $\phi = \sum_{n=0}^{\infty} \mu_n(\phi) \cdot \mu_n$, where $\mu_n \in X'$ is a generalized eigenfunction. Further, it is shown that for the case (ii), the decomposition is reduced to a finite sum because of a certain degeneracy of the space $X = H_+$. We further investigate the semigroup generated by T_1 and the projection to the eigenspace in the space of generalized functions X'. It is proved that the semigroup $e^{T_1 t}$ behaves as $e^{T_1 t}\phi = \sum_{n=0}^{\infty} e^{\lambda_n t}\mu_n(\phi) \cdot \mu_n$ for any $\phi \in X$, where λ_n's are resonance poles. This completely determines the dynamics of the linearized Kuramoto model. In particular, when $0 < K < K_c$, all resonance poles lie on the left half plane: $\mathrm{Re}(\lambda_n) < 0$, which proves the linear stability of the de-synchronous state. When $K = K_c$, there are resonance poles on the imaginary axis. We define a generalized center subspace $\mathbf{E}_c$ on X' to be a space spanned by generalized eigenfunctions associated with resonance poles on the imaginary axis. It is remarkable that though the center subspace in a usual sense is of infinite dimensional because of the continuous spectrum on the imaginary axis, the dimension of the generalized center subspace on X' is finite in general. The projection operator to the generalized center subspace will be investigated in detail.

Note that the spectral decomposition based on a rigged Hilbert space was originally proposed by Gelfand *et al.* [18, 29]. They proposed a spectral decomposition of a selfadjoint operator by using a system of generalized eigenfunctions, however, it involves an integral; that is, eigenfunctions are uncountable. Our results are quite different from Gelfand's one in that our operator T_1 is not selfadjoint and its spectral decomposition consists of a countable eigenfunctions.

Finally, we apply the center manifold reduction to the infinite dimensional Kuramoto model by extending it to an evolution equation on X'. Since the generalized center subspace is of finite dimensional, a corresponding center manifold on X' seems to be a finite dimensional manifold. However, there are no existence theorems of center manifolds on X' because X' is *not* a Banach space. To prove the existence of a center manifold, we introduce a topology on X in a technical way so that the dual space X' becomes a complete metric space. With this topology, X' becomes a topological vector space called Montel space, which is obtained as a projective limit of Banach spaces. This topology proves to have a very convenient property that every weakly convergent series in X' is also convergent with respect the metric. By using this topology and the spectral decomposition, the existence of a finite dimensional center manifold for the Kuramoto model will be proved. The dynamics on the center manifold will be derived when $g(\omega)$ is an even and unimodal function. In this case, the center manifold on X' is of one dimensional, and we can show that the synchronous solution (a solution such that $r > 0$) emerges through the pitchfork bifurcation, which proves the Kuramoto's conjecture. A bifurcation structure depends on the shape of $g(\omega)$. When $g(\omega)$ is an even and bimodal function, it is shown that the synchronous solution appears through a subcritical bifurcation as is shown in Fig.2 (b).

This paper is organized as follows: In Sec.2, the continuous model for the Kuramoto model is introduced and its basic properties are reviewed. In Sec.3, Kuramoto's transition point K_c is derived and it is proved that if $K > K_c$, the de-synchronous state is unstable because of eigenvalues on the right half plane. In Sec.4, the linear stability of the de-synchronous state is investigated. We will show that when $0 < K < K_c$, the order parameter decays exponentially to zero as $t \to \infty$ because of the existence of resonance poles. In Sec.5, the spectral theory of resonance poles on a rigged Hilbert space is developed. We investigate properties of the operator T_1, the semigroup, eigenfunctions, projections by means of the rigged Hilbert space. In Sec.6, the nonlinear stability of the de-synchronous state is proved as an application of the spectral decomposition on the rigged Hilbert space. It is shown that when $0 < K < K_c$, the order parameter tends to zero as $t \to \infty$ without neglecting the nonlinear term. The center manifold theory will be developed in Sec.7. Sec.7.1 to Sec.7.4 are devoted to the proof of the existence of a center manifold on the dual space X'. In Sec.7.5, the dynamics on the center manifold is derived, and the Kuramoto's conjecture is solved.

2 Continuous model

In this section, we introduce a continuous model of the Kuramoto model and show a few properties of it.

For the N-dimensional Kuramoto model (1.1), taking the continuous limit $N \to \infty$, we obtain the continuous model of the Kuramoto model, which is an evolution equation of a probability measure $\rho_t = \rho_t(\theta, \omega)$ on $S^1 = [0, 2\pi)$ parameterized by $t \in \mathbf{R}$ and $\omega \in \mathbf{R}$, defined as

$$\left\{ \begin{array}{l} \dfrac{\partial \rho_t}{\partial t} + \dfrac{\partial}{\partial \theta}\left(\left(\omega + \dfrac{K}{2\sqrt{-1}} (\eta(t) e^{-\sqrt{-1}\theta} - \overline{\eta(t)} e^{\sqrt{-1}\theta}) \right) \rho_t \right) = 0, \\ \eta(t) := \displaystyle\int_{\mathbf{R}} \int_0^{2\pi} e^{\sqrt{-1}\theta} \rho_t(\theta, \omega) g(\omega) d\theta d\omega, \\ \rho_0(\theta, \omega) = h(\theta), \end{array} \right. \tag{2.1}$$

where $h(\theta)$ is an initial condition and $g(\omega)$ is a given probability density function for natural frequencies. We are assuming that the initial condition $h(\theta)$ is independent of ω. This assumption corresponds to the assumption for the discrete model (1.1) that initial values $\{\theta_j(0)\}_{j=1}^N$ and natural frequencies $\{\omega_j\}_{j=1}^N$ are independently distributed, and is a physically natural assumption often used in literature. However, we will also consider ω-dependent initial conditions $h(\theta, \omega)$, a probability measure on S^1 parameterized by ω, for mathematical reasons, in Sec.7. Roughly speaking, $\rho_t(\theta, \omega)$ denotes a probability that an oscillator having a natural frequency ω is placed at a position θ (for example, see [1, 14] for how to derive Eq.(2.1)). Since h and ρ_t are measures on S^1, they should be denoted as $dh(\theta)$ and $d\rho_t(\theta, \cdot)$, however, we use the present notation for simplicity. The $\eta(t)$ is a continuous version of (1.2), and we also call it the *order parameter*. $\overline{\eta(t)}$ denotes the complex conjugate of $\eta(t)$. We can prove that Eq.(2.1) is a proper continuous model in the sense that the order parameter (1.2) of the N-dimensional Kuramoto model converges to $\eta(t)$ as $N \to \infty$ under some assumptions, see Chiba [10]. The purpose in this paper is to investigate the dynamics of Eq.(2.1).

A few properties of Eq.(2.1) are in order. It is easy to prove the low of conservation of mass:

$$\int_{\mathbf{R}}\int_0^{2\pi} \rho_t(\theta, \omega) g(\omega) d\theta d\omega = \int_{\mathbf{R}}\int_0^{2\pi} h(\theta) g(\omega) d\theta d\omega = 1. \tag{2.2}$$

By using the characteristic curve method, Eq.(2.1) is formally integrated as follows: Consider the equation

$$\frac{dx}{dt} = \omega + \frac{K}{2\sqrt{-1}}(\eta(t)e^{-\sqrt{-1}x} - \overline{\eta(t)}e^{\sqrt{-1}x}), \;\; x \in [0, 2\pi), \tag{2.3}$$

which defines a characteristic curve. Let $x = x(t, s; \theta, \omega)$ be a solution of Eq.(2.3) satisfying the initial condition $x(s, s; \theta, \omega) = \theta$ at an initial time s. Then, along the characteristic curve, Eq.(2.1) is integrated to yield

$$\rho_t(\theta, \omega) = h(x(0, t; \theta, \omega)) \exp\Big[\frac{K}{2}\int_0^t (\eta(s)e^{-\sqrt{-1}x(s,t;\theta,\omega)} + \overline{\eta(s)}e^{\sqrt{-1}x(s,t;\theta,\omega)}) ds\Big]. \tag{2.4}$$

By using Eq.(2.4), it is easy to show the equality

$$\int_0^{2\pi} a(\theta, \omega)\rho_t(\theta, \omega) d\theta = \int_0^{2\pi} a(x(t, 0; \theta, \omega), \omega) h(\theta) d\theta, \tag{2.5}$$

for any measurable function $a(\theta, \omega)$. In particular, the order parameter $\eta(t)$ are rewritten as

$$\eta(t) = \int_{\mathbf{R}}\int_0^{2\pi} e^{\sqrt{-1}x(t,0;\theta,\omega)} g(\omega) h(\theta) d\theta d\omega. \tag{2.6}$$

Substituting it into Eqs.(2.3) and (2.4), we obtain

$$\frac{d}{dt}x(t, s; \theta, \omega) = \omega + K\int_{\mathbf{R}}\int_0^{2\pi} \sin\Big(x(t, 0; \theta', \omega') - x(t, s; \theta, \omega)\Big) g(\omega') h(\theta') d\theta' d\omega', \tag{2.7}$$

and

$$\begin{aligned} \rho_t(\theta, \omega) &= h(x(0, t; \theta, \omega)) \times \\ &\exp\Big[K\int_0^t ds \cdot \int_{\mathbf{R}}\int_0^{2\pi} \cos\Big(x(s, 0; \theta', \omega') - x(s, t; \theta, \omega)\Big) h(\theta') g(\omega') d\theta' d\omega'\Big], \end{aligned} \tag{2.8}$$

respectively. They define a system of integro-ordinary differential equations which is equivalent to Eq.(2.1). Even if $h(\theta)$ is not differentiable, we consider Eq.(2.8) to be a weak solution of Eq.(2.1). Indeed, even if h and ρ_t are not differentiable, the quantity (2.5) is differentiable with respect to t when $a(\theta, \omega)$ is differentiable. It is natural to consider the dynamics of weak solutions because ρ_t is a probability measure and we are interested in the dynamics of its moments, in particular the order parameter. In [10], the existence and uniqueness of weak solutions of Eq.(2.1) is proved. In this paper, the existence and uniqueness of solutions of a perturbed system of (2.1), which includes (2.1) itself as a particular case, will be proved in Appendix B. The perturbed system is introduced in Sec.7 to prove the existence of a center manifold.

3 Transition point formula and the linear instability

A trivial solution of the continuous model (2.1), which is independent of θ and t, is given by the uniform distribution $\rho_t(\theta, \omega) = 1/(2\pi)$. In this case, $\eta(t) \equiv 0$. This solution is called the *incoherent state* or the *de-synchronous state*. In this section and the next section, we investigate the linear stability of the de-synchronous state. The nonlinear stability will be discussed in Sec.6.

Let

$$Z_j(t, \omega) := \int_0^{2\pi} e^{\sqrt{-1}j\theta}\rho_t(\theta, \omega)d\theta = \int_0^{2\pi} e^{\sqrt{-1}jx(t,0;\theta,\omega)}h(\theta)d\theta \tag{3.1}$$

be the Fourier coefficients of $\rho_t(\theta, \omega)$. Then, $Z_0(t, \omega) = 1$ and Z_j satisfy the differential equations

$$\frac{dZ_1}{dt} = \sqrt{-1}\omega Z_1 + \frac{K}{2}\eta(t) - \frac{K}{2}\overline{\eta(t)}Z_2, \tag{3.2}$$

and

$$\frac{dZ_j}{dt} = j\sqrt{-1}\omega Z_j + \frac{jK}{2}(\eta(t)Z_{j-1} - \overline{\eta(t)}Z_{j+1}), \tag{3.3}$$

for $j = 2, 3, \cdots$. The order parameter $\eta(t)$ is the integral of $Z_1(t, \omega)$ with the weight $g(\omega)$. The de-synchronous state corresponds to the trivial solution $Z_j \equiv 0$ for $j = 1, 2, \cdots$. Eq.(3.1) shows $|Z_j(t, \omega)| \leq 1$ and thus $Z_j(t, \omega)$ is in the weighted Lebesgue space $L^2(\mathbf{R}, g(\omega)d\omega)$ for every t :

$$||Z_j(t, \cdot)||^2_{L^2(\mathbf{R},g(\omega)d\omega)} = \int_{\mathbf{R}} |Z_j(t, \omega)|^2 g(\omega)d\omega \leq 1.$$

As evolution equations on $L^2(\mathbf{R}, g(\omega)d\omega)$, the above equations are linearized around the origin as

$$\frac{dZ_1}{dt} = \left(\sqrt{-1}\mathcal{M} + \frac{K}{2}\mathcal{P}\right)Z_1, \tag{3.4}$$

and

$$\frac{dZ_j}{dt} = j\sqrt{-1}\mathcal{M}Z_j, \tag{3.5}$$

for $j = 2, 3, \cdots$, where $\mathcal{M} : q(\omega) \mapsto \omega q(\omega)$ is the multiplication operator on $L^2(\mathbf{R}, g(\omega)d\omega)$ and $\mathcal{P}$ is the projection on $L^2(\mathbf{R}, g(\omega)d\omega)$ defined to be

$$\mathcal{P}q(\omega) = \int_{\mathbf{R}} q(\omega)g(\omega)d\omega. \tag{3.6}$$

If we put $P_0(\omega) \equiv 1$, $\mathcal{P}$ is also expressed as $\mathcal{P}q(\omega) = (q, P_0)$, where $(\ ,\ )$ is the inner product on $L^2(\mathbf{R}, g(\omega)d\omega)$ defined as

$$(q_1, q_2) := \int_{\mathbf{R}} q_1(\omega)\overline{q_2(\omega)}g(\omega)d\omega. \tag{3.7}$$

Note that the order parameter is given as $\eta(t) = \mathcal{P}Z_1 = (Z_1, P_0)$. To determine the linear stability of the de-synchronous state and the order parameter, we have to investigate the spectrum and the semigroup of the operator $T_1 := \sqrt{-1}\mathcal{M} + \dfrac{K}{2}\mathcal{P}$.

Remark. We need not assume that the Fourier series $\sum_{-\infty}^{\infty} Z_j(t, \omega)e^{\sqrt{-1}j\theta}$ converges to $\rho_t(\theta, \omega)$ in any sense. It is known that there is a one-to-one correspondence between a measure on S^1 and its Fourier coefficients (see Shohat and Tamarkin [40]). Thus the dynamics of $\{Z_j(t, \omega)\}_{-\infty}^{\infty}$ uniquely determines the dynamics of $\rho_t(\theta, \omega)$, and vice versa. In particular, since a weak solution of the initial value problem (2.1) is unique (Chiba [10]), so is Eqs.(3.2),(3.3).

3.1 Analysis of the operator $\sqrt{-1}\mathcal{M}$

Before investigating the operator T_1, we give a few properties of the multiplication operator $\mathcal{M} : q(\omega) \mapsto \omega q(\omega)$ on $L^2(\mathbf{R}, g(\omega)d\omega)$. The domain $\mathsf{D}(\mathcal{M})$ of $\mathcal{M}$ is dense in $L^2(\mathbf{R}, g(\omega)d\omega)$. It is well known that its spectrum is given by $\sigma(\mathcal{M}) = \mathrm{supp}(g) \subset \mathbf{R}$, where $\mathrm{supp}(g)$ is a support of the function g. Thus the spectrum of $\sqrt{-1}\mathcal{M}$ is

$$\sigma(\sqrt{-1}\mathcal{M}) = \sqrt{-1} \cdot \mathrm{supp}(g) = \{\sqrt{-1}\lambda \,|\, \lambda \in \mathrm{supp}(g)\} \subset \sqrt{-1}\mathbf{R}. \tag{3.8}$$

The semigroup $e^{\sqrt{-1}\mathcal{M}t}$ generated by $\sqrt{-1}\mathcal{M}$ is given as $e^{\sqrt{-1}\mathcal{M}t}q(\omega) = e^{\sqrt{-1}\omega t}q(\omega)$. In particular, we obtain

$$(e^{\sqrt{-1}\mathcal{M}t}q_1, q_2) = \int_{\mathbf{R}} e^{\sqrt{-1}\omega t}q_1(\omega)\overline{q_2(\omega)}g(\omega)d\omega, \tag{3.9}$$

for any $q_1, q_2 \in L^2(\mathbf{R}, g(\omega)d\omega)$. This is the Fourier transform of the function $q_1(\omega)\overline{q_2(\omega)}g(\omega)$. Thus if $q_1(\omega)\overline{q_2(\omega)}g(\omega)$ is real analytic on $\mathbf{R}$ and has an analytic continuation to the upper half plane, then $(e^{\sqrt{-1}\mathcal{M}t}q_1, q_2)$ decays exponentially as $t \to \infty$, while if $q_1(\omega)\overline{q_2(\omega)}g(\omega)$ is C^r, then it decays as $O(1/t^r)$ (see Vilenkin [48]). This means that $e^{\sqrt{-1}\mathcal{M}t}$ does not decay in $L^2(\mathbf{R}, g(\omega)d\omega)$, however, it decays to zero in a suitable weak topology. A weak topology will play an important role in this paper. These facts are summarized as follows:

Proposition 3.1. A solution of the equation (3.5) with an initial value $q(\omega) \in L^2(\mathbf{R}, g(\omega)d\omega)$

is given by $Z_j(t,\omega) = e^{j\sqrt{-1}\mathcal{M}t}q(\omega) = e^{j\sqrt{-1}\omega t}q(\omega)$. The quantity $(e^{j\sqrt{-1}\mathcal{M}t}q_1, q_2)$ decays exponentially to zero as $t \to \infty$ if $g(\omega), q_1(\omega)$ and $q_2(\omega)$ have analytic continuations to the upper half plane.

This proposition suggests that analyticity of $g(\omega)$ and initial conditions also plays an important role for analysis of the operator T_1. The resolvent $(\lambda - \sqrt{-1}\mathcal{M})^{-1}$ of the operator $\sqrt{-1}\mathcal{M}$ is calculated as

$$((\lambda - \sqrt{-1}\mathcal{M})^{-1}q_1, q_2) = \int_{\mathbf{R}} \frac{1}{\lambda - \sqrt{-1}\omega} q_1(\omega)\overline{q_2(\omega)}g(\omega)d\omega. \tag{3.10}$$

We define the function $D(\lambda)$ to be

$$D(\lambda) = ((\lambda - \sqrt{-1}\mathcal{M})^{-1}P_0, P_0) = \int_{\mathbf{R}} \frac{1}{\lambda - \sqrt{-1}\omega} g(\omega)d\omega \tag{3.11}$$

(recall that $P_0(\omega) \equiv 1$). It is holomorphic in $\mathbf{C}\backslash\sigma(\sqrt{-1}\mathcal{M})$ and will be used in later calculations.

3.2 Eigenvalues of the operator T_1 and the transition point formula

The domain of $T_1 = \sqrt{-1}\mathcal{M} + \frac{K}{2}\mathcal{P}$ is given by $\mathsf{D}(\mathcal{M}) \cap \mathsf{D}(\mathcal{P}) = \mathsf{D}(\mathcal{M})$, which is dense in $L^2(\mathbf{R}, g(\omega)d\omega)$. Since $\mathcal{M}$ is selfadjoint and since $\mathcal{P}$ is bounded, T_1 is a closed operator [22]. Let $\varrho(T_1)$ be the resolvent set of T_1 and $\sigma(T_1) = \mathbf{C}\backslash\varrho(T_1)$ the spectrum. Since T_1 is closed, T_1 has no residual spectrum. Let $\sigma_p(T_1)$ and $\sigma_c(T_1)$ be the point spectrum (the set of eigenvalues) and the continuous spectrum of T_1, respectively.

Proposition 3.2. (i) Eigenvalues λ of T_1 are given as roots of

$$D(\lambda) = \frac{2}{K}, \quad \lambda \in \mathbf{C}\backslash\sigma(\sqrt{-1}\mathcal{M}). \tag{3.12}$$

(ii) The continuous spectrum of T_1 is given by

$$\sigma_c(T_1) = \sigma(\sqrt{-1}\mathcal{M}) = \sqrt{-1}\cdot \mathrm{supp}(g). \tag{3.13}$$

Proof. (i) Suppose that $\lambda \in \sigma_p(T_1)\backslash\sigma(\sqrt{-1}\mathcal{M})$. Then, there exists $x \in L^2(\mathbf{R}, g(\omega)d\omega)$ such that

$$\lambda x = (\sqrt{-1}\mathcal{M} + \frac{K}{2}\mathcal{P})x, \quad x \neq 0.$$

Since $\lambda \notin \sigma(\sqrt{-1}\mathcal{M})$, $(\lambda - \sqrt{-1}\mathcal{M})^{-1}$ is defined and the above is rewritten as

$$\begin{aligned} x &= (\lambda - \sqrt{-1}\mathcal{M})^{-1}\frac{K}{2}\mathcal{P}x \\ &= \frac{K}{2}(x, P_0)(\lambda - \sqrt{-1}\mathcal{M})^{-1}P_0(\omega). \end{aligned}$$

By taking the inner product with $P_0(\omega)$, we obtain

$$1 = \frac{K}{2}((\lambda - \sqrt{-1}\mathcal{M})^{-1}P_0, P_0) = \frac{K}{2}D(\lambda). \tag{3.14}$$

This proves that roots of Eq.(3.12) is in $\sigma_p(T_1)\backslash\sigma(\sqrt{-1}\mathcal{M})$. The corresponding eigenvector is given by $x = (\lambda - \sqrt{-1}\mathcal{M})^{-1}P_0(\omega) = 1/(\lambda - \sqrt{-1}\omega)$. If $\lambda \in \sqrt{-1}\mathbf{R}$, $x \notin L^2(\mathbf{R}, g(\omega)d\omega)$. Thus there are no eigenvalues on the imaginary axis. In particular, there are no eigenvalues on $\sigma(\sqrt{-1}\mathcal{M})$.
(ii) This follows from the fact that the essential spectrum is stable under the bounded perturbation (see Kato [22]) : the essential spectrum of T_1 is the same as $\sigma(\sqrt{-1}\mathcal{M})$. Since there are no eigenvalues on $\sigma(\sqrt{-1}\mathcal{M})$, it coincides with the continuous spectrum. ■

Our next task is to calculate roots of Eq.(3.12) to obtain eigenvalues of $T_1 = \sqrt{-1}\mathcal{M} + \frac{K}{2}\mathcal{P}$. By putting $\lambda = x + \sqrt{-1}y$ with $x, y \in \mathbf{R}$, Eq.(3.12) is rewritten as

$$\begin{cases} \displaystyle\int_{\mathbf{R}} \frac{x}{x^2 + (\omega - y)^2} g(\omega)d\omega = \frac{2}{K}, \\ \displaystyle\int_{\mathbf{R}} \frac{\omega - y}{x^2 + (\omega - y)^2} g(\omega)d\omega = 0. \end{cases} \tag{3.15}$$

The next lemma is easily obtained.

Lemma 3.3.
(i) λ satisfies $\mathrm{Re}(\lambda) > 0$ for any $K > 0$ if it exists.
(ii) If $K > 0$ is sufficiently large, there exists at least one eigenvalue λ near infinity.
(iii) If $K > 0$ is sufficiently small, there are no eigenvalues.

Proof. Part (i) of the lemma immediately follows from the first equation of Eq.(3.15): Since the right hand side is positive, x in the left had side has to be positive. To prove part (ii) of the lemma, note that if $|\lambda|$ is large, Eq.(3.12) is expanded as

$$\frac{1}{\lambda} + O(\frac{1}{\lambda^2}) = \frac{2}{K}.$$

Thus the Rouché's theorem proves that Eq.(3.12) has a root $\lambda \sim K/2$ if $K > 0$ is sufficiently large. To prove part (iii) of the lemma, we see that the left hand side of the first equation of Eq.(3.15) is bounded for any $x, y \in \mathbf{R}$. To do so, let $G(\omega)$ be the primitive function of $g(\omega)$ and fix $\delta > 0$ small. The left hand side of the first equation of Eq.(3.15) is calculated as

$$\begin{aligned} & \int_{\mathbf{R}} \frac{xg(\omega)d\omega}{x^2 + (\omega - y)^2} \\ = \ & \int_{y+\delta}^{\infty} \frac{xg(\omega)d\omega}{x^2 + (\omega - y)^2} + \int_{-\infty}^{y-\delta} \frac{xg(\omega)d\omega}{x^2 + (\omega - y)^2} + \int_{y-\delta}^{y+\delta} \frac{xg(\omega)d\omega}{x^2 + (\omega - y)^2} \\ = \ & \int_{y+\delta}^{\infty} \frac{xg(\omega)d\omega}{x^2 + (\omega - y)^2} + \int_{-\infty}^{y-\delta} \frac{xg(\omega)d\omega}{x^2 + (\omega - y)^2} \\ & + \frac{x}{x^2 + \delta^2}(G(y + \delta) - G(y - \delta)) + \int_{y-\delta}^{y+\delta} \frac{2x(\omega - y)}{(x^2 + (\omega - y)^2)^2} G(\omega)d\omega. \end{aligned}$$

The first three terms in the right hand side above are bounded for any $x, y \in \mathbf{R}$. By the mean value theorem, there exists a number ξ such that the last term is estimated as

$$\begin{aligned}
&\int_{y-\delta}^{y+\delta} \frac{2x(\omega - y)}{(x^2 + (\omega - y)^2)^2} G(\omega) d\omega \\
= &\int_0^{\delta} \frac{2x\omega}{(x^2+\omega^2)^2} (G(y+\omega) - G(y-\omega)) d\omega \\
= &(G(y+0) - G(y-0)) \int_0^{\xi} \frac{2x\omega}{(x^2+\omega^2)^2} d\omega + (G(y+\delta) - G(y-\delta)) \int_{\xi}^{\delta} \frac{2x\omega}{(x^2+\omega^2)^2} d\omega.
\end{aligned} \tag{3.16}$$

Since G is continuous, the above is calculated as

$$(G(y+\delta) - G(y-\delta)) \left(\frac{x}{x^2+\xi^2} - \frac{x}{x^2+\delta^2} \right).$$

If $\xi \neq 0$, this is bounded for any $x, y \in \mathbf{R}$. If $\xi = 0$, Eq.(3.16) yields

$$\int_0^{\delta} \frac{2x\omega}{(x^2+\omega^2)^2} (G(y+\omega) - G(y-\omega)) d\omega = (G(y+\delta) - G(y-\delta)) \int_0^{\delta} \frac{2x\omega}{(x^2+\omega^2)^2} d\omega.$$

Since $G(\omega)$ is monotonically increasing, we obtain

$$G(y+\omega) - G(y-\omega) = G(y+\delta) - G(y-\delta)$$

for $0 \leq \omega \leq \delta$. In particular, putting $\omega = 0$ gives $G(y+\delta) - G(y-\delta) = 0$. Thus $G(y+\omega) - G(y-\omega) = 0$ for $0 \leq \omega \leq \delta$. This proves that the quantity (3.16) is zero. Now we have proved that the left hand side of the first equation of Eq.(3.15) is bounded for any $x, y \in \mathbf{R}$, although the right hand side diverges as $K \to +0$. Thus Eq.(3.12) has no roots if $K > 0$ is sufficiently small. ■

Lemma 3.3 shows that if $K > 0$ is sufficiently large, the trivial solution $Z_1 = 0$ of the equation $dZ_1/dt = T_1 Z_1$ is unstable because of eigenvalues with positive real parts. Our purpose in this section is to determine the bifurcation point K_c such that if $K < K_c$, the operator T_1 has no eigenvalues, while if K exceeds K_c, eigenvalues appear on the right half plane (K_c should be positive because of Lemma 3.3 (iii)). To calculate eigenvalues $\lambda = \lambda(K)$ explicitly is difficult in general. However, since zeros of a holomorphic function do not vanish because of the argument principle, $\lambda(K)$ disappears if and only if it is absorbed into the continuous spectrum $\sigma(\sqrt{-1}\mathcal{M})$, on which $D(\lambda)$ is not holomorphic, as K decreases. This fact suggests that to determine K_c, it is sufficient to investigate Eq.(3.12) or Eq.(3.15) near the imaginary axis. Thus take the limit $x \to +0$ in Eq.(3.15):

$$\left\{ \begin{array}{l}
\displaystyle \lim_{x \to +0} \int_{\mathbf{R}} \frac{x}{x^2 + (\omega - y)^2} g(\omega) d\omega = \frac{2}{K}, \\
\displaystyle \lim_{x \to +0} \int_{\mathbf{R}} \frac{\omega - y}{x^2 + (\omega - y)^2} g(\omega) d\omega = 0.
\end{array} \right. \tag{3.17}$$



Fig. 3: A schematic view of behavior of roots λ of Eq.(3.12) when K decreases. Thick lines denote the continuous spectrum. As K decreases, eigenvalues $\lambda_1, \lambda_2, \cdots$ converge to $\sqrt{-1}y_1, \sqrt{-1}y_2, \cdots$ and disappear at some $K = K_1, K_2, \cdots$, respectively.

These equations determine K_j and y_j such that one of the eigenvalues $\lambda = \lambda_j(K)$ converges to $\sqrt{-1}y_j$ as $K \to K_j + 0$ (see Fig.3). To calculate them, we need the next lemma.

In this paper, the probability density function $g(\omega)$ is called piecewise continuous when it satisfies following: If $g(\omega)$ is discontinuous at ω_0, then $g(\omega_0 + 0)$ and $g(\omega_0 - 0)$ exist, and for any $\varepsilon > 0$, there exists $\delta > 0$ such that when $-\delta < \omega - \omega_0 < 0$, $|g(\omega) - g(\omega_0 - 0)| < \varepsilon$ and when $0 < \omega - \omega_0 < \delta$, $|g(\omega) - g(\omega_0 + 0)| < \varepsilon$. If $g(\omega)$ is piecewise continuous except for a measure zero set, then we also call it piecewise continuous. Note that when $g(\omega_0 - 0) = g(\omega_0 + 0) \neq g(\omega_0)$, we can modify $g(\omega)$ on a point $\{\omega_0\}$ so that it becomes continuous at ω_0. Thus if g is discontinuous at ω_0, we can assume without loss of generality that $g(\omega_0 - 0) \neq g(\omega_0 + 0)$.

Lemma 3.4. (i) Suppose that $g(\omega)$ is piecewise continuous. If one of the eigenvalues $\lambda(K)$ satisfies $\lambda(K) \to \sqrt{-1}y_j$ as $K \to K_j$. Then, $g(\omega)$ is continuous at $\omega = y_j$.
(ii) If $g(\omega)$ is continuous at $\omega = y$, then

$$\lim_{x \to +0} \int_{\mathbf{R}} \frac{x}{x^2 + (\omega - y)^2} g(\omega) d\omega = \pi g(y). \tag{3.18}$$

Proof. To prove (i), suppose that $g(\omega)$ is discontinuous at $\omega = 0$ without loss of generality. We will show that a root $\lambda(K)$ of Eq.(3.12) can not approach to zero for any $K > 0$. Put $g(+0) = h_+$, $g(-0) = h_-$ and $h_+ \neq h_-$. For any $\varepsilon > 0$, there exists $\delta > 0$ such that if $-\delta < \omega < 0$, then $|g(\omega) - h_-| < \varepsilon$ and if $0 < \omega < \delta$, then $|g(\omega) - h_+| < \varepsilon$. For Eq.(3.12), we suppose $|\lambda| = |x + \sqrt{-1}y| < \delta$ and $y > 0$. The case $y < 0$ is treated in a similar manner.

We calculate $D(\lambda)$ as

$$\begin{aligned} D(\lambda) &= \int_\delta^\infty \frac{g(\omega)}{\lambda - \sqrt{-1}\omega} d\omega + \int_{-\infty}^{-\delta} \frac{g(\omega)}{\lambda - \sqrt{-1}\omega} d\omega \\ &\quad + \int_0^\delta \frac{1}{\lambda - \sqrt{-1}\omega} (g(\omega) - h_+ + h_+)\, d\omega + \int_{-\delta}^0 \frac{1}{\lambda - \sqrt{-1}\omega} (g(\omega) - h_- + h_-)\, d\omega \\ &= \int_\delta^\infty \frac{g(\omega)}{\lambda - \sqrt{-1}\omega} d\omega + \int_\delta^\infty \frac{g(-\omega)}{\lambda + \sqrt{-1}\omega} d\omega \\ &\quad + \int_0^\delta \frac{1}{\lambda - \sqrt{-1}\omega} (g(\omega) - h_+)\, d\omega + \int_0^\delta \frac{1}{\lambda + \sqrt{-1}\omega} (g(-\omega) - h_-)\, d\omega \\ &\quad + \int_0^\delta \frac{1}{\lambda - \sqrt{-1}\omega} (h_+ - h_-)\, d\omega + h_- \int_0^\delta \frac{d\omega}{\lambda - \sqrt{-1}\omega} + h_- \int_0^\delta \frac{d\omega}{\lambda + \sqrt{-1}\omega}. \end{aligned} \tag{3.19}$$

Since $|\lambda| < \delta$, there exists a positive number M, which is independent of λ, such that

$$\int_\delta^\infty \frac{g(\pm\omega)}{\lambda \mp \sqrt{-1}\omega} d\omega < M.$$

Thus $|D(\lambda)|$ is estimated as

$$\begin{aligned} |D(\lambda)| &\geq |h_+ - h_-| \int_0^\delta \frac{d\omega}{|\lambda - \sqrt{-1}\omega|} - \varepsilon \int_0^\delta \frac{d\omega}{|\lambda - \sqrt{-1}\omega|} - \varepsilon \int_0^\delta \frac{d\omega}{|\lambda + \sqrt{-1}\omega|} \\ &\quad - \left| h_- \log\left(\frac{\lambda + \sqrt{-1}\delta}{\lambda - \sqrt{-1}\delta} \right) \right| - 2M. \end{aligned}$$

Since $y > 0$, $|\lambda - \sqrt{-1}\omega| \leq |\lambda + \sqrt{-1}\omega|$ when $\omega \geq 0$. This shows that

$$|D(\lambda)| \geq (|h_+ - h_-| - 2\varepsilon) \int_0^\delta \frac{d\omega}{|\lambda - \sqrt{-1}\omega|} - \left| h_- \log\left(\frac{\lambda + \sqrt{-1}\delta}{\lambda - \sqrt{-1}\delta} \right) \right| - 2M. \tag{3.20}$$

The right hand side tends to infinity as $\lambda \to 0$ if $2\varepsilon < |h_+ - h_-|$. This proves that as K decreases, a root of Eq.(3.12) can not approach to zero for positive K.

(ii) The formula Eq.(3.18) is famous and given in Ahlfors [3]. ■

Recall that the second equation of Eq.(3.17) determines imaginary parts to which $\lambda(K)$ converges as $\mathrm{Re}(\lambda(K)) \to +0$. Let $y_1, y_2, \cdots$ be roots of the second equation of Eq.(3.17), at which $g(\omega)$ is continuous. Since $g(\omega)$ is continuous at $\omega = y_j$, substituting it into the first equation of Eq.(3.17) yields

$$K_j = \frac{2}{\pi g(y_j)}, \quad j = 1, 2, \cdots, \tag{3.21}$$

which gives the value such that $\mathrm{Re}(\lambda(K)) \to 0$ as $K \to K_j + 0$. Now we obtain the next theorem.

Theorem 3.5. Let $y_1, y_2, \cdots$ be roots of the second equation of Eq.(3.17). Suppose that g is continuous at $y_1, y_2, \cdots$ (this is true when g is piecewise continuous). Put

$$K_c := \inf_j K_j = \frac{2}{\pi \sup_j g(y_j)}. \tag{3.22}$$

If $0 < K \leq K_c$, the operator T_1 has no eigenvalues, while if K exceeds K_c, eigenvalues of T_1 appear on the right half plane. In this case, the trivial solution $Z_1 = 0$ of Eq.(3.4) is unstable.

In general, there exists $K_c^{(2)}$ such that T_1 has eigenvalues when $K_c < K < K_c^{(2)}$ but they disappear again at $K = K_c^{(2)}$; i.e. the stability of the trivial solution $Z_1 = 0$ may change many times (see Example 3.9). Such $K_c^{(2)}$ is one of the values K_j's. However, if $g(\omega)$ is an even and unimodal function, it is easy to prove that T_1 has eigenvalues on the right half plane for any $K > K_c$, and they are real as is shown in Mirollo and Strogatz [32]. Indeed, the second equation of Eq.(3.15) is calculated as

$$0 = \int_{\mathbf{R}} \frac{\omega - y}{x^2 + (\omega - y)^2} g(\omega) d\omega = \int_0^\infty \frac{\omega}{x^2 + \omega^2} (g(y+\omega) - g(y-\omega)) d\omega.$$

If g is even, $y = 0$ is a root of this equation. If g is unimodal, $g(y+\omega) - g(y-\omega) > 0$ when $y < 0, \omega > 0$ and $g(y+\omega) - g(y-\omega) < 0$ when $y > 0, \omega > 0$. Hence, $y = 0$ is a unique root. This implies that an eigenvalue should be on the real axis, and $(K, y) = (K_c, 0)$ is a unique solution of Eq.(3.17). As a corollary, we obtain the transition point (bifurcation point to the synchronous state) conjectured by Kuramoto [26]:

Corollary 3.6 (Kuramoto's transition point). Suppose that the probability density function $g(\omega)$ is even, unimodal and continuous at $y = 0$. Then, K_c defined as above is given by

$$K_c = \frac{2}{\pi g(0)}. \tag{3.23}$$

When $K > K_c$, the solution $Z_1 = 0$ of Eq.(3.4) is unstable. In particular, the order parameter $\eta(t) = (Z_1, P_0)$ is linearly unstable.

Example 3.7. If $g(\omega)$ is given by

$$g(\omega) = \begin{cases} 1 & (-1 \leq \omega \leq -1/2) \\ 1/2 & (1/4 \leq \omega \leq 5/4) \\ 0 & (\text{otherwise}), \end{cases} \tag{3.24}$$

(see Fig.4 (a)), then roots of the second equation of Eq.(3.17) are $y \sim -0.706, -0.044, 0.567$. Thus K_c is given as

$$K_c = \frac{2}{\pi g(-0.706)} = \frac{2}{\pi}.$$



Fig. 4: Probability density functions $g(\omega)$ for (a) Example 3.7 and (b) Example 3.8.

More generally, suppose that $g(\omega)$ is of the form

$$g(\omega) = \begin{cases} c_1 & (a_1 \leq \omega \leq b_1) \\ c_2 & (a_2 \leq \omega \leq b_2) \\ \vdots & \qquad \vdots \\ c_n & (a_n \leq \omega \leq b_n) \\ 0 & \text{(otherwise)}, \end{cases} \tag{3.25}$$

where $c_i \neq 0$ and $a_1 < b_1 < a_2 < \cdots < b_n$ are constants. In this case, we can prove that the second equation of Eq.(3.17) has roots y_j on the intervals (a_j, b_j) for $j = 1, \cdots, n$. Thus K_c is given by

$$K_c = \frac{2}{\pi \max_{1\leq j\leq n} c_j}. \tag{3.26}$$

Indeed, the left hand side of the second equation of Eq.(3.17) is calculated as

$$\lim_{x\to 0} \int_{\mathbf{R}} \frac{\omega - y}{x^2 + (\omega - y)^2} g(\omega) d\omega = \sum_{j=1}^{n} c_j \log\left(\frac{y - a_j}{b_j - y}\right).$$

This tends to $-\infty$ as $y \to a_j + 0$ and to $+\infty$ as $y \to b_j - 0$. Now the intermediate-value theorem proves the desired result.

Example 3.8. If $g(\omega)$ is given as Fig.4 (b), the second equation of Eq.(3.17) has roots $y = 0$ and $y \sim \pm 0.7459$. Thus K_c is approximately given by

$$K_c = \frac{2}{\pi g(0.7459)} = \frac{2}{\pi \cdot 1.9671}. \tag{3.27}$$

It is notable that the value 0.7459 is slightly different from 0.75 which gives the maximum of $g(\omega)$. This result gives a counter-example for the Kuramoto's argument $K_c = 2/(\pi \max g(\omega))$ for bimodal distributions [26]. Such a counter-example is also given by Martens *et al.* [30].

Example 3.9. We give an example that the stability of $Z_1 = 0$ changes many times.

Suppose that $g(\omega)$ is given by

$$g(\omega) = \begin{cases} 7 & (-3 < \omega < -2) \\ -0.1\omega + 3.8 & (-2 < \omega < -1) \\ 3 & (-1 < \omega < 1) \\ 0.1\omega + 3.8 & (1 < \omega < 2) \\ 7 & (2 < \omega < 3) \\ 0 & (\text{otherwise}), \end{cases} \tag{3.28}$$

see Fig.5 (a). Behavior of eigenvalues $\lambda(K)$ is represented as Fig.5 (b), in which arrows indicate the direction that K increases. Solutions of the second equation of Eq.(3.17) are approximately given by

$$y_{\pm 1} = \pm 2.1032,\ y_{\pm 2} = \pm 1.6882,\ y_{\pm 3} = \pm 1.4046,\ y_4 = 0. \tag{3.29}$$

This means that $Z_1 = 0$ is unstable if and only if

$$0.0909 \sim \frac{2}{\pi g(y_1)} = K_c < K < K_c^{(2)} = \frac{2}{\pi g(y_2)} \sim 0.1604 \tag{3.30}$$

and

$$0.1616 \sim \frac{2}{\pi g(y_3)} < K < \infty. \tag{3.31}$$

Thus $Z_1 = 0$ recovers the stability in the small interval $0.1604 < K < 0.1616$. Indeed, an eigenvalue emerges at $\sqrt{-1}y_1$ when $K = K_c$, it goes to the right side as K increases for a moment, but it again approaches the imaginary axis and disappears at $\sqrt{-1}y_2$ when $K = K_c^{(2)}$. After that, another eigenvalue emerges at $\sqrt{-1}y_3$ when $K = K_3$. This eigenvalue remains on the right half plane for any $K > K_3$ (by using the theory of resonance poles developed in Sec.4, we can show that the eigenvalue disappearing at $\sqrt{-1}y_2$ is connected to the eigenvalue emerging from $\sqrt{-1}y_3$ in the second Riemann sheet of the resolvent of T_1, see Fig.6).

Remark. Although we suppose in this paper that the distribution $g(\omega)$ for natural frequencies is a probability density function, it is interesting to consider a more general case that natural frequencies are distributed according to a probability measure $\hat{g}$, not a function. In this case, we can prove that $K_c = 0$ if and only if there is a point $y \in \mathbf{R}$ such that $\hat{g}(\{y\}) \neq 0$. The proof is done in a similar way to that of Lemma 3.3 (iii), in which $g(\omega)d\omega$ and $G(\omega)$ are replaced by $d\hat{g}$ and $\hat{g}((-\infty, \omega])$, respectively. If $\hat{g}(\{y\}) \neq 0$, then $G(y + 0) \neq G(y - 0)$. As a result, the first term in the right hand side of Eq.(3.16) is estimated as $O(1/x)$ as $x \to +0$. This and Rouché's theorem prove that Eq.(3.12) has a root λ for any small $K > 0$ satisfying $\mathrm{Re}(\lambda) \sim O(K)$ as $K \to +0$. Thus the operator T_1 has eigenvalues on the right half plane for any $K > 0$. This means that if a lot of oscillators share the same natural frequency, they may synchronize for any small $K > 0$.



Fig. 5: (a) A probability density function $g(\omega)$ for Example 3.9. (b) A schematic view of behavior of eigenvalues $\lambda(K)$ for Example 3.9.

4 Linear stability theory

Theorem 3.5 shows that K_c is the least bifurcation point and the trivial solution $Z_1 = 0$ of Eq.(3.4) is unstable if K is slightly larger than K_c. If $0 < K \leq K_c$, there are no eigenvalues and the continuous spectrum of T_1 lies on the imaginary axis: $\sigma(T_1) = \sigma(\sqrt{-1}\mathcal{M})$. In this section, we investigate the dynamics of Eq.(3.4) for $0 < K < K_c$. We will see that the order parameter $\eta(t)$ may decay exponentially even if the spectrum lies on the imaginary axis because of the existence of resonance poles.

4.1 Resonance poles

Since $\sqrt{-1}\mathcal{M}$ has the semigroup $e^{\sqrt{-1}\mathcal{M}t}$ and since $\mathcal{P}$ is bounded, the operator $T_1 = \sqrt{-1}\mathcal{M} + \frac{K}{2}\mathcal{P}$ also generates the semigroup $e^{T_1 t}$ (Kato [22]) on $L^2(\mathbf{R}, g(\omega)d\omega)$. A solution of Eq.(3.4) with an initial value $\phi(\omega) \in L^2(\mathbf{R}, g(\omega)d\omega)$ is given by $e^{T_1 t}\phi(\omega)$. The semigroup $e^{T_1 t}$ is calculated by using the Laplace inversion formula

$$e^{T_1 t} = \lim_{y\to\infty} \frac{1}{2\pi\sqrt{-1}} \int_{x-\sqrt{-1}y}^{x+\sqrt{-1}y} e^{\lambda t}(\lambda - T_1)^{-1} d\lambda, \tag{4.1}$$

for $t > 0$, where $x > 0$ is chosen so that the contour (see Fig.7 (a)) is to the right of the spectrum of T_1 (Hille and Phillips [21], Yosida [49]). The resolvent $(\lambda - T_1)^{-1}$ is given as follows.

Lemma 4.1. For any $\phi(\omega), \psi(\omega) \in L^2(\mathbf{R}, g(\omega)d\omega)$, the equality

$$\begin{aligned} &((\lambda - T_1)^{-1}\phi, \psi)\\ &= ((\lambda - \sqrt{-1}\mathcal{M})^{-1}\phi, \psi) + \frac{K/2}{1 - KD(\lambda)/2}((\lambda - \sqrt{-1}\mathcal{M})^{-1}\phi, P_0)((\lambda - \sqrt{-1}\mathcal{M})^{-1}P_0, \psi) \end{aligned} \tag{4.2}$$

holds.

Proof. Put

$$R(\lambda)\phi := (\lambda - T_1)^{-1}\phi = (\lambda - \sqrt{-1}\mathcal{M} - \frac{K}{2}\mathcal{P})^{-1}\phi,$$

which yields

$$\begin{aligned}(\lambda - \sqrt{-1}\mathcal{M})R(\lambda)\phi &= \phi + \frac{K}{2}\mathcal{P}R(\lambda)\phi \\ &= \phi + \frac{K}{2}(R(\lambda)\phi, P_0)P_0.\end{aligned}$$

This is rearranged as

$$R(\lambda)\phi = (\lambda - \sqrt{-1}\mathcal{M})^{-1}\phi + \frac{K}{2}(R(\lambda)\phi, P_0)(\lambda - \sqrt{-1}\mathcal{M})^{-1}P_0. \tag{4.3}$$

By taking the inner product with P_0, we obtain

$$(R(\lambda)\phi, P_0) = ((\lambda - \sqrt{-1}\mathcal{M})^{-1}\phi, P_0) + \frac{K}{2}(R(\lambda)\phi, P_0)D(\lambda).$$

This provides

$$(R(\lambda)\phi, P_0) = \frac{1}{1 - KD(\lambda)/2}((\lambda - \sqrt{-1}\mathcal{M})^{-1}\phi, P_0).$$

Substituting it into Eq.(4.3), we obtain Lemma 4.1. ■

Eq.(4.1) and Lemma 4.1 show that $(e^{T_1 t}\phi, \psi)$ is given by

$$\begin{aligned}(e^{T_1 t}\phi, \psi) &= \lim_{y\to\infty} \frac{1}{2\pi\sqrt{-1}} \int_{x-\sqrt{-1}y}^{x+\sqrt{-1}y} e^{\lambda t}\Big((\lambda - \sqrt{-1}\mathcal{M})^{-1}\phi, \psi) \\ &\quad + \frac{K/2}{1 - KD(\lambda)/2}((\lambda - \sqrt{-1}\mathcal{M})^{-1}\phi, P_0)((\lambda - \sqrt{-1}\mathcal{M})^{-1}P_0, \psi)\Big)d\lambda. \end{aligned} \tag{4.4}$$

In particular, the order parameter $\eta(t) = (Z_1, P_0)$ for the linearized system (3.4) with the initial condition $Z_1(0, \omega) = \phi(\omega)$ is given by $\eta(t) = (e^{T_1 t}\phi, P_0)$.

One of the effective ways to calculate the integral above is to use the residue theorem. Recall that the resolvent $(\lambda - T_1)^{-1}$ is holomorphic on $\mathbf{C}\backslash\sigma(T_1)$. When $0 < K \leq K_c$, T_1 has no eigenvalues and the continuous spectrum lies on the imaginary axis : $\sigma(T_1) = \sigma(\sqrt{-1}\mathcal{M}) = \sqrt{-1}\cdot \mathrm{supp}(g)$. Thus the integrand $e^{\lambda t}((\lambda - T_1)^{-1}\phi, \psi)$ in Eq.(4.4) is holomorphic on the right half plane and may not be holomorphic on $\sigma(T_1)$. However, under assumptions below, we can show that the integrand has an analytic continuation through the line $\sigma(T_1)$ from right to left. Then, the analytic continuation may have poles on the left half plane (on the second Riemann sheet of the resolvent), which are called *resonance poles* [38]. The resonance pole λ affects the integral in Eq.(4.4) through the residue theorem (see Fig.7 (b)). In this manner, the order parameter $\eta(t)$ can decay with the exponential rate $e^{\mathrm{Re}(\lambda)t}$. Such an exponential decay caused by resonance poles is well known in the theory of Schrödinger operators [38], and for the Kuramoto model, it is investigated numerically by Strogatz *et al.* [44] and Balmforth *et al.* [4].

At first, we construct an analytic continuation of the function $F_0(\lambda) := ((\lambda - T_1)^{-1}\phi, \overline{\psi})$. In the next lemma, we consider $((\lambda - T_1)^{-1}\phi, \overline{\psi})$ instead of $((\lambda - T_1)^{-1}\phi, \psi)$ so that the complex conjugate of $\psi(\omega)$ does not appear in the right hand side of Eq.(4.5) (by the definition of the inner product, $(\,\cdot\,, \psi)$ involves the conjugate of ψ, while $(\,\cdot\,, \overline{\psi})$ does not).

Lemma 4.2. Suppose that the probability density function $g(\omega)$ and functions $\phi(\omega), \psi(\omega)$ are real analytic on $\mathbf{R}$ and they have meromorphic continuations to the upper half plane. Then the function $F_0(\lambda) := ((\lambda - T_1)^{-1}\phi, \overline{\psi})$ defined on the right half plane has the meromorphic continuation $F_1(\lambda)$ to the left half plane, which is given by

$$\begin{aligned} F_1(\lambda) &= ((\lambda - \sqrt{-1}\mathcal{M})^{-1}\phi, \overline{\psi}) + 2\pi\phi(-\sqrt{-1}\lambda)\psi(-\sqrt{-1}\lambda)g(-\sqrt{-1}\lambda) \\ &\qquad + \frac{K/2}{1 - KD(\lambda)/2 - \pi K g(-\sqrt{-1}\lambda)} Q[\lambda, \phi] Q[\lambda, \psi], \end{aligned} \tag{4.5}$$

where $Q[\lambda, \phi]$ is defined to be

$$Q[\lambda, \phi] = ((\lambda - \sqrt{-1}\mathcal{M})^{-1}\phi, P_0) + 2\pi g(-\sqrt{-1}\lambda)\phi(-\sqrt{-1}\lambda). \tag{4.6}$$

Note that $Q[\lambda, \,\cdot\,]$ defines a linear functional for each $\lambda \in \mathbf{C}$. Actually, we will define a suitable function space in Sec.5 so that $Q[\lambda, \,\cdot\,]$ becomes a continuous linear functional (generalized function).

Proof. Define a function $\widetilde{F}(\lambda)$ to be

$$\widetilde{F}(\lambda) = \begin{cases} ((\lambda - \sqrt{-1}\mathcal{M})^{-1}\phi, \overline{\psi}) & (\mathrm{Re}(\lambda) > 0), \\ ((\lambda - \sqrt{-1}\mathcal{M})^{-1}\phi, \overline{\psi}) + 2\pi\phi(-\sqrt{-1}\lambda)\psi(-\sqrt{-1}\lambda)g(-\sqrt{-1}\lambda) & (\mathrm{Re}(\lambda) < 0). \end{cases} \tag{4.7}$$

By the formula (3.18), we obtain

$$\begin{aligned} &\lim_{\mathrm{Re}(\lambda)\to+0} ((\lambda - \sqrt{-1}\mathcal{M})^{-1}\phi, \overline{\psi}) - \lim_{\mathrm{Re}(\lambda)\to-0} ((\lambda - \sqrt{-1}\mathcal{M})^{-1}\phi, \overline{\psi}) \\ &= 2\pi\phi(\mathrm{Im}(\lambda)) \cdot \psi(\mathrm{Im}(\lambda)) \cdot g(\mathrm{Im}(\lambda)), \end{aligned} \tag{4.8}$$

which proves that $\lim_{\mathrm{Re}(\lambda)\to+0} \widetilde{F}(\lambda) = \lim_{\mathrm{Re}(\lambda)\to-0} \widetilde{F}(\lambda)$. Therefore, if we show that $\widetilde{F}(\lambda)$ is continuous on the imaginary axis, then $\widetilde{F}(\lambda)$ is meromorphic on $\mathbf{C}$ by the Schwarz's principle of reflection. To see this, put $\phi(\omega)\psi(\omega)g(\omega) = q(\omega)$. By the formula (3.18),

$$\begin{aligned} &\lim_{x\to+0} \int_{-\infty}^{\infty} \frac{1}{\lambda - \sqrt{-1}\omega} q(\omega) d\omega \\ = \;& \lim_{x\to+0} \int_{-\infty}^{\infty} \frac{x}{x^2 + (\omega - y)^2} q(\omega) d\omega + \sqrt{-1} \lim_{x\to+0} \int_{-\infty}^{\infty} \frac{\omega - y}{x^2 + (\omega - y)^2} q(\omega) d\omega \\ = \;& \pi q(y) - \pi\sqrt{-1} V(y), \end{aligned}$$

where $\lambda = x + \sqrt{-1}y$ and $V(y)$ is the Hilbert transform of q defined by

$$V(y) = \text{p.v.}\frac{1}{\pi}\int_{-\infty}^{\infty}\frac{1}{t}q(y-t)dt, \tag{4.9}$$

see Chap.VI of Stein and Weiss [41]. Since $q(y)$ is Lipschitz continuous, so is $V(y)$ (Thm.106 of Titchmarsh [45]). This proves that $\lim_{x\to+0}\int_{-\infty}^{\infty}(\lambda - \sqrt{-1}\omega)^{-1}q(\omega)d\omega$ is continuous in y. Therefore, $\widetilde{F}(\lambda)$ is meromorphic on $\mathbf{C}$. Now we have obtained the meromorphic continuation of $((\lambda - \sqrt{-1}\mathcal{M})^{-1}\phi, \overline{\psi})$ from right to left. Applying this to Eq.(4.2), we obtain the meromorphic continuation of $F_0(\lambda)$ as Eq.(4.5). ■

Eq.(4.5) is rewritten as

$$\begin{aligned}
F_1(\lambda) \;=\; & \frac{K/2}{1 - KD(\lambda)/2 - \pi K g(-\sqrt{-1}\lambda)}\Bigg(\\
& (2/K - D(\lambda))((\lambda - \sqrt{-1}\mathcal{M})^{-1}\phi, \overline{\psi}) + ((\lambda - \sqrt{-1}\mathcal{M})^{-1}\phi, P_0)\cdot((\lambda - \sqrt{-1}\mathcal{M})^{-1}\psi, P_0) \\
& + \; 2\pi g(-\sqrt{-1}\lambda)\Big(\frac{2}{K}\phi(-\sqrt{-1}\lambda)\psi(-\sqrt{-1}\lambda) - D(\lambda)\phi(-\sqrt{-1}\lambda)\psi(-\sqrt{-1}\lambda) \\
& \qquad - ((\lambda - \sqrt{-1}\mathcal{M})^{-1}\phi, \overline{\psi}) + ((\lambda - \sqrt{-1}\mathcal{M})^{-1}\phi, P_0)\cdot\psi(-\sqrt{-1}\lambda) \\
& \qquad + ((\lambda - \sqrt{-1}\mathcal{M})^{-1}\psi, P_0)\cdot\phi(-\sqrt{-1}\lambda)\Big)\Bigg).
\end{aligned} \tag{4.10}$$

This expression shows that poles of g are removable. Therefore, poles of $F_1(\lambda)$ on the left half plane and the imaginary axis are given as roots of the equation

$$\left\{\begin{array}{ll} D(\lambda) + 2\pi g(-\sqrt{-1}\lambda) = \dfrac{2}{K}, & \text{Re}(\lambda) < 0, \\ \lim\limits_{\text{Re}(\lambda)\to+0} D(\lambda) = \lim\limits_{\text{Re}(\lambda)\to-0}\Big(D(\lambda) + 2\pi g(-\sqrt{-1}\lambda)\Big) = \dfrac{2}{K}, & \text{Re}(\lambda) = 0, \end{array}\right. \tag{4.11}$$

and poles of the functions $\phi(-\sqrt{-1}\lambda)$ and $\psi(-\sqrt{-1}\lambda)$. To avoid dynamics caused by a special choice of ϕ and ψ, in what follows, we will assume that continuations of ϕ and ψ have no poles.

Definition 4.3. Roots of Eq.(4.11) on the left half plane and the imaginary axis are called *resonance poles* of the operator T_1.

Since the left hand side of Eq.(4.11) is an analytic continuation of that of Eq.(3.12), at least one of the resonance poles is obtained as a continuation of an eigenvalue $\lambda(K)$ coming from the right half plane when K decreases from K_c (see Fig. 6 and a comment in Example 3.9). However, if $g(\lambda)$ has an essential singularity, there exist infinitely many resonance poles in general, which are not obtained as continuations of eigenvalues.

We want to calculate the Laplace inversion formula (4.4) by deforming the contour as Fig.7 (b), and pick up the residues at resonance poles. We should show that the integral along the arc C_4 converges to zero as the radius tends to infinity. For this purpose, we



Fig. 6: As is discussed in Sec.3, an eigenvalue $\lambda(K)$ disappears from the original complex plane at $K = K_c$. But it still exists as a resonance pole on the second Riemann sheet of the resolvent.

have to make some assumptions for growth rates of $\phi(\lambda)$ and $\psi(\lambda)$ as $|\lambda| \to \infty$. Since suitable assumptions depend on the growth rate of $g(\lambda)$, we calculate the Laplace inversion formula by dividing into two cases: In Sec.4.2, $g(\omega)$ is assumed to decay faster than any exponential functions, such as the Gaussian distribution. In Sec.4.3, we consider the case that $g(\omega)$ is a rational function.

Remark. In Lemma 4.2 and Def.4.3, $g(\omega)$ is assumed to be analytic on the real axis. If $g(\omega)$ is piecewise analytic on the real axis such as Examples 3.7 to 3.9, we can show that an analytic continuation of the resolvent has logarithmic singularities: if $g(\omega)$ is not analytic at $\omega = y \in \mathbf{R}$, then $\sqrt{-1}y$ is a logarithmic singularity of $F_0(\lambda)$. In this case, there exist infinitely many Riemann sheets and resonance poles exist on every Riemann sheets in general. Resonance poles on the n-th Riemann sheet are given as roots of the equation

$$D(\lambda) + 2\pi n g(-\sqrt{-1}\lambda) = \frac{2}{K}, \quad \mathrm{Re}(\lambda) < 0.$$

In this paper, we do not deal with such problems.

4.2 Gaussian-type case

In this subsection, we suppose that the probability density function $g(\omega)$ for natural frequencies satisfies that

(A1) $g(\omega)$ is real analytic on $\mathbf{R}$ and has a meromorphic continuation $g(\lambda)$ to the upper half plane.
(A2) for any $\gamma > 0$, there exists a positive number C such that

$$|g(\omega)| < Ce^{-\gamma|\omega|}, \quad \omega \in \mathbf{R}. \tag{4.12}$$

(A3) there exists a positive number δ such that in the angular domains

$$|\arg(\lambda)| \leq \delta, \ \ |\arg(\lambda) - \pi| \leq \delta, \tag{4.13}$$

near the real axis, $g(\lambda)$ satisfies Eq.(4.12) for any $\gamma > 0$.

Note that since $g(\omega)$ is real-valued on $\mathbf{R}$, (A1) implies that $g(\omega)$ also has a meromorphic continuation to the lower half plane due to the Schwarz's principle of reflection. The assumption (A3) implies that there are no poles of g in the angular domains (4.13). These assumptions are satisfied, for example, when g is of the form $g(\omega) = C' e^{P(\omega)}$ with a polynomial $P(\omega) = -\omega^{2m} + a_1\omega^{2m-1} + \cdots + a_{2m}$ and some constant C', such as the Gaussian distribution. Obviously (A3) includes (A2), however, they are divided to emphasize that (A3) is an assumption for an analytic continuation of g. The assumption (A2) holds while the assumption (A3) fails, for instance, if $g(\omega)$ decays as $\exp(-\exp\omega^2)$ on $\mathbf{R}$.

To make assumptions for ϕ and ψ, we prepare a certain function space. Let $\mathrm{Exp}_+(\beta, n)$ be the set of holomorphic functions on the region $\mathbf{C}_n := \{z \in \mathbf{C} \,|\, \mathrm{Im}(z) \geq -1/n\}$ such that the norm

$$||\phi||_{\beta,n} := \sup_{\mathrm{Im}(z)\geq -1/n} e^{-\beta|z|}|\phi(z)| \tag{4.14}$$

is finite. With this norm, $\mathrm{Exp}_+(\beta, n)$ is a Banach space. Let $\mathrm{Exp}_+(\beta)$ be their inductive limit with respect to $n = 1, 2, \cdots$

$$\mathrm{Exp}_+(\beta) = \varinjlim_{n\geq 1} \mathrm{Exp}_+(\beta, n) = \bigcup_{n\geq 1} \mathrm{Exp}_+(\beta, n). \tag{4.15}$$

Thus $\mathrm{Exp}_+(\beta)$ is the set of holomorphic functions near the upper half plane that can grow at most the rate $e^{\beta|z|}$. Next, define Exp_+ to be their inductive limit with respect to $\beta = 0, 1, 2, \cdots$

$$\mathrm{Exp}_+ = \varinjlim_{\beta\geq 0} \mathrm{Exp}_+(\beta) = \bigcup_{\beta\geq 0} \mathrm{Exp}_+(\beta). \tag{4.16}$$

Thus Exp_+ is the set of holomorphic functions near the upper half plane that can grow at most exponentially (see also Fig. 9); $\phi(z)$ in Exp_+ satisfies $||\phi||_{\beta,n} < \infty$ for some β, n, and such β and n can depend on ϕ. Topological properties of Exp_+ will be discussed in Sec.5.2. In this section, the topology on Exp_+ is not used.

Finally, we make the assumption for the parameter K. Although we are interested in a positive K from a physical viewpoint, any complex number is considered in the proof of the next theorem. In the theory of functions, if a function $f(\lambda)$ converges to $w \in \mathbf{C}$ along a path γ tending to ∞, w and γ are called an asymptotic value and an asymptotic path, respectively. Let $\mathcal{K} \subset \mathbf{C} \cup \{\infty\}$ be the set of numbers K such that $2/K$ is an asymptotic value of the function $2\pi g(\lambda)$. Since $D(\lambda) \to \infty$ as $|\lambda| \to 0$ (see Claim below), $2/K$ is an asymptotic value of $2\pi g(\lambda)$ if and only if it is an asymptotic value of the function $D(\lambda) + 2\pi g(-\sqrt{-1}\lambda)$. Define a subset $\mathcal{K}_0 \subset \mathcal{K}$ as follows: $K \in \mathcal{K}_0$ if and only if there is an asymptotic path γ included in the angular domain $\pi/2 + \delta < \arg(\lambda) < 3\pi/2 - \delta$ such that

$$D(\lambda) + 2\pi g(-\sqrt{-1}\lambda) = \frac{2}{K} + o(\frac{1}{|\lambda|^n}), \quad |\lambda| \to \infty, \tag{4.17}$$

holds along γ for every $n = 0, 1, 2, \cdots$, where $\delta > 0$ is the constant appeared in the assumption (A3). That is, if the function $D(\lambda) + 2\pi g(-\sqrt{-1}\lambda)$ converges to a value $2/K$ faster than any polynomials along a path γ in the angular domain, then $K \in \mathcal{K}_0$. Put $\mathcal{K}_1 = \mathcal{K}_0 \cup \{0\} \cup \{\infty\}$. It is well known as the Denjoy-Ahlfors theorem that if $g(\lambda)$ is an entire function of order p, then the number of asymptotic values is at most $2p$ [50].

In this case, $\mathcal{K}_1$ is a finite set. If $g(\lambda)$ is an entire function of infinite order or $g(\lambda)$ is a meromorphic function, the set of asymptotic values $\mathcal{K}$ can have the cardinality of the continuum (see Gross [19] and Eremenko [16] for such examples; see also Bergweiler and Eremenko [6] for a condition that a meromorphic function has at most finite asymptotic values). However, since the condition (4.17) and the condition that $g(\lambda)$ is an integrable real-valued function on $\mathbf{R}$ are too strong, it seems that $\mathcal{K}_1 = \{0, \infty\}$ in most cases (actually, the author does not know an example of a probability density function $g(\omega)$ satisfying (A1) to (A3) whose cardinality of $\mathcal{K}_1$ is larger than two).

The main theorem in this section is stated as follows.

Theorem 4.4. Suppose that $K \notin \mathcal{K}_1$ and $g(\omega)$ satisfies the assumptions (A1) to (A3). For any $\phi, \psi \in \mathrm{Exp}_+$, there exists a positive number t_0 such that the semigroup $e^{T_1 t}$ satisfies the equality

$$(e^{T_1 t}\phi, \overline{\psi}) = \sum_{n=0}^{M} S_n[t, \phi, \psi] e^{\xi_n t} + \sum_{n=0}^{\infty} R_n[t, \phi, \psi] e^{\lambda_n t}, \tag{4.18}$$

for $t > t_0$, where $\xi_0, \cdots, \xi_M$ are eigenvalues of T_1 on the right half plane, $S_n[t, \phi, \psi] e^{\xi_n t}$ are corresponding residues of $F_0(\lambda) e^{\lambda t}$, and where $\lambda_0, \lambda_1, \cdots$ are resonance poles of T_1 such that $|\lambda_0| \leq |\lambda_1| \leq \cdots$, and $R_n[t, \phi, \psi] e^{\lambda_n t}$ are corresponding residues of $F_1(\lambda) e^{\lambda t}$. In particular when $0 < K \leq K_c$, it is written as

$$(e^{T_1 t}\phi, \overline{\psi}) = \sum_{n=0}^{\infty} R_n[t, \phi, \psi] e^{\lambda_n t}. \tag{4.19}$$

If λ_n is a pole of order k, then R_n is a polynomial of degree at most $k-1$ in t. When $0 < K < K_c$, all resonance poles lie on the left half plane. In particular, the order parameter $\eta(t) = (e^{T_1 t}\phi, P_0)$ for the linearized system (3.4) decays to zero exponentially as $t \to \infty$.

To prove Eq.(4.18), we need the next estimate.

Claim. For any $\phi, \psi \in \mathrm{Exp}_+$, $((\lambda - \sqrt{-1}\mathcal{M})^{-1}\phi, \overline{\psi})$ tends to zero as $|\lambda| \to \infty$.

Proof. This is a well known property of the resolvent when λ is included in the resolvent set. Indeed, by the definition, we have

$$\begin{aligned} ((\lambda - \sqrt{-1}\mathcal{M})^{-1}\phi, \overline{\psi}) &= \int_{\mathbf{R}} \frac{1}{\lambda - \sqrt{-1}\omega} \phi(\omega)\psi(\omega)g(\omega)d\omega \\ &= \int_{\mathbf{R}} \frac{x}{x^2 + (\omega - y)^2} q(\omega)d\omega + \sqrt{-1}\int_{\mathbf{R}} \frac{\omega - y}{x^2 + (\omega - y)^2} q(\omega)d\omega, \end{aligned}$$

where we put $\lambda = x + \sqrt{-1}y$ and $q(\omega) = \phi(\omega)\psi(\omega)g(\omega)$. Since $\phi, \psi \in \mathrm{Exp}_+$, there exist $C_1, C_2, \beta_1, \beta_2 \geq 0$ such that $|\phi(z)| \leq C_1 e^{\beta_1 |z|}$, $|\psi(z)| \leq C_2 e^{\beta_2 |z|}$. Thus we obtain

$$|((\lambda - \sqrt{-1}\mathcal{M})^{-1}\phi, \overline{\psi})| \leq C_1 C_2 \int_{\mathbf{R}} \frac{1}{|\lambda - \sqrt{-1}\omega|} e^{(\beta_1 + \beta_2)|\omega|} g(\omega)d\omega.$$

By the assumption (A2) for $g(\omega)$, this integral exists when $\lambda \notin \sqrt{-1}\mathbf{R}$. When $\lambda \notin \sqrt{-1}\mathbf{R}$, $1/|\lambda - \sqrt{-1}\omega| \to 0$ as $|\lambda| \to \infty$ uniformly in $\omega \in \mathbf{R}$, which proves that $|((\lambda -$

$\sqrt{-1}\mathcal{M})^{-1}\phi, \overline{\psi})| \to 0$. In particular, an upper bound of $|((\lambda - \sqrt{-1}\mathcal{M})^{-1}\phi, \overline{\psi})|$ is determined by C_1, C_2, β_1 and β_2.

To estimate the case $\lambda \in \sqrt{-1}\mathbf{R}$, take the limit $x \to +0$ to yield

$$\lim_{x\to+0}((\lambda - \sqrt{-1}\mathcal{M})^{-1}\phi, \overline{\psi}) = \pi q(y) - \pi\sqrt{-1}V(y),$$

where $V(y)$ is defined by Eq.(4.9). Due to the assumption (A2), $|q(y)| \leq C_1 C_2 e^{(\beta_1+\beta_2)|y|} g(y)$ tends to zero as $y \to \pm\infty$. Next, let us prove $V(y) \to 0$ as $y \to \pm\infty$. Due to the assumption (A3), $|q(\lambda)|$ tends to zero as $|\lambda| \to \infty$ in the angular domain (4.13). This fact and Cauchy's theorem allow us to rewrite $V(y)$ as

$$\begin{aligned} V(y) &= \frac{1}{\pi}\int_0^\infty \frac{1}{t}\left(q(y-t) - q(y+t)\right)dt \\ &= \frac{1}{\pi}\int_0^{\sqrt{-1}a} \frac{1}{t}\left(q(y-t) - q(y+t)\right)dt + \frac{1}{\pi}\int_{\sqrt{-1}a}^{\sqrt{-1}a+\infty} \frac{1}{t}\left(q(y-t) - q(y+t)\right)dt, \end{aligned}$$

where a is a sufficiently small positive number. By the assumption (A3), the second term above tends to zero as $y \to \pm\infty$. For the first term, there exists a number $\kappa \in \mathbf{C}$ such that

$$\int_0^{\sqrt{-1}a} \frac{1}{t}\left(q(y-t) - q(y+t)\right)dt = 2\int_0^{\sqrt{-1}a} \frac{dq}{dy}(y + \kappa t)dt. \tag{4.20}$$

Since $q(z)$ is holomorphic on the upper half plane except for poles of $g(z)$, Cauchy's formula is applied to provide

$$\frac{dq}{dy}(z_0) = \frac{1}{2\pi\sqrt{-1}}\int_C \frac{1}{(z-z_0)^2} q(z)dz = \frac{1}{2\pi}\int_0^{2\pi} \frac{1}{ae^{\sqrt{-1}\theta}} q(z_0 + ae^{\sqrt{-1}\theta})d\theta$$

with $z_0 = y + \kappa t$. By the assumption (A3), $q(y + \kappa t + ae^{\sqrt{-1}\theta}) \to 0$ as $y \to \pm\infty$ uniformly in $0 \leq \theta \leq 2\pi$. Therefore, Eq.(4.20) tends to zero as $y \to \pm\infty$. This completes the proof of Claim. ■

From the above proof, we see that $((\lambda - \sqrt{-1}\mathcal{M})^{-1}\phi, \overline{\psi})$ has an upper bound determined by constants C_1, C_2, β_1 and β_2.

Proof of Thm.4.4. Let us prove Eq.(4.18). Take a sequence $\{r_n\}_{n=0}^\infty$ of positive numbers as follows: if $2/K$ is not an asymptotic value of the function $D(\lambda) + 2\pi g(-\sqrt{-1}\lambda)$, then there exist a positive constant A and a sequence $\{r_n\}_{n=0}^\infty$ with $r_n \to \infty$ such that

$$\left|1 - \pi K g(-\sqrt{-1}\lambda) - \frac{K}{2}D(\lambda)\right| \geq A, \tag{4.21}$$

for $\lambda = r_n e^{\sqrt{-1}\theta}$, $\pi/2 + \delta < \theta < 3\pi/2 - \delta$. Even if $2/K$ is an asymptotic value of the function $D(\lambda) + 2\pi g(-\sqrt{-1}\lambda)$, because of Eq.(4.17), there exist a positive constant A, an integer $m \geq 0$ and a sequence $\{r_n\}_{n=0}^\infty$ with $r_n \to \infty$ such that

$$\left|1 - \pi K g(-\sqrt{-1}\lambda) - \frac{K}{2}D(\lambda)\right| \geq \frac{A}{r_n^m}, \tag{4.22}$$



(a) (b)

Fig. 7: The contour for the Laplace inversion formula.

holds for $\lambda = r_n e^{\sqrt{-1}\theta}$, $\pi/2 + \delta < \theta < 3\pi/2 - \delta$.

Recall that eigenvalues of T_1 are roots of the equation $D(\lambda) = 2/K$. Since $D(\lambda) \to 0$ as $|\lambda| \to \infty$, the number of eigenvalues on the right half plane is finite, say $\xi_0, \cdots, \xi_M$. Take a positive number $d > 0$ so that $\mathrm{Re}(\xi_j) < d$ for $j = 0, \cdots, M$. With these d and r_n, take paths C_1 to C_6 as are shown in Fig.7 (b):

$$\begin{aligned}
C_1 &= \{d + \sqrt{-1}y \,|\, -r_n \leq y \leq r_n\},\\
C_2 &= \{x + \sqrt{-1}r_n \,|\, 0 \leq x \leq d\},\\
C_3 &= \{r_n e^{\sqrt{-1}\theta} \,|\, \pi/2 \leq \theta \leq \pi/2 + \delta\},\\
C_4 &= \{r_n e^{\sqrt{-1}\theta} \,|\, \pi/2 + \delta \leq \theta \leq 3\pi/2 - \delta\},
\end{aligned}$$

and C_5 and C_6 are defined in a similar way to C_3 and C_2, respectively. We put $C^{(n)} = \sum_{j=1}^{6} C_j$.

Let $\lambda_0, \lambda_1, \cdots, \lambda_{\#(n)}$ be resonance poles inside the closed curve $C^{(n)}$. By the definition of r_n, there are no resonance poles on the curve $C^{(n)}$. By the residue theorem, we have

$$2\pi\sqrt{-1}\left(\sum_{j=0}^{M} S_j[t,\phi,\psi]e^{\xi_j t} + \sum_{j=0}^{\#(n)} R_j[t,\phi,\psi]e^{\lambda_j t}\right) = \int_{C_6+C_1+C_2} e^{\lambda t}F_0(\lambda)d\lambda + \int_{C_3+C_4+C_5} e^{\lambda t}F_1(\lambda)d\lambda,$$

when r_n is sufficiently large so that $C^{(n)}$ includes all eigenvalues $\xi_0, \cdots, \xi_M$. Since the integral $\int_{C_1} e^{\lambda t}F_0(\lambda)d\lambda/(2\pi\sqrt{-1})$ converges to $(e^{T_1 t}\phi, \overline{\psi})$ as $n \to \infty$,

$$\begin{aligned}
(e^{T_1 t}\phi, \overline{\psi}) = &\sum_{j=0}^{M} S_j[t,\phi,\psi]e^{\xi_j t} + \lim_{n\to\infty}\sum_{j=0}^{\#(n)} R_j[t,\phi,\psi]e^{\lambda_j t}\\
&- \frac{1}{2\pi\sqrt{-1}}\lim_{n\to\infty}\int_{C_6+C_2} e^{\lambda t}F_0(\lambda)d\lambda - \frac{1}{2\pi\sqrt{-1}}\lim_{n\to\infty}\int_{C_3+C_4+C_5} e^{\lambda t}F_1(\lambda)d\lambda. \quad (4.23)
\end{aligned}$$

It is easy to verify that the integrals along C_2, C_3, C_5, C_6 tend to zero as $n \to \infty$ because of the assumption (A3). For example, the integral along C_2 is estimated as

$$\begin{aligned}\Big|\int_{C_2} e^{\lambda t} F_0(\lambda) d\lambda\Big| &= \Big|\int_d^0 e^{(x+\sqrt{-1}r_n)t} F_0(x+\sqrt{-1}r_n) dx\Big| \\ &\leq e^{dt}\int_0^d |F_0(x+\sqrt{-1}r_n)| dx,\end{aligned}$$

where F_0 is given as (4.2). By using Claim above, the integral along C_2 proves to be zero as $n \to \infty$ ($r_n \to \infty$). The integrals along C_3, C_5, C_6 are estimated in a similar manner. The integral along C_4 is estimated as

$$\begin{aligned}\left|\int_{C_4} e^{\lambda t} F_1(\lambda) d\lambda\right| &\leq \int_{\pi/2+\delta}^{3\pi/2-\delta} r_n e^{r_n t \cos\theta} |F_1(r_n e^{\sqrt{-1}\theta})| d\theta \\ &\leq \max_{\pi/2+\delta \leq \theta \leq 3\pi/2-\delta} |F_1(r_n e^{\sqrt{-1}\theta})| \int_\delta^{\pi/2} 2r_n e^{-r_n t \sin\theta} d\theta \\ &\leq \max_{\pi/2+\delta \leq \theta \leq 3\pi/2-\delta} |F_1(r_n e^{\sqrt{-1}\theta})| \int_\delta^{\pi/2} 2r_n e^{-2r_n t\theta/\pi} d\theta \\ &\leq \max_{\pi/2+\delta \leq \theta \leq 3\pi/2-\delta} |F_1(r_n e^{\sqrt{-1}\theta})| \cdot \frac{\pi}{t}\left(e^{-2r_n t\delta/\pi} - e^{-r_n t}\right). \qquad (4.24)\end{aligned}$$

Since $|(\lambda - \sqrt{-1}\mathcal{M})^{-1}| \to 0$ as $|\lambda| \to \infty$, $F_1(\lambda)$ given by Eq.(4.10) is estimated as

$$|F_1(\lambda)| \leq \frac{D_0 + |D_1 + D_2\phi(-\sqrt{-1}\lambda) + D_3\psi(-\sqrt{-1}\lambda) + D_4\phi(-\sqrt{-1}\lambda)\psi(-\sqrt{-1}\lambda)| \cdot |g(-\sqrt{-1}\lambda)|}{|1 - \pi K g(-\sqrt{-1}\lambda) - \frac{K}{2}D(\lambda)|},$$

where D_0 to D_4 are some positive constants. Since $\phi, \psi \in \mathrm{Exp}_+$, there exist $C_1, C_2, \beta_1, \beta_2 \geq 0$ such that $|\phi(z)| \leq C_1 e^{\beta_1 |z|}$, $|\psi(z)| \leq C_2 e^{\beta_2 |z|}$:

$$|F_1(\lambda)| \leq \frac{D_0 + \left(D_1 + D_2 C_1 e^{\beta_1|\lambda|} + D_3 C_2 e^{\beta_2|\lambda|} + D_4 C_1 C_2 e^{(\beta_1+\beta_2)|\lambda|}\right) \cdot |g(-\sqrt{-1}\lambda)|}{|1 - \pi K g(-\sqrt{-1}\lambda) - \frac{K}{2}D(\lambda)|}.$$

Suppose that $g(-\sqrt{-1}r_n e^{\sqrt{-1}\theta})$ diverges as $n \to \infty$ ($r_n \to \infty$). Then, there exists $D_5 > 0$, which has an upper bound determined by constants C_1, C_2, β_1 and β_2, such that $|F_1(r_n e^{\sqrt{-1}\theta})|$ is estimated as

$$|F_1(r_n e^{\sqrt{-1}\theta})| \leq D_5 e^{(\beta_1+\beta_2)r_n}. \qquad (4.25)$$

If $g(-\sqrt{-1}r_n e^{\sqrt{-1}\theta})$ is bounded, Eq.(4.22) shows that there exists $D_6 > 0$, which has an upper bound determined by constants C_1, C_2, β_1 and β_2, such that

$$|F_1(r_n e^{\sqrt{-1}\theta})| \leq \frac{D_6}{A} r_n^m e^{(\beta_1+\beta_2)r_n} \cdot |g(-\sqrt{-1}r_n e^{\sqrt{-1}\theta})|. \qquad (4.26)$$

Therefore, we obtain

$$\left|\int_{C_4} e^{\lambda t} F_1(\lambda) d\lambda\right| \leq \frac{D_7 r_n^m}{t}\left(e^{(\beta_1+\beta_2-2\delta t/\pi)r_n} - e^{(\beta_1+\beta_2-t)r_n}\right), \qquad (4.27)$$

with some $D_7 > 0$. Thus if $t > t_0 := \pi(\beta_1 + \beta_2)/(2\delta)$, this integral tends to zero as $n \to \infty$, which proves Eq.(4.18). From the above calculation, it turns out that the integrals along C_2 to C_6 have upper bounds determined by C_1, C_2, β_1 and β_2 once g is fixed. This fact will be used to prove Thm.5.10.

In particular when $0 < K \leq K_c$, there are no eigenvalues on the right half plane (Thm.3.5). Thus Eq.(4.18) is reduced to Eq.(4.19).

Finally, we prove that all resonance poles are on the left half plane when $0 < K < K_c$. Resonance poles are roots of Eq.(4.11), which is reduced to Eq.(3.17) on the imaginary axis. We have already shown that K_c is the least value such that Eq.(3.17) has a root y. Therefore, when $0 < K < K_c$, there are no resonance poles on the imaginary axis. ■

The condition $K \notin \mathcal{K}_1$ is used to prove that the infinite series in Eq.(4.18) converges. However, to prove the exponential decay of the order parameter $\eta(t) = (e^{T_1 t}\phi, P_0)$, we do not need the infinite series such as Eq.(4.18). In Sec.5.3, we will give another expression of the semigroup $(e^{T_1 t}\phi, \overline{\psi})$ consisting of a finite sum and a certain integral, which proves that $\eta(t)$ decays exponentially without the assumption $K \notin \mathcal{K}_1$. Nevertheless, for almost all choices of $g(\omega)$, $\mathcal{K}_1 = \{0, \infty\}$ and thus $K \notin \mathcal{K}_1$ is not a restriction.

Remark. Because of the assumption (A3), there are no resonance poles of T_1 on the area

$$\{z \in \mathbf{C} \,|\, |\arg(z) - \pi/2| \leq \delta,\ |\arg(z) - 3\pi/2| \leq \delta,\ |z| \geq r\}, \tag{4.28}$$

if $r > 0$ is sufficiently large. Further, the left hand side of Eq.(4.11) is meromorphic. This shows that on each vertical line $\{\lambda \,|\, \mathrm{Re}(\lambda) = a \leq 0\}$, there are only finitely many resonance poles. Thus $\mathrm{Re}(\lambda_n) \to -\infty$ as $n \to \infty$, and we can renumber the index of resonance poles λ_n so that

$$0 \geq \mathrm{Re}(\lambda_0) \geq \mathrm{Re}(\lambda_1) \geq \mathrm{Re}(\lambda_2) \geq \cdots .$$

Note that $\mathrm{Exp}_+(0)$ is the set of bounded holomorphic functions near the upper half plane. From the proof above, we immediately obtain the following.

Corollary 4.5. Suppose that $K \notin \mathcal{K}_1$ and $g(\omega)$ satisfies the assumptions (A1) to (A3). If $\phi, \psi \in \mathrm{Exp}_+(0)$, then Eq.(4.18) is true for $t > 0$.

4.3 Rational case

In this subsection, we suppose that $g(\omega)$ is a rational function. Since $g(\omega)$ does not decay so fast as $|\omega| \to \infty$, we should choose moderate functions for ϕ and ψ. Let $\mathbf{C}_+ = \{z \in \mathbf{C} \,|\, \mathrm{Im}(z) \geq 0\}$ be the real axis and the upper half plane. Let H_+ be the set of bounded holomorphic functions on $\mathbf{C}_+$. With the norm

$$\|\phi\| := \sup_{\mathrm{Im}(z) \geq 0} |\phi(z)|, \tag{4.29}$$

H_+ is a Banach space.

It is remarkable that if $g(\omega)$ is a rational function, Eq.(4.11) is reduced to an algebraic equation. Thus the number of resonance poles is finite. The proof of the following theorem is similar to that of Thm.4.4 and omitted here (since $F_0(\lambda)$ and $F_1(\lambda)$ are rational functions, estimates of integrals in Eq.(4.23) are too easy. We need not introduce the set $\mathcal{K}$).

Theorem 4.6. Suppose that $0 < K \leq K_c$ and $g(\omega)$ is a rational function. For any $\phi, \psi \in H_+$, the semigroup $e^{T_1 t}$ satisfies the equality

$$(e^{T_1 t}\phi, \overline{\psi}) = \sum_{n=0}^{M} R_n[t, \phi, \psi] e^{\lambda_n t}, \tag{4.30}$$

for $t > 0$, where $\lambda_0, \cdots, \lambda_M$ are resonance poles of T_1 and $R_0[t, \phi, \psi]e^{\lambda_0 t}, \cdots, R_M[t, \phi, \psi]e^{\lambda_M t}$ are corresponding residues of $F_1(\lambda)e^{\lambda t}$. In particular, when $0 < K < K_c$, $\lambda_0, \cdots, \lambda_M$ are on the left half plane and the order parameter $\eta(t) = (e^{T_1 t}\phi, P_0)$ for the linearized system (3.4) decays to zero exponentially as $t \to \infty$.

Since the right hand side of Eq.(4.30) is a finite sum, the semigroup $e^{T_1 t}$ looks like an exponential of a matrix. The reason of this fact will be revealed in Sec.5.4 by means of the theory of rigged Hilbert spaces.

Example 4.7. If $g(\omega) = 1/(\pi(1 + \omega^2))$ is the Lorentzian distribution, a resonance pole is given by $\lambda = K/2 - 1$ (a root of Eq.(4.11)). Therefore $\eta(t)$ decays with the exponential rates $e^{(K/2-1)t}$. Note that the transition point is $K_c = 2/\pi/g(0) = 2$.

5 Spectral theory

We have proved that when $K > K_c$, the de-synchronous state ($\eta(t) \equiv 0$) is linearly unstable because of eigenvalues on the right half plane, while when $0 < K < K_c$, it is linearly stable because of resonance poles on the left half plane. Next, we want to investigate bifurcations at $K = K_c$. However, a center manifold in a usual sense is of infinite dimensional because the continuous spectrum lies on the imaginary axis. To handle this difficulty, we develop a spectral theory of resonance poles based on a rigged Hilbert space.

5.1 Rigged Hilbert space

Let X be a topological vector space and X' its dual space. X' is a set of continuous anti-linear functionals on X. We use the Dirac's notation; for $\mu \in X'$ and $\varphi \in X$, $\mu(\varphi)$ is denoted by $\langle \varphi \,|\, \mu \rangle$. For any $a, b \in \mathbf{C}$ and $\phi, \psi \in X$, $\mu, \xi \in X'$, the equalities

$$\langle a\phi + b\psi \,|\, \mu \rangle = a\langle \phi \,|\, \mu \rangle + b\langle \psi \,|\, \mu \rangle, \tag{5.1}$$

$$\langle \phi \,|\, a\mu + b\xi \rangle = \overline{a}\langle \phi \,|\, \mu \rangle + \overline{b}\langle \phi \,|\, \xi \rangle, \tag{5.2}$$

hold. Several topologies can be defined on the dual space X'. Two of the most usual topologies are the weak dual topology and the strong dual topology [46]. A sequence $\{\mu_j\} \subset X'$ is said to be weakly convergent to $\mu \in X'$ if $\langle \phi \,|\, \mu_j \rangle \to \langle \phi \,|\, \mu \rangle$ for each $\phi \in X$; a sequence $\{\mu_j\} \subset X'$ is said to be strongly convergent to $\mu \in X'$ if $\langle \phi \,|\, \mu_j \rangle \to \langle \phi \,|\, \mu \rangle$ uniformly on any bounded subset $A \subset X$.

Let H be a Hilbert space with the inner product $(\cdot\,,\,\cdot)_H$ such that X is a dense subspace of H. Since a Hilbert space is isomorphic to its dual space, we obtain $H \subset X'$ through $H \simeq H'$.

Definition 5.1. If a topological vector space X is a dense subspace of a Hilbert space H and a topology of X is stronger than that of H, the triplet

$$X \subset H \subset X' \tag{5.3}$$

is called the *rigged Hilbert space* or the *Gelfand triplet*. The *canonical inclusion* $i : X \to X'$ is defined as follows; for $\psi \in X$, we denote $i(\psi)$ by $|\psi\rangle$, which is defined to be

$$i(\psi)(\phi) = \langle \phi \,|\, \psi \rangle = (\phi, \psi)_H, \tag{5.4}$$

for any $\phi \in X$. If the inner product on H is Hermitian, the alternative definition

$$i(\psi)(\phi) = \langle \phi \,|\, \overline{\psi} \rangle = (\phi, \overline{\psi})_H. \tag{5.5}$$

is used so that $i(a\psi)(\phi) = ai(\psi)(\phi)$ for $a \in \mathbf{C}$. After section 5.2, we will use the latter one. Thus if $H = L^2(\mathbf{R}, g(\omega)d\omega)$, then

$$i(\psi)(\phi) = \int_{\mathbf{R}} \phi(\omega)\psi(\omega)g(\omega)d\omega.$$

Let $A : X \to X$ be a continuous linear operator on X. Then, the dual operator $A^{\times} : X' \to X'$ defined through

$$\langle \phi \,|\, A^{\times}\mu \rangle = \langle A\phi \,|\, \mu \rangle, \quad \phi \in X,\ \mu \in X' \tag{5.6}$$

is also continuous on X' for both of the weak dual topology and the strong dual topology. We can show the equality

$$A^{\times} i(\psi) = i(A\psi), \tag{5.7}$$

for any $\psi \in X$.

It is easy to show that the canonical inclusion is injective if and only if X is a dense subspace of H, and the canonical inclusion is continuous (for both of the weak dual topology and the strong dual topology) if and only if a topology of X is stronger than that of H (see Tréves [46]). If X is not dense in H, two functionals on H may not be distinguished as functionals on X. As a result, $H' \not\subset X'$ in general.

Definition 5.2. When $X \subset H$ is not a dense subspace of H, the triplet (X, H, X') is called the *degenerate* rigged Hilbert space.

For applications to the Kuramoto model, we investigate two triplets, $\mathrm{Exp}_+ \subset L^2(\mathbf{R}, g(\omega)d\omega) \subset \mathrm{Exp}_+'$, and a degenerate one $(H_+, L^2(\mathbf{R}, g(\omega)d\omega), H_+')$.

5.2 Spectral theory on $\mathrm{Exp}_+ \subset L^2(\mathbf{R}, g(\omega)d\omega) \subset \mathrm{Exp}'_+$

In this subsection, we suppose that $g(\omega)$ satisfies the assumptions (A1) to (A3). Since g decays faster than any exponential functions $e^{-\beta|\omega|}$, we have $\mathrm{Exp}_+ \subset L^2(\mathbf{R}, g(\omega)d\omega)$, and indeed, Exp_+ is dense in $L^2(\mathbf{R}, g(\omega)d\omega)$ and the topology of Exp_+ is stronger than that of $L^2(\mathbf{R}, g(\omega)d\omega)$ (see Prop.5.3 below). Thus the rigged Hilbert space $\mathrm{Exp}_+ \subset L^2(\mathbf{R}, g(\omega)d\omega) \subset \mathrm{Exp}'_+$ is well defined. Recall that $\mathrm{Exp}_+(\beta, n)$ is the Banach space of holomorphic functions on $\mathbf{C}_n$ with the norm $|| \cdot ||_{\beta,n}$, and $\mathrm{Exp}_+(\beta)$ is their inductive limit with respect to $n \geq 1$. By the definition of the inductive limit, the topology of $\mathrm{Exp}_+(\beta)$ is defined as follows: a set $U \subset \mathrm{Exp}_+(\beta)$ is open if and only if $U \cap \mathrm{Exp}_+(\beta, n)$ is open for every $n \geq 1$. This implies that the inclusions $\mathrm{Exp}_+(\beta, n) \to \mathrm{Exp}_+(\beta)$ are continuous for every $n \geq 1$. Similarly, Exp_+ is an inductive limit of $\mathrm{Exp}_+(\beta)$, and its topology is induced from that of $\mathrm{Exp}_+(\beta)$: a set $U \subset \mathrm{Exp}_+$ is open if and only if $U \cap \mathrm{Exp}_+(\beta)$ is open for every $\beta = 0, 1, \cdots$. The inclusions $\mathrm{Exp}_+(\beta) \to \mathrm{Exp}_+$ are continuous for every $\beta = 0, 1, \cdots$. On the dual space Exp'_+, both of the weak dual topology and the strong dual topology can be introduced as was explained (see also Fig. 9).

A topological vector space X is called Montel if every bounded set of X is relatively compact. A Montel space has a convenient property that on a bounded set A of a dual space of a Montel space, the weak dual topology coincides with the strong dual topology. In particular, a weakly convergent series in a dual of a Montel space also converges with respect to the strong dual topology (see Tréves [46]). This property is very important to develop a theory of generalized functions.

The topology of Exp_+ has following properties.

Proposition 5.3. Exp_+ is a topological vector space satisfying

(i) Exp_+ is a complete Montel space.
(ii) if $\{\phi_j\}_{j=1}^\infty$ is a convergent series in Exp_+, there exist $n \geq 1$ and $\beta \geq 0$ such that $\{\phi_j\}_{j=1}^\infty \subset \mathrm{Exp}_+(\beta, n)$ and $\{\phi_j\}_{j=1}^\infty$ converges with respect to the norm $|| \cdot ||_{\beta,n}$.
(iii) Exp_+ is a dense subspace of $L^2(\mathbf{R}, g(\omega)d\omega)$.
(iv) the topology of Exp_+ is stronger than that of $L^2(\mathbf{R}, g(\omega)d\omega)$.

Proof. (i) At first, we prove that $\mathrm{Exp}_+(\beta)$ is Montel. To do so, it is sufficient to show that the inclusion $\mathrm{Exp}_+(\beta, n) \to \mathrm{Exp}_+(\beta, n+1)$ is a compact operator for every n (see Grothendieck [20], Chap.4.3.3). To prove it, let A be a bounded set of $\mathrm{Exp}_+(\beta, n)$. There exists a constant C such that $||\phi||_{\beta,n} = \sup_{z\in\mathbf{C}_n} e^{-\beta|z|}|\phi(z)| < C$ for any $\phi \in A$. This means that the set A is locally bounded in the interior of $\mathbf{C}_n$. Therefore, for any sequence $\{\phi_j\}_{j=1}^\infty \subset A$, there exists a subsequence converging to some holomorphic function ψ uniformly on compact subsets in $\mathbf{C}_n$ (Montel's theorem). In particular, the subsequence converges to ψ on $\mathbf{C}_{n+1}$, and it satisfies $||\psi||_{\beta,n+1} < C$ and $\psi \in \mathrm{Exp}_+(\beta, n+1)$. This proves that the inclusion $\mathrm{Exp}_+(\beta, n) \to \mathrm{Exp}_+(\beta, n+1)$ is compact and thus $\mathrm{Exp}_+(\beta)$ is Montel. In a similar manner, we can prove by using the Montel's theorem that the inclusion $\mathrm{Exp}_+(\beta) \to \mathrm{Exp}_+(\beta+1)$ is a compact operator for every $\beta = 0, 1, \cdots$, which proves that Exp_+ is also Montel. Next, we show that Exp_+ is complete. Since $\mathrm{Exp}_+(\beta, n)$ is a Banach space, in particular it is a DF space, their inductive limit $\mathrm{Exp}_+(\beta)$ is a DF space by virtue of Prop.5 in Chap.4.3.3 of [20], in which it is shown that an inductive limit of DF spaces is DF. The same proposition also shows that the inductive limit Exp_+ of DF spaces $\mathrm{Exp}_+(\beta)$ is a DF space. Since Exp_+

is Montel and DF, it is complete because of Cor.2 in Chap.4.3.3 of [20].

(ii) It is known that if the inclusion $\mathrm{Exp}_+(\beta) \to \mathrm{Exp}_+(\beta+1)$ is a compact operator for every $\beta = 0, 1, \cdots$, then, for any bounded set $A \subset \mathrm{Exp}_+$, there exists β such that $A \subset \mathrm{Exp}_+(\beta)$ and A is bounded on $\mathrm{Exp}_+(\beta)$ (see Komatsu [23] and references therein). By using the same fact again, it turns out that for any bounded set $A \subset \mathrm{Exp}_+$, there exist β and n such that $A \subset \mathrm{Exp}_+(\beta, n)$. In particular, since a convergent series $\{\phi_j\}_{j=1}^{\infty}$ is bounded, there exists β and n such that $\{\phi_j\}_{j=1}^{\infty} \subset \mathrm{Exp}_+(\beta, n)$ and it converges with respect to the topology of $\mathrm{Exp}_+(\beta, n)$.

To prove (iii), note that $L^2(\mathbf{R}, g(\omega)d\omega)$ is obtained by the completion of the set of polynomials because the probability density $g(\omega)$ has all moments due to the assumption (A2). Obviously Exp_+ includes all polynomials, and thus Exp_+ is dense in $L^2(\mathbf{R}, g(\omega)d\omega)$.

For (iv), note that Exp_+ satisfies the first axiom of countability because it is defined through the inductive limits of Banach spaces. Therefore, to prove (iv), it is sufficient to show that the inclusion $\mathrm{Exp}_+ \to L^2(\mathbf{R}, g(\omega)d\omega)$ is sequentially continuous. Let $\{\phi_j\}_{j=1}^{\infty}$ be a sequence in Exp_+ which converges to zero. By (ii), there exist β and n such that $\{\phi_j\}_{j=1}^{\infty}$ converges in the topology of $\mathrm{Exp}_+(\beta, n)$: $||\phi_j||_{\beta,n} \to 0$. Then,

$$\begin{aligned}
||\phi_j||^2_{L^2(\mathbf{R},g(\omega)d\omega)} &= \int_{-\infty}^{\infty} |\phi_j(\omega)|^2 g(\omega)d\omega \\
&\leq \sup_{\omega\in\mathbf{R}} e^{-2\beta|\omega|}|\phi_j(\omega)|^2 \int_{-\infty}^{\infty} e^{2\beta|\omega|} g(\omega)d\omega \\
&\leq ||\phi_j||^2_{\beta,n} \int_{-\infty}^{\infty} e^{2\beta|\omega|} g(\omega)d\omega.
\end{aligned}$$

By the assumption (A2) for $g(\omega)$, the right hand side exists and tends to zero as $j \to \infty$. This means that the inclusion $\mathrm{Exp}_+ \to L^2(\mathbf{R}, g(\omega)d\omega)$ is continuous. ■

The topology of the dual space Exp_+' has following properties.

Proposition 5.4.

(i) Exp_+' is a complete Montel space with respect to the strong dual topology.
(ii) Exp_+' is sequentially complete with respect to the weak dual topology; that is, for a sequence $\{\mu_j\}_{j=1}^{\infty}$, if $\langle \phi \,|\, \mu_j \rangle$ converges to some complex number C_ϕ for every $\phi \in \mathrm{Exp}_+$ as $j \to \infty$, then there exists $\mu \in \mathrm{Exp}_+'$ such that $C_\phi = \langle \phi \,|\, \mu \rangle$ and $\mu_j \to \mu$ with respect to the strong dual topology.

Proof. (i) It is known that the strong dual of a Montel space is Montel and complete, see Tréves [46]. (ii) Suppose that $\langle \phi \,|\, \mu_j \rangle$ converges to some complex number C_ϕ for every $\phi \in \mathrm{Exp}_+$. This means that the set $\{\mu_j\}_{j=1}^{\infty}$ is weakly bounded and is a Cauchy sequence with respect to the weak dual topology. As was explained before, on a bounded set of a dual space of a Montel space, the weak dual topology and the strong dual topology coincide with one another. Thus $\{\mu_j\}_{j=1}^{\infty}$ is a Cauchy sequence with respect to the strong dual topology. Since Exp_+' is complete with respect to the strong dual topology, μ_j converges to some element $\mu \in \mathrm{Exp}_+'$. In particular, $\langle \phi \,|\, \mu_j \rangle$ converges to $\langle \phi \,|\, \mu \rangle = C_\phi$. ■

Next, we restrict the domain of the operator $T_1 = \sqrt{-1}\mathcal{M} + \frac{K}{2}\mathcal{P}$ to Exp_+. We simply denote $T_1|_{\mathrm{Exp}_+}$ by T_1. We will see that T_1 is quite moderate if restricted to Exp_+.

Proposition 5.5.
(i) The operator $T_1 : \mathrm{Exp}_+ \to \mathrm{Exp}_+$ is continuous (note that it is not continuous on $L^2(\mathbf{R}, g(\omega)d\omega)$).
(ii) The operator $T_1 : \mathrm{Exp}_+ \to \mathrm{Exp}_+$ generates a holomorphic semigroup $e^{T_1 t} : \mathrm{Exp}_+ \to \mathrm{Exp}_+$ on the positive t axis (note that it is not holomorphic on $L^2(\mathbf{R}, g(\omega)d\omega)$).
(iii) Eigenvalues of $T_1 : \mathrm{Exp}_+ \to \mathrm{Exp}_+$ are the same as those of $T_1 : L^2(\mathbf{R}, g(\omega)d\omega) \to L^2(\mathbf{R}, g(\omega)d\omega)$ (*i.e.* the roots of $D(\lambda) = 2/K$ on $\mathrm{Re}(\lambda) > 0$). The residual spectrum of $T_1 : \mathrm{Exp}_+ \to \mathrm{Exp}_+$ is the imaginary axis and the left half plane. There is no continuous spectrum of $T_1 : \mathrm{Exp}_+ \to \mathrm{Exp}_+$.

Proof. (i) It is easy to see by the definition that if $\phi \in \mathrm{Exp}_+$, then $T_1\phi \in \mathrm{Exp}_+$. Let $\{\phi_j\}_{j=1}^{\infty}$ be a sequence in Exp_+ converging to zero. By Prop.5.3 (ii), there exist $\beta \geq 0$ and $n \geq 1$ such that $||\phi_j||_{\beta,n} \to 0$. For any $\varepsilon > 0$, $||T_1\phi_j||_{\beta+\varepsilon,n}$ is estimated as

$$\begin{aligned}
||T_1\phi_j||_{\beta+\varepsilon,n} &\leq || \sqrt{-1}\omega\phi_j||_{\beta+\varepsilon,n} + \frac{K}{2}|(\phi_j, P_0)| \cdot ||P_0||_{\beta+\varepsilon,n} \\
&\leq \sup_{\omega\in\mathbf{C}_n} e^{-(\beta+\varepsilon)|\omega|}|\omega\phi_j(\omega)| + \frac{K}{2}||\phi_j||_{L^2(\mathbf{R},g(\omega)d\omega)} \\
&\leq ||\phi_j||_{\beta,n} \cdot \sup_{\omega\in\mathbf{C}_n} e^{-\varepsilon|\omega|}|\omega| + \frac{K}{2}||\phi_j||_{L^2(\mathbf{R},g(\omega)d\omega)},
\end{aligned}$$

which tends to zero as $j \to \infty$. This proves that $T_1\phi_j$ tends to zero as $j \to \infty$ with respect to the topology of Exp_+, and thus $T_1 : \mathrm{Exp}_+ \to \mathrm{Exp}_+$ is continuous.
(ii) We know that the operator T_1 generates the semigroup $e^{T_1 t}$ as an operator on $L^2(\mathbf{R}, g(\omega)d\omega)$ (see Sec.4.1). In other words, the differential equation

$$\frac{d}{dt}x(t,\omega) = T_1 x(t,\omega) = \sqrt{-1}\omega x_1(t,\omega) + \frac{K}{2}(x(t,\cdot), P_0) \tag{5.8}$$

has a unique solution $x(t,\omega) = e^{T_1 t}\phi(\omega)$ in $L^2(\mathbf{R}, g(\omega)d\omega)$ if an initial condition ϕ is in $L^2(\mathbf{R}, g(\omega)d\omega)$. We have to prove that if $\phi \in \mathrm{Exp}_+$, then $x(t,\cdot) \in \mathrm{Exp}_+$. For this purpose, we integrate the above equation as

$$e^{T_1 t}\phi(\omega) = e^{\sqrt{-1}\omega t}\phi(\omega) + \frac{K}{2}\int_0^t e^{\sqrt{-1}\omega(t-s)}(e^{T_1 s}\phi, P_0)ds. \tag{5.9}$$

From this expression, it is obvious that if $\phi \in \mathrm{Exp}_+(\beta, n)$, then $e^{T_1 t}\phi(\omega) \in \mathrm{Exp}_+$. Next, we show that $e^{T_1 t}$ is a holomorphic semigroup. The proof is done in the same way as the standard proof of the existence of holomorphic semigroups [22]. Let δ be a positive number appeared in the assumption (A3). In the proof of Thm.4.4, we see that the Laplace integral along C_2, C_3, C_5 and C_6 tends to zero as $r_n \to \infty$ (see Fig.7 (b) and Eq.(4.23)). This means that the contour (the straight line represented as Fig.7 (a)) for the Laplace inversion formula (4.4) can be deformed to the contour Γ represented in Fig.8, which goes to infinity along the rays $\arg(\lambda) = \pm(\pi/2 + \delta)$:

$$(e^{T_1 t}\phi, \psi) = \frac{1}{2\pi\sqrt{-1}}\int_\Gamma e^{\lambda t}((\lambda - T_1)^{-1}\phi, \psi)d\lambda, \tag{5.10}$$



Fig. 8: The contour for the Laplace inversion formula.

for $t > 0$. We show that this integral exists even if t is a complex number. For this purpose, put $\lambda = re^{\pm\sqrt{-1}(\delta+\pi/2)} \in \Gamma$ and $t = |t|e^{\sqrt{-1}\theta}$ with $|\theta| < \delta - \varepsilon$, where $\varepsilon > 0$ is an arbitrarily small number. Then

$$|e^{\lambda t}| = |e^{r|t|\cos(\theta\pm(\delta+\pi/2))}| < e^{-r|t|\sin\varepsilon}.$$

Further, $((\lambda - T_1)^{-1}\phi, \psi)$ decays as $O(1/|\lambda|)$ as $|\lambda| \to \infty$ along Γ because of Eq.(4.5), the assumption (A3) and Claim in the proof of Thm.4.4. This proves that the integral in Eq.(5.10) exists for $t = |t|e^{\sqrt{-1}\theta}$, $|\theta| < \delta - \varepsilon$, which gives an analytic continuation of $(e^{T_1 t}\phi, \psi)$ to the sector $\{t \in \mathbf{C} \mid \arg(t) < \delta\}$. In particular, $(e^{T_1 t}\phi, P_0)$ is holomorphic in the sector. Now Eq.(5.9) proves that $e^{T_1 t}\phi$ is also holomorphic and included in Exp_+ in the sector $\{t \in \mathbf{C} \mid \arg(t) < \delta\}$.
(iii) Eigenvalues are calculated in the same way as Prop.3.2 (i). Note that corresponding eigenvectors $1/(\lambda - \sqrt{-1}\omega)$ are in Exp_+ when $\mathrm{Re}(\lambda) > 0$. Next, from the proof of Lemma 4.1, it turns out that the resolvent is given by

$$(\lambda - T_1)^{-1}\phi = \frac{1}{\lambda - \sqrt{-1}\omega}\left(\phi(\omega) + \frac{K}{2}\frac{1}{1 - KD(\lambda)/2}\int_{-\infty}^{\infty}\frac{1}{\lambda - \sqrt{-1}\omega}\phi(\omega)g(\omega)d\omega\right), \quad (5.11)$$

for $\phi \in \mathrm{Exp}_+$. Suppose that $\mathrm{Re}(\lambda) \leq 0$. Since $1/(\lambda - \sqrt{-1}\omega) \notin \mathrm{Exp}_+$, $(\lambda - T_1)^{-1}\phi \in \mathrm{Exp}_+$ if and only if ϕ satisfies

$$\phi(\omega) + \frac{K}{2}\frac{1}{1 - KD(\lambda)/2}\int_{-\infty}^{\infty}\frac{1}{\lambda - \sqrt{-1}\omega}\phi(\omega)g(\omega)d\omega = (\lambda - \sqrt{-1}\omega)\psi(\omega)$$

for some $\psi \in \mathrm{Exp}_+$. In particular, ϕ has to satisfy

$$\left(1 - \frac{K}{2}D(\lambda)\right)\phi(-\sqrt{-1}\lambda) = -\frac{K}{2}\int_{-\infty}^{\infty}\frac{1}{\lambda - \sqrt{-1}\omega}\phi(\omega)g(\omega)d\omega. \quad (5.12)$$

For fixed λ, define continuous linear functionals F_1, F_2 on Exp_+ to be

$$\begin{aligned} F_1[\phi] &= \left(1 - \frac{K}{2}D(\lambda)\right)\phi(-\sqrt{-1}\lambda), \\ F_2[\phi] &= -\frac{K}{2}\int_{-\infty}^{\infty}\frac{1}{\lambda - \sqrt{-1}\omega}\phi(\omega)g(\omega)d\omega, \end{aligned}$$

respectively. If $F_1 = F_2$ on a dense subset of Exp_+, then $F_1 = F_2$ for any $\phi \in \mathrm{Exp}_+$ because F_1 and F_2 are continuous. However, it is obvious that $F_1[1] \neq F_2[1]$. This proves that the set of functions ϕ satisfying (5.12) is not dense in Exp_+, and thus the domain of $(\lambda - T_1)^{-1}$ is not dense in Exp_+ when $\mathrm{Re}(\lambda) \leq 0$. Therefore, the imaginary axis and the left half plane are included in the residual spectrum. Finally, suppose $\mathrm{Re}(\lambda) > 0$ and λ is not an eigenvalue. Then, $(\lambda - T_1)^{-1}\phi \in \mathrm{Exp}_+$ for any $\phi \in \mathrm{Exp}_+$. From Eq.(5.11), it is easy to prove that if a sequence $\{\phi_j\}_{j=1}^{\infty}$ tends to zero in Exp_+, then $\|(\lambda - T_1)^{-1}\phi_j\|_{\beta,n}$ tends to zero for some β and n, which proves that $(\lambda - T_1)^{-1}\phi_j \to 0$ in Exp_+. Therefore, $(\lambda - T_1)^{-1}$ is continuous and T_1 has no continuous spectrum on the right half plane. ■

Eigenvalues of T_1 are given as roots of the equation $D(\lambda) = 2/K$, $\mathrm{Re}(\lambda) > 0$, and corresponding eigenvectors are

$$v_\lambda(\omega) = \frac{1}{\lambda - \sqrt{-1}\omega} \in \mathrm{Exp}_+. \tag{5.13}$$

If we regard it as a functional on Exp_+ through the canonical inclusion $i : \mathrm{Exp}_+ \to \mathrm{Exp}_+'$, it acts on Exp_+ as

$$i(v_\lambda)(\phi) = \langle \phi \,|\, \overline{v}_\lambda \rangle = (\phi, \overline{v}_\lambda) = \int_{\mathbf{R}} \phi(\omega)v_\lambda(\omega)g(\omega)d\omega = \int_{\mathbf{R}} \frac{1}{\lambda - \sqrt{-1}\omega}\phi(\omega)g(\omega)d\omega, \tag{5.14}$$

for $\phi \in \mathrm{Exp}_+$. Due to Eq.(4.7), the analytic continuation of this value from right to left is given as

$$\int_{\mathbf{R}} \frac{1}{\lambda - \sqrt{-1}\omega}\phi(\omega)g(\omega)d\omega + 2\pi\phi(-\sqrt{-1}\lambda)g(-\sqrt{-1}\lambda). \tag{5.15}$$

Motivated by this observation, let us define a linear functional $\mu(\lambda) \in \mathrm{Exp}_+'$ to be

$$\langle \phi \,|\, \mu(\lambda) \rangle = \int_{\mathbf{R}} \frac{1}{\lambda - \sqrt{-1}\omega}\phi(\omega)g(\omega)d\omega + 2\pi\phi(-\sqrt{-1}\lambda)g(-\sqrt{-1}\lambda), \tag{5.16}$$

when $\mathrm{Re}(\lambda) < 0$, and

$$\langle \phi \,|\, \mu(\lambda) \rangle = \lim_{x \to +0} \int_{\mathbf{R}} \frac{1}{(x + \sqrt{-1}y) - \sqrt{-1}\omega}\phi(\omega)g(\omega)d\omega, \tag{5.17}$$

when $\lambda = \sqrt{-1}y$, $y \in \mathbf{R}$. Indeed, $\mu(\lambda)$ is continuous and thus an element of Exp_+'. Let $\{\phi_m\}_{m=1}^{\infty}$ be a sequence in Exp_+ converging to zero. By the definition of the topology on Exp_+, $\phi_m(\omega)$ converges to zero uniformly on compact sets on the upper half plane. Hence, the second term $\phi(-\sqrt{-1}\lambda)g(-\sqrt{-1}\lambda)$ of Eq.(5.16) tends to zero as $m \to \infty$. The integral of the first term of Eq.(5.16) is decomposed into three terms $\int_{-M}^{M} \cdots d\omega$, $\int_{M}^{\infty} \cdots d\omega$ and $\int_{-\infty}^{-M} \cdots d\omega$. Since $\phi_m(\omega) \to 0$ uniformly on compact sets, the term $\int_{-M}^{M} \cdots d\omega$ tends to zero. Since the integral $\int_{\mathbf{R}} \cdots d\omega$ exists, two terms $\int_{M}^{\infty} \cdots d\omega$ and $\int_{-\infty}^{-M} \cdots d\omega$ become sufficiently small if M is sufficiently large. This proves that $\langle \phi_m \,|\, \mu(\lambda) \rangle \to 0$ as $m \to \infty$. Similarly, we can verify for the case $\lambda = \sqrt{-1}y$ that $\mu(\lambda)$ is a continuous functional (note that the existence of the limit in Eq.(5.17) was shown in the proof of Lemma 4.2).

We expect that $\mu(\lambda)$ plays a similar role to the eigenvectors. Indeed, we can prove the following theorem.

Theorem 5.6. Let $\lambda_0, \lambda_1, \cdots$ be the resonance poles of the operator T_1 and $T_1^{\times}$ the dual operator of T_1 defined through

$$\langle \phi \,|\, T_1^{\times} \xi \rangle = \langle T_1 \phi \,|\, \xi \rangle, \quad \phi \in \mathrm{Exp}_+,\ \xi \in \mathrm{Exp}_+'. \tag{5.18}$$

Then, the equality

$$T_1^{\times} \mu(\lambda_n) = \overline{\lambda}_n \mu(\lambda_n) \tag{5.19}$$

holds for $n = 0, 1, 2, \cdots$. In this sense, $\overline{\lambda}_n$ is an eigenvalue of $T_1^{\times}$, and $\mu(\lambda_n)$ is an eigenvector. In what follows, $\mu(\lambda_n)$ is denoted by μ_n and we call it the *generalized eigenfunction* associated with the resonance pole λ_n.

Proof. The proof is straightforward. Suppose that $\mathrm{Re}(\lambda_n) < 0$. For any $\phi \in \mathrm{Exp}_+$,

$$\begin{aligned}
\langle \phi \,|\, T_1^{\times} \mu_n \rangle &= \langle T_1 \phi \,|\, \mu_n \rangle \\
&= \int_{\mathbf{R}} \frac{1}{\lambda_n - \sqrt{-1}\omega} (T_1\phi)(\omega) g(\omega) d\omega + 2\pi (T_1 \phi)(-\sqrt{-1}\lambda_n) g(-\sqrt{-1}\lambda_n) \\
&= \int_{\mathbf{R}} \frac{\sqrt{-1}\omega}{\lambda_n - \sqrt{-1}\omega} \phi(\omega) g(\omega) d\omega + \frac{K}{2} \int_{\mathbf{R}} \frac{1}{\lambda_n - \sqrt{-1}\omega} g(\omega) d\omega \cdot \int_{\mathbf{R}} \phi(\omega) g(\omega) d\omega \\
&\quad + 2\pi \left(\lambda_n \phi(-\sqrt{-1}\lambda_n) + \frac{K}{2} \int_{\mathbf{R}} \phi(\omega) g(\omega) d\omega \right) g(-\sqrt{-1}\lambda_n) \\
&= \lambda_n \left(\int_{\mathbf{R}} \frac{1}{\lambda_n - \sqrt{-1}\omega} \phi(\omega) g(\omega) d\omega + 2\pi \phi(-\sqrt{-1}\lambda_n) g(-\sqrt{-1}\lambda_n) \right) \\
&\quad + \frac{K}{2} \int_{\mathbf{R}} \phi(\omega) g(\omega) d\omega \left(D(\lambda_n) + 2\pi g(-\sqrt{-1}\lambda_n) - \frac{2}{K} \right) \\
&= \lambda_n \langle \phi \,|\, \mu_n \rangle + \frac{K}{2} \int_{\mathbf{R}} \phi(\omega) g(\omega) d\omega \left(D(\lambda_n) + 2\pi g(-\sqrt{-1}\lambda_n) - \frac{2}{K} \right).
\end{aligned} \tag{5.20}$$

Since λ_n is a resonance pole, it is a root of Eq.(4.11). Thus we obtain

$$\langle \phi \,|\, T_1^{\times} \mu_n \rangle = \lambda_n \langle \phi \,|\, \mu_n \rangle = \langle \phi \,|\, \overline{\lambda}_n \mu_n \rangle,$$

which proves the theorem. The proof for the case $\mathrm{Re}(\lambda_n) = 0$ is done in the same way. ■

Define a dual semigroup $(e^{T_1 t})^\times$ through

$$\langle \phi \,|\, (e^{T_1 t})^\times \mu \rangle = \langle e^{T_1 t}\phi \,|\, \mu \rangle. \tag{5.21}$$

for any $\phi \in \mathrm{Exp}_+$ and $\mu \in \mathrm{Exp}_+'$.

Proposition 5.7. (i) A solution of the initial value problem

$$\frac{d}{dt}\xi = T_1^\times \xi, \quad \xi(0) = \mu \in \mathrm{Exp}_+' \tag{5.22}$$

in Exp_+', which is continuous in an initial condition with respect to the weak and the strong dual topologies, is uniquely given by $\xi(t) = (e^{T_1 t})^\times \mu$.
(ii) $(e^{T_1 t})^\times$ has eigenvalues $e^{\overline{\lambda}_0 t}, e^{\overline{\lambda}_1 t}, \cdots$, where $\lambda_0, \lambda_1, \cdots$ are resonance poles of T_1.

Proof. (i) For any $\phi \in \mathrm{Exp}_+$, $(e^{T_1 t})^\times \mu$ satisfies

$$\begin{aligned}\frac{d}{dt}\langle \phi \,|\, (e^{T_1 t})^\times \mu \rangle &= \frac{d}{dt}\langle e^{T_1 t}\phi \,|\, \mu \rangle = \langle T_1 e^{T_1 t}\phi \,|\, \mu \rangle \\ &= \langle e^{T_1 t} T_1 \phi \,|\, \mu \rangle = \langle \phi \,|\, T_1^\times (e^{T_1 t})^\times \mu \rangle.\end{aligned}$$

This provides

$$\frac{d}{dt}(e^{T_1 t})^\times \mu = T_1^\times (e^{T_1 t})^\times \mu. \tag{5.23}$$

It is easy to verify from Eq.(5.21) that when $t = 0$, $(e^{T_1 t})^\times$ is the identity. This proves that $(e^{T_1 t})^\times \mu$ is a solution of the initial value problem (5.22).

To prove the uniqueness of a solution, let us show that a solution of the equation $d\phi/dt = T_1\phi$ in Exp_+ is unique (this fact was mentioned in the proof of Thm.5.5 (ii)). Indeed, let ϕ_1 and ϕ_2 be two solutions of the equation with the same initial condition in Exp_+. Then, Eq.(5.9) provides

$$\phi_1(t,\omega) - \phi_2(t,\omega) = \frac{K}{2}\int_0^t e^{\sqrt{-1}\omega(t-s)} (\phi_1(s,\,\cdot\,) - \phi_2(s,\,\cdot\,), P_0) ds.$$

This is estimated in $L^2(\mathbf{R}, g(\omega)d\omega)$ as

$$\begin{aligned}||\phi_1(t,\,\cdot\,) - \phi_2(t,\,\cdot\,)||_{L^2(\mathbf{R},g(\omega)d\omega)} &\leq \frac{K}{2}\int_0^t |(\phi_1(s,\,\cdot\,) - \phi_2(s,\,\cdot\,), P_0)| ds \\ &\leq \frac{K}{2}\int_0^t ||\phi_1(s,\,\cdot\,) - \phi_2(s,\,\cdot\,)||_{L^2(\mathbf{R},g(\omega)d\omega)} ds.\end{aligned}$$

Now the Gronwall inequality proves that $||\phi_1(t,\,\cdot\,) - \phi_2(t,\,\cdot\,)||_{L^2(\mathbf{R},g(\omega)d\omega)} = 0$, and thus a solution is unique in $L^2(\mathbf{R}, g(\omega)d\omega)$. Since $\mathrm{Exp}_+ \subset L^2(\mathbf{R}, g(\omega)d\omega)$, a solution is unique in Exp_+. For a solution $\phi(t,\omega) \in \mathrm{Exp}_+$, the canonical inclusion yields

$$\frac{d}{dt} i(\phi) = i(T_1\phi) = T_1^\times i(\phi).$$

Since $i : \mathrm{Exp}_+ \to \mathrm{Exp}'_+$ is injective, it turns out that a solution of (5.22) in $i(\mathrm{Exp}_+)$ is unique. Now we suppose that the initial value problem (5.22) has two solutions $f_1(t, \mu)$ and $f_2(t, \mu)$, which are continuous in $\mu \in \mathrm{Exp}'_+$. There exists a sequence $\{\zeta_n\}_{n=1}^{\infty} \subset i(\mathrm{Exp}_+)$ such that $\zeta_n \to \mu$ with respect to the weak dual topology and the strong dual topology because $i(\mathrm{Exp}_+)$ is dense in Exp'_+. Since a solution in $i(\mathrm{Exp}_+)$ is unique, $f_1(t, \zeta_n) = f_2(t, \zeta_n)$. Taking the limit yields $f_1(t, \mu) = f_2(t, \mu)$, and they coincide with $(e^{T_1 t})^{\times}\mu$.
(ii) Eq.(5.19) implies

$$\frac{d}{dt}\langle \phi \,|\, e^{\overline{\lambda}_n t}\mu_n \rangle = \langle \phi \,|\, e^{\overline{\lambda}_n t}\overline{\lambda}_n \mu_n \rangle = \langle \phi \,|\, e^{\overline{\lambda}_n t} T_1^{\times}\mu_n \rangle = \langle \phi \,|\, T_1^{\times}(e^{\overline{\lambda}_n t}\mu_n)\rangle.$$

Thus $\xi(t) = e^{\overline{\lambda}_n t}\mu_n$ is a solution of the equation $d\xi/dt = T_1^{\times}\xi$. Obviously $e^{\overline{\lambda}_n t}\mu_n$ is continuous in $\mu_n \in \mathrm{Exp}'_+$. By the uniqueness of a solution, we obtain $e^{\overline{\lambda}_n t}\mu_n = (e^{T_1 t})^{\times}\mu_n$. ■

If we define a semigroup $e^{T_1^{\times} t}$ generated by $T_1^{\times}$ to be the flow of (5.22), then Prop.5.7 (i) means $e^{T_1^{\times} t} = (e^{T_1 t})^{\times}$. Prop.5.7 (i) also implies that a solution of the inhomogeneous equation

$$\frac{d}{dt}\xi = T_1^{\times}\xi + f(t), \quad f(t) \in \mathrm{Exp}'_+, \tag{5.24}$$

which is continuous in an initial condition $\mu \in \mathrm{Exp}'_+$, is uniquely given by

$$\xi(t) = (e^{T_1 t})^{\times}\mu + \int_0^t (e^{T_1(t-s)})^{\times} f(s)ds. \tag{5.25}$$

This formula will be used so often when analyzing the nonlinear system (3.2),(3.3).

In what follows, we suppose that $0 < K \leq K_c$ and $K \notin \mathcal{K}_1$ so that Eq.(4.19) is applicable. To develop a spectral decomposition, we make the additional assumption.

(A4) All resonance poles are poles of first order (*i.e.* all resonance poles are simple roots of Eq.(4.11)).

Note that this is the assumption only for $g(\omega)$. For example, the Gaussian distribution satisfies (A4), see Appendix A for the proof. Under the assumption (A4), $R_n[t, \phi, \psi]$ in Eq.(4.19) is independent of t. Since $R_n[t, \phi, \psi]$ is the residue of $F_1(\lambda)$ given as Eq.(4.5), it is calculated as

$$R_n[t, \phi, \psi] = \frac{K}{2D_n}\langle \phi \,|\, \mu_n \rangle \langle \psi \,|\, \mu_n \rangle, \tag{5.26}$$

where

$$D_n := \lim_{\lambda \to \lambda_n} \frac{1}{\lambda - \lambda_n}\left(1 - \frac{K}{2}D(\lambda) - \pi K g(-\sqrt{-1}\lambda)\right) \tag{5.27}$$

is a constant which is independent of ϕ, ψ. Note that $Q[\lambda, \phi]$ given by Eq.(4.6) is just the definition of the functional $\mu(\lambda)$. Thus Eq.(4.19) is rewritten as

$$(e^{T_1 t}\phi, \overline{\psi}) = \sum_{n=0}^{\infty} \frac{K}{2D_n} e^{\lambda_n t}\langle \phi \,|\, \mu_n \rangle \langle \psi \,|\, \mu_n \rangle, \tag{5.28}$$

for $t > t_0$. By using the canonical inclusion i, the left hand side is rewritten as

$$(e^{T_1 t}\phi, \overline{\psi}) = \langle e^{T_1 t}\phi \,|\, \overline{\psi} \rangle = \langle \phi \,|\, (e^{T_1 t})^{\times}\overline{\psi} \rangle. \tag{5.29}$$

Therefore, we obtain

$$(e^{T_1 t})^{\times}|\overline{\psi}\rangle = \sum_{n=0}^{\infty} \frac{K}{2\overline{D}_n} e^{\overline{\lambda}_n t} \overline{\langle \psi \,|\, \mu_n \rangle} \cdot |\, \mu_n \rangle, \tag{5.30}$$

for $t > t_0$. Note that the complex conjugates arise in the coefficients because functionals are anti-linear. Since Eq.(5.30) comes from Eq.(5.28), the infinite series in the right hand side of Eq.(5.30) converges with respect to the weak dual topology on Exp'_+. However, since Exp_+ is Montel, it also converges with respect to the strong dual topology. We divide the infinite sum in Eq.(5.28) into two parts as

$$(e^{T_1 t}\phi, \overline{\psi}) = \sum_{n=0}^{M} \frac{K}{2D_n} e^{\lambda_n t} \langle \phi \,|\, \mu_n \rangle \langle \psi \,|\, \mu_n \rangle + \sum_{n=M+1}^{\infty} \frac{K}{2D_n} e^{\lambda_n t} \langle \phi \,|\, \mu_n \rangle \langle \psi \,|\, \mu_n \rangle, \tag{5.31}$$

where $M \in \mathbf{N}$ is any natural number. The second part $\sum_{n=M+1}^{\infty} [\cdots]$ does not converge when $0 < t < t_0$ in general. However, since $(e^{T_1 t}\phi, \overline{\psi})$ is holomorphic in $t > 0$ and continuous at $t = 0$, we obtain

$$(\phi, \overline{\psi}) = \sum_{n=0}^{M} \frac{K}{2D_n} \langle \phi \,|\, \mu_n \rangle \langle \psi \,|\, \mu_n \rangle + \lim_{t \to +0} \sum_{n=M+1}^{\infty} \frac{K}{2D_n} e^{\lambda_n t} \langle \phi \,|\, \mu_n \rangle \langle \psi \,|\, \mu_n \rangle, \tag{5.32}$$

where the second term has a meaning in the sense of an analytic continuation in t. Through the canonical inclusion, the above yields

$$|\,\overline{\psi}\rangle = \sum_{n=0}^{M} \frac{K}{2\overline{D}_n} \overline{\langle \psi \,|\, \mu_n \rangle} \cdot |\, \mu_n \rangle + \mathcal{R}_M[\psi], \tag{5.33}$$

$$\mathcal{R}_M[\psi] := \lim_{t \to +0} \sum_{n=M+1}^{\infty} \frac{K}{2\overline{D}_n} e^{\overline{\lambda}_n t} \overline{\langle \psi \,|\, \mu_n \rangle} \cdot |\, \mu_n \rangle,$$

which gives the spectral decomposition of $|\,\overline{\psi}\rangle \in i(\mathrm{Exp}_+)$.

Theorem 5.8 (Spectral decomposition). Suppose that $0 < K \leq K_c$, $K \notin \mathcal{K}_1$ and $g(\omega)$ satisfies the assumptions (A1) to (A4).
(i) A system of generalized eigenfunctions $\{\mu_n\}_{n=0}^{\infty}$ is complete in the sense that if $\langle \phi \,|\, \mu_n \rangle = 0$ for $n = 0, 1, \cdots$, then $\phi = 0$.
(ii) $\mu_0, \mu_1, \cdots$ are linearly independent of each other: if $\sum_{n=0}^{\infty} a_n |\, \mu_n \rangle = 0$ with $a_n \in \mathbf{C}$, then $a_n = 0$ for every n.
(iii) Let V_M be a complementary subspace of $\mathrm{span}\{\mu_0, \cdots, \mu_M\}$ in Exp'_+, which includes μ_j for every $j = M+1, M+2, \cdots$. Then, any $|\,\overline{\psi}\rangle \in i(\mathrm{Exp}_+)$ is uniquely decomposed with respect to the direct sum $\mathrm{span}\{\mu_0, \cdots, \mu_M\} \oplus V_M$ as Eq.(5.33), and this decomposition is independent of the choice of the complementary subspace V_M.

Proof. (i) If $\langle \phi \,|\, \mu_n \rangle = 0$ for all n, Eq.(5.28) provides $(e^{T_1 t}\phi, \overline{\psi}) = 0$ for any $\psi \in \mathrm{Exp}_+$. Since Exp_+ is dense in $L^2(\mathbf{R}, g(\omega)d\omega)$, we obtain $e^{T_1 t}\phi = 0$ for $t > t_0$. Since $e^{T_1 t}$ is holomorphic in $t > 0$ and strongly continuous at $t = 0$, we obtain $\phi = 0$ by taking the limit $t \to +0$.

(ii) Suppose that $\sum_{n=0}^{\infty} a_n|\mu_n\rangle = 0$. Since $(e^{T_1 t})^\times$ is continuous,

$$0 = (e^{T_1 t})^\times \sum_{n=0}^{\infty} a_n|\mu_n\rangle = \sum_{n=0}^{\infty} a_n (e^{T_1 t})^\times |\mu_n\rangle = \sum_{n=0}^{\infty} a_n e^{\overline{\lambda}_n t}|\mu_n\rangle.$$

We can assume that

$$0 \geq \mathrm{Re}(\lambda_0) \geq \mathrm{Re}(\lambda_1) \geq \mathrm{Re}(\lambda_2) \geq \cdots,$$

without loss of generality. Further, on each vertical line $\{\lambda \,|\, \mathrm{Re}(\lambda) = a \leq 0\}$, there are only finitely many resonance poles (see Remark in Sec.4.2). Suppose that $\mathrm{Re}(\lambda_0) = \cdots = \mathrm{Re}(\lambda_k)$ and $\mathrm{Re}(\lambda_k) > \mathrm{Re}(\lambda_{k+1})$. Then, the above equality provides

$$0 = \sum_{n=0}^{k} a_n e^{-\sqrt{-1}\mathrm{Im}(\lambda_n)t}|\mu_n\rangle + \sum_{n=k+1}^{\infty} a_n e^{(\overline{\lambda}_n - \mathrm{Re}(\lambda_0))t}|\mu_n\rangle.$$

Taking the limit $t \to \infty$ yields

$$0 = \lim_{t\to\infty} \sum_{n=0}^{k} a_n e^{-\sqrt{-1}\mathrm{Im}(\lambda_n)t}|\mu_n\rangle.$$

Since the finite set $\mu_0, \cdots, \mu_k$ of eigenvectors are linearly independent as in a finite-dimensional case, we obtain $a_n = 0$ for $n = 0, \cdots, k$. The same procedure is repeated to prove $a_n = 0$ for every n.

(iii) Let V_M and V'_M be two complementary subspaces of $\mathrm{span}\{\mu_0, \cdots, \mu_M\}$, both of which include $\mu_{M+1}, \mu_{M+2}, \cdots$. Let

$$\mathrm{Exp}'_+ = \mathrm{span}\{\mu_0, \cdots, \mu_M\} \oplus V_M = \mathrm{span}\{\mu_0, \cdots, \mu_M\} \oplus V'_M$$

be two direct sums and let

$$|\overline{\psi}\rangle = \sum_{n=0}^{M} a_n|\mu_n\rangle + v = \sum_{n=0}^{M} a'_n|\mu_n\rangle + v', \quad v \in V_M,\ v' \in V'_M$$

be corresponding decompositions. Then, $(e^{T_1 t})^\times|\overline{\psi}\rangle$ is given by

$$(e^{T_1 t})^\times|\overline{\psi}\rangle = \sum_{n=0}^{M} a_n e^{\overline{\lambda}_n t}|\mu_n\rangle + (e^{T_1 t})^\times v = \sum_{n=0}^{M} a'_n e^{\overline{\lambda}_n t}|\mu_n\rangle + (e^{T_1 t})^\times v'.$$

They give decompositions of $(e^{T_1 t})^\times|\overline{\psi}\rangle$ with respect to two direct sums

$$\mathrm{span}\{(e^{T_1 t})^\times\mu_0, \cdots, (e^{T_1 t})^\times\mu_n\} \oplus (e^{T_1 t})^\times V_M, \quad \mathrm{span}\{(e^{T_1 t})^\times\mu_0, \cdots, (e^{T_1 t})^\times\mu_n\} \oplus (e^{T_1 t})^\times V'_M,$$

respectively. Since $(e^{T_1 t})^\times|\mu_n\rangle = e^{\overline{\lambda}_n t}|\mu_n\rangle$, $\mathrm{span}\{(e^{T_1 t})^\times\mu_0, \cdots, (e^{T_1 t})^\times\mu_n\} = \mathrm{span}\{\mu_0, \cdots, \mu_n\}$, and the sets $(e^{T_1 t})^\times V_M$ and $(e^{T_1 t})^\times V'_M$ also include $\mu_{M+1}, \mu_{M+2}, \cdots$. Because of part (ii) of the theorem, the decomposition of $(e^{T_1 t})^\times|\overline{\psi}\rangle$ with respect to above direct sums is uniquely given as Eq.(5.30) for $t > t_0$. This implies

$$a_n = a'_n \ (= \frac{K}{2\overline{D}_n}\overline{\langle \psi \,|\, \mu_n\rangle}), \quad (e^{T_1 t})^\times v = (e^{T_1 t})^\times v' \ (= \sum_{n=M+1}^{\infty} \frac{K}{2\overline{D}_n} e^{\overline{\lambda}_n t}\overline{\langle \psi \,|\, \mu_n\rangle}\cdot |\,\mu_n\rangle).$$

Since $(e^{T_1 t})^\times$ is continuous in $t \geq 0$, we obtain $v = v'$. ■

When $K > K_c$, the spectral decomposition involves eigenvalues and eigenvectors in Exp_+. In particular, if all eigenvalues are not degenerate (that is, all roots of Eq.(3.12) are single roots), then Eq.(4.18) proves

$$(e^{T_1 t})^\times|\overline{\psi}\rangle = \sum_{n=0}^{M} \frac{K}{2\overline{E}_n} e^{\overline{\xi}_n t}\overline{\langle \psi \,|\, \overline{v}_n\rangle}\cdot |\,\overline{v}_n\rangle + \sum_{n=0}^{\infty} \frac{K}{2\overline{D}_n} e^{\overline{\lambda}_n t}\overline{\langle \psi \,|\, \mu_n\rangle}\cdot |\,\mu_n\rangle, \tag{5.34}$$

for $t > t_0$, and

$$|\,\overline{\psi}\rangle = \sum_{n=0}^{M} \frac{K}{2\overline{E}_n} \overline{\langle \psi \,|\, \overline{v}_n\rangle}\cdot |\,\overline{v}_n\rangle + \sum_{n=0}^{M'} \frac{K}{2\overline{D}_n} \overline{\langle \psi \,|\, \mu_n\rangle}\cdot |\,\mu_n\rangle + \mathcal{R}_{M'}[\psi], \tag{5.35}$$

where $\xi_0, \cdots, \xi_M$ are eigenvalues of T_1 on the right half plane, $v_n := v_{\xi_n}(\omega) \in \mathrm{Exp}_+$ are corresponding eigenvectors defined by Eq.(5.13), and where E_n are defined to be

$$E_n = \lim_{\xi \to \xi_n} \frac{1}{\xi - \xi_n}\left(1 - \frac{K}{2}D(\xi)\right). \tag{5.36}$$

It is easy to extend Thm.5.8 to the case that $K > K_c$: a system of eigenvectors $\{v_n\}_{n=0}^{M}$ and generalized eigenfunctions $\{\mu_n\}_{n=0}^{\infty}$ is complete, and they are linearly independent. If eigenvalues and resonance poles are degenerate, the right hand sides of Eq.(5.33) and (5.35) become more complex involving polynomials in t. In this paper, we do not consider such cases for simplicity, though it will be treated in the same way as a finite dimensional case because multiplicities of eigenvalues and resonance poles are finite.

This theorem suggests the expression of the projection to the generalized eigenspace.

Definition 5.9. Denote by $\Pi_n : \mathrm{Exp}'_+ \to \mathrm{span}\{\mu_n\} \subset \mathrm{Exp}'_+$ the projection to the generalized eigenspace. For $|\,\overline{\psi}\rangle \in i(\mathrm{Exp}_+)$, it is given as

$$\Pi_n|\,\overline{\psi}\rangle = \frac{K}{2\overline{D}_n}\overline{\langle \psi \,|\, \mu_n\rangle}\cdot |\,\mu_n\rangle. \tag{5.37}$$

Unfortunately, the projection Π_n is not a continuous operator . For example, put $\psi_m(\omega) = e^{-\sqrt{-1}m\omega}$. Then, $i(\psi_m) = |\,e^{\sqrt{-1}m\omega}\rangle$ converges to zero as $m \to \pm\infty$ with respect to the weak dual topology by virtue of the Riemann-Lebesgue lemma. However,

$$\langle \psi_m \,|\, \mu_n\rangle = \int_{\mathbf{R}} \frac{1}{\lambda_n - \sqrt{-1}\omega} e^{-\sqrt{-1}m\omega} g(\omega)d\omega + 2\pi e^{-m\lambda_n} g(-\sqrt{-1}\lambda_n) \tag{5.38}$$

does not tend to zero. It diverges as $m \to \infty$ when $\mathrm{Re}(\lambda_n) < 0$. This means that $\Pi_n : \mathrm{Exp}'_+ \to \mathrm{Exp}'_+$ is not continuous with respect to the weak dual topology. To avoid such a difficulty caused by "weakness" of the topology of the domain, we restrict the domain of Π_n. For constants $C \geq 1$ and $\alpha \geq 0$, define a subset $V_{C,\alpha} \subset \mathrm{Exp}_+$ to be

$$V_{C,\alpha} = \{\phi \in \mathrm{Exp}_+ \,|\, |\phi(z)|e^{-\alpha|z|} \leq C \text{ when } \mathrm{Im}(z) \geq 0\}. \tag{5.39}$$

We restrict the domain of the projection to $i(V_{C,\alpha}) \subset \mathrm{Exp}'_+$ and denote it by $\Pi_n^{C,\alpha} = \Pi_n|_{i(V_{C,\alpha})}$. Note that the set $\{e^{-\sqrt{-1}m\omega}\}_{m=0}^{\infty}$ above is *not* included in $V_{C,\alpha}$. To discuss the continuity of $\Pi_n^{C,\alpha}$, let us introduce the projective topology on Exp'_+ (see also Fig. 9 and Table 1). In the dual space Exp'_+, the weak dual topology and the strong dual topology are defined. Another topology called the projective topology is defined as follows: Recall that $\mathrm{Exp}_+(\beta, n)$ is a Banach space, and its strong dual $\mathrm{Exp}_+(\beta, n)'$ is a Banach space with the norm

$$||\mu||^*_{\beta,n} := \sup_{||\phi||_{\beta,n}=1} |\langle \phi \,|\, \mu \rangle|. \tag{5.40}$$

We introduce the projective topology on $\mathrm{Exp}_+(\beta)' = \bigcap_{n\geq 1} \mathrm{Exp}_+(\beta, n)'$ as follows: $U \subset \mathrm{Exp}_+(\beta)'$ is open if and only if there exist open sets $U_n \subset \mathrm{Exp}_+(\beta, n)'$ such that $U_n \cap \mathrm{Exp}_+(\beta)' = U$ for every $n \geq 1$. It is known that the projective topology is equivalent to that induced by the metric

$$d_\beta(\mu_1, \mu_2) := \sum_{n=1}^{\infty} \frac{1}{2^n} \frac{||\mu_1 - \mu_2||^*_{\beta,n}}{1 + ||\mu_1 - \mu_2||^*_{\beta,n}}, \tag{5.41}$$

see Gelfand and Shilov [17]. When the projective topology is equipped, $\mathrm{Exp}_+(\beta)'$ is called the projective limit of $\mathrm{Exp}_+(\beta, n)'$ and denoted by $\mathrm{Exp}_+(\beta)' = \varprojlim \mathrm{Exp}_+(\beta, n)'$. In a similar manner, the projective topology on $\mathrm{Exp}'_+ = \bigcap_{\beta\geq 0} \mathrm{Exp}_+(\beta)'$ is introduced so that $U \subset \mathrm{Exp}'_+$ is open if and only if there exist open sets $U_\beta \subset \mathrm{Exp}_+(\beta)'$ such that $U_\beta \cap \mathrm{Exp}'_+ = U$ for every $\beta \geq 0$. This topology coincides with the topology induced by the metric

$$d(\mu_1, \mu_2) := \sum_{\beta=0}^{\infty} \frac{1}{2^\beta} \frac{d_\beta(\mu_1, \mu_2)}{1 + d_\beta(\mu_1, \mu_2)}. \tag{5.42}$$

In this manner, Exp'_+ equipped with the projective topology is a complete metric vector space. When the projective topology is equipped, Exp_+ is called the projective limit of $\mathrm{Exp}_+(\beta)'$ and denoted by $\mathrm{Exp}'_+ = \varprojlim \mathrm{Exp}_+(\beta)'$.

By the definition, the projective topology on Exp'_+ is weaker than the strong dual topology and stronger than the weak dual topology. Since Exp_+ is a Montel space, the weak dual topology coincides with the strong dual topology on any bounded set of Exp'_+. This implies that the projective topology also coincides with the weak dual topology and the strong dual topology on any bounded set of Exp'_+. In particular, a weakly convergent series in Exp'_+ also converges with respect to the metric d and the strong dual topology. Note that the projective topology is stronger than that induced by the norm $||\cdot||^*_{\beta,n}$. Therefore, a convergent series with respect to the metric d also converges with respect to the

$$\begin{tikzcd}[column sep=small, row sep=tiny]
\mathrm{Exp}_+(0) \subset \cdots \subset & \mathrm{Exp}_+(\beta) & \subset \cdots \subset & \mathrm{Exp}_+ & \subset L^2(\mathbf{R}, g(\omega)d\omega) \subset & \mathrm{Exp}'_+ & \subset \cdots \subset & \mathrm{Exp}_+(\beta)' & \subset \cdots \subset \mathrm{Exp}_+(0)' \\
 & \cup & & \cup & & \cup & & \cap & \\
 & \vdots & & V_{C,\beta} \arrow[rr, "\text{canonical}\\\text{inclusion}"'] & & i(V_{C,\beta}) & & \vdots & \\
 & \cup & & & & & & \cap & \\
 & \mathrm{Exp}_+(\beta, n) & & & & & & \mathrm{Exp}_+(\beta, n)' & \\
 & \cup & & & & & & \cap & \\
 & \vdots & & & & & & \vdots & \\
 & \cup & & & & & & \cap & \\
 & \mathrm{Exp}_+(\beta, 1) & & & & & & \mathrm{Exp}_+(\beta, 1)' &
\end{tikzcd}$$

Fig. 9: A diagram for the rigged Hilbert space $\mathrm{Exp}_+ \subset L^2(\mathbf{R}, g(\omega)d\omega) \subset \mathrm{Exp}'_+$.

$\mathrm{Exp}_+(\beta, n)$	Banach space: $\|\phi\|_{\beta,n} = \sup\limits_{\mathrm{Im}(z)\geq -1/n} \|\phi(z)\|e^{-\beta\|z\|}$
$\mathrm{Exp}_+(\beta, n)'$	Banach space: $\|\xi\|^*_{\beta,n} = \sup\limits_{\|\phi\|_{\beta,n}=1} \|\langle \phi \,\|\, \xi \rangle\|$
$\mathrm{Exp}_+(\beta)' = \varprojlim \mathrm{Exp}_+(\beta, n)'$	$d_\beta(\xi, \zeta) = \sum\limits_{n=1}^{\infty} \dfrac{1}{2^n} \dfrac{\|\xi - \zeta\|^*_{\beta,n}}{1 + \|\xi - \zeta\|^*_{\beta,n}}$
$\mathrm{Exp}'_+ = \varprojlim \mathrm{Exp}_+(\beta)'$	$d(\xi, \zeta) = \sum\limits_{\beta=0}^{\infty} \dfrac{1}{2^\beta} \dfrac{d_\beta(\xi, \zeta)}{1 + d_\beta(\xi, \zeta)}$

Table 1: Metric vector spaces used in Section 5.

norm $|| \cdot ||^*_{\beta,n}$ for every β and n.

Theorem 5.10. For any $C \geq 1$ and $\alpha \geq 0$, the following holds.
(i) On $i(V_{C,\alpha}) \subset \mathrm{Exp}'_+$, the weak dual topology, the projective topology and the strong dual topology coincide with one another.
(ii) The closure $\overline{i(V_{C,\alpha})}$ of $i(V_{C,\alpha})$ is a connected, compact metric space.
(iii) When $K \notin \mathcal{K}_1$, $\Pi_j^{C,\alpha} : i(V_{C,\alpha}) \to \mathrm{Exp}'_+$ is continuous for every $j = 0, 1, \cdots$ with respect to the projective topology and sequentially continuous with respect to the weak dual topology.
(iv) For the system (2.1), give an initial condition $\rho_0(\theta, \omega) = h(\theta)$, where h is an arbitrary measure on S^1. Then, corresponding solutions of (3.2),(3.3) satisfy $Z_j(t, \cdot) \in V_{1,0}$ for any $t \geq 0$ and $j = 1, 2, \cdots$.

Proof. (i) At first, we show that the set $i(V_{C,\alpha}) \subset \mathrm{Exp}'_+$ is equicontinuous. For any small $\varepsilon > 0$, we define a neighborhood $U = U(\varepsilon) \subset \mathrm{Exp}_+$ of the origin so that if $\phi \in U \cap \mathrm{Exp}_+(\gamma)$, then

$$\sup_{\mathrm{Im}(z)\geq 0} e^{-\gamma|z|}|\phi(z)| < \varepsilon D(\alpha, \gamma),$$

where $D(\alpha, \gamma)$ is a positive number to be determined. Then, for any $\phi \in U \cap \mathrm{Exp}_+(\gamma)$ and

$|\overline{\psi}\rangle \in i(V_{C,\alpha})$,

$$\begin{aligned}|\langle\phi\,|\,\overline{\psi}\rangle| &\leq \int_{\mathbf{R}} |\phi(\omega)|\cdot|\psi(\omega)|g(\omega)d\omega \\ &\leq \int_{\mathbf{R}} e^{-\gamma|\omega|}|\phi(\omega)|\cdot e^{-\alpha|\omega|}|\psi(\omega)|\cdot e^{(\alpha+\gamma)|\omega|}g(\omega)d\omega \\ &\leq \varepsilon CD(\alpha,\gamma)\int_{\mathbf{R}} e^{(\alpha+\gamma)|\omega|}g(\omega)d\omega.\end{aligned}$$

Since $g(\omega)$ decays faster than exponential functions, the integral $E(\alpha,\gamma) := \int_{\mathbf{R}} e^{(\alpha+\gamma)|\omega|}g(\omega)d\omega$ exists. If we put $D(\alpha,\gamma) = 1/CE(\alpha,\gamma)$, we obtain $|\langle\phi\,|\,\overline{\psi}\rangle| < \varepsilon$ for any $\phi \in U$ and $|\overline{\psi}\rangle \in i(V_{C,\alpha})$. This proves that $i(V_{C,\alpha})$ is an equicontinuous set. In particular, $i(V_{C,\alpha})$ is a bounded set of Exp_+' for both of the weak dual topology and the strong dual topology (see Prop.32.5 of Tréves [46]). Since Exp_+ is Montel, the weak dual topology, the projective topology and the strong dual topology coincide on the bounded set $i(V_{C,\alpha})$. Thus it is sufficient to prove (ii) for one of these topologies.

(ii) Obviously $V_{C,\alpha} \subset \mathrm{Exp}_+$ is connected (actually it is a convex set). Since the canonical inclusion i is continuous, $i(V_{C,\alpha})$ and $\overline{i(V_{C,\alpha})}$ are connected. Since the strong dual Exp_+' is Montel (Prop.5.4), every bounded set of Exp_+' is relatively compact, which proves that $\overline{i(V_{C,\alpha})}$ is compact. By the projective topology, $\overline{i(V_{C,\alpha})}$ is a metrizable space with the metric (5.42).

(iii) We show that $\Pi_n^{C,\alpha}$ is continuous at $0 \in i(V_{C,\alpha})$ with respect to the projective topology. Then, $\Pi_n^{C,\alpha}$ proves to be continuous on $i(V_{C,\alpha})$ because $\Pi_n^{C,\alpha}$ is linear. Since the projective topology is metrizable, it is sufficient to show that $\Pi_n^{C,\alpha}$ is sequentially continuous. Since every weakly convergent series in Exp_+' also converges with respect to the projective topology, it is sufficient to show that $\Pi_n^{C,\alpha}$ is sequentially continuous with respect to the weak dual topology. Take a sequence $\{\psi_m\}_{m=1}^{\infty} \subset V_{C,\alpha}$ such that $i(\psi_m) = |\overline{\psi_m}\rangle \to 0 \in i(V_{C,\alpha})$ as $m \to \infty$ with respect to the projective topology. Since the weak dual topology is weaker than the projective topology, it also converges with respect to the weak dual topology. In the proof of Thm.4.4, we have obtained Eq.(4.23) for each $\psi \in \mathrm{Exp}_+$. Putting $\psi = \psi_m$ in Eq.(4.23) yields

$$\begin{aligned}(e^{T_1 t}\phi, \overline{\psi}_m) &= \sum_{j=0}^{M} \frac{K}{2E_j} e^{\xi_j t}\langle\phi\,|\,\overline{\nu}_j\rangle\langle\psi_m\,|\,\overline{\nu}_j\rangle + \lim_{n\to\infty}\sum_{j=0}^{\#(n)} \frac{K}{2D_j} e^{\lambda_j t}\langle\phi\,|\,\mu_j\rangle\langle\psi_m\,|\,\mu_j\rangle \\ &\quad -\frac{1}{2\pi\sqrt{-1}}\lim_{n\to\infty}\int_{C_2+C_6} e^{\lambda t}F_0(\lambda)d\lambda - \frac{1}{2\pi\sqrt{-1}}\lim_{n\to\infty}\int_{C_3+C_4+C_5} e^{\lambda t}F_1(\lambda)d\lambda.\end{aligned}$$

In the proof of Thm.4.4, we proved that $\int_{C_2+C_6} e^{\lambda t}F_0(\lambda)d\lambda$ and $\int_{C_3+C_4+C_5} e^{\lambda t}F_1(\lambda)d\lambda$ tend to zero as $n \to \infty$. In particular, we showed that when $|\phi(z)| \leq C_1 e^{\beta_1|z|}$ and $|\psi(z)| \leq C_2 e^{\beta_2|z|}$ on the contour, these integrals have upper bounds determined by C_1, C_2, β_1 and β_2. Thus if $\psi_m \in V_{C,\alpha}$, these integrals have upper bounds which is independent of m, and it is easy to verify that they tend to zero uniformly in m. As a result, the infinite series in Eq.(4.18) converges uniformly in m, and we obtain

$$\begin{aligned}\lim_{m\to\infty}(e^{T_1 t}\phi, \overline{\psi}_m) &= \lim_{m\to\infty}\sum_{j=0}^{M}\frac{K}{2E_j}e^{\xi_j t}\langle \phi \,|\, \overline{v}_j\rangle\langle \psi_m \,|\, \overline{v}_j\rangle + \lim_{m\to\infty}\sum_{j=0}^{\infty}\frac{K}{2D_j}e^{\lambda_j t}\langle \phi \,|\, \mu_j\rangle\langle \psi_m \,|\, \mu_j\rangle \\ &= \sum_{j=0}^{M}\frac{K}{2E_j}e^{\xi_j t}\langle \phi \,|\, \overline{v}_j\rangle \lim_{m\to\infty}\langle \psi_m \,|\, \overline{v}_j\rangle + \sum_{j=0}^{\infty}\frac{K}{2D_j}e^{\lambda_j t}\langle \phi \,|\, \mu_j\rangle \lim_{m\to\infty}\langle \psi_m \,|\, \mu_j\rangle. \end{aligned} \tag{5.43}$$

Since $|\,\overline{\psi}_m\rangle \to 0$ with respect to the weak dual topology, the left hand side above is zero:

$$\lim_{m\to\infty}(e^{T_1 t}\phi, \overline{\psi}_m) = \lim_{m\to\infty}\langle e^{T_1 t}\phi \,|\, \overline{\psi}_m\rangle = 0.$$

Therefore, we obtain

$$0 = \sum_{j=0}^{M}\frac{K}{2\overline{E}_j}e^{\overline{\xi}_j t}\lim_{m\to\infty}\overline{\langle \psi_m \,|\, \overline{v}_j\rangle}\cdot |\,\overline{v}_j\rangle + \sum_{j=0}^{\infty}\frac{K}{2\overline{D}_j}e^{\overline{\lambda}_j t}\lim_{m\to\infty}\overline{\langle \psi_m \,|\, \mu_j\rangle}\cdot |\,\mu_j\rangle.$$

Now Thm.5.8 (ii) shows that (see also Eq.(5.36) below)

$$\lim_{m\to\infty}\langle \psi_m \,|\, \mu_j\rangle = 0 \tag{5.44}$$

for all j. This proves that

$$\lim_{m\to\infty}\Pi_j^{C,\alpha}|\,\overline{\psi}_m\rangle = \lim_{m\to\infty}\frac{K}{2\overline{D}_j}\overline{\langle \psi_m \,|\, \mu_j\rangle}\cdot |\,\mu_j\rangle = 0,$$

with respect to the weak dual topology. This implies that $\Pi_j^{C,\alpha}$ is sequentially continuous at $0 \in i(V_{C,\alpha})$ with respect to the weak dual topology. Since Exp_+ is Montel, a weakly convergent series in Exp_+' also converges with respect to the projective topology. Thus $\Pi_j^{C,\alpha}$ is continuous at $0 \in i(V_{C,\alpha})$ for all j with respect to the projective topology. Note that if $|\,\overline{\psi}_m\rangle \notin i(V_{C,\alpha})$, Eq.(5.44) may fail in general as is shown in Eq.(5.38).

(iv) To prove $Z_j \in V_{1,0}$, recall that Z_j is defined by Eq.(3.1). We want to estimate the analytic continuation of $Z_j(t,\omega)$ with respect to ω. Put $X(t) = e^{\sqrt{-1}x(t,0;\theta,\omega)}$. From Eq.(2.3), it turns out that X satisfies the equation

$$\left\{\begin{array}{l} \dfrac{dX}{dt} = \sqrt{-1}\omega X + \dfrac{K}{2}\eta(t) - \dfrac{K}{2}\overline{\eta(t)}X^2, \\ X(0) = e^{\sqrt{-1}\theta}. \end{array}\right.$$

Put $X(t) = \xi(t)e^{\sqrt{-1}p(t)}$, $\eta(t) = \zeta(t)e^{\sqrt{-1}q(t)}$ with $\xi, \zeta, p, q \in \mathbf{R}$. Then, the above equation is rewritten as

$$\frac{d\xi}{dt} + \sqrt{-1}\frac{dp}{dt}\xi = (\sqrt{-1}\mathrm{Re}(\omega) - \mathrm{Im}(\omega))\xi + \frac{K}{2}\zeta e^{\sqrt{-1}(q-p)} - \frac{K}{2}\zeta\xi^2 e^{-\sqrt{-1}(q-p)},$$

which yields

$$\frac{d\xi}{dt} = -\mathrm{Im}(\omega)\xi + \frac{K}{2}\zeta(1-\xi^2)\cos(q-p). \tag{5.45}$$

This equation shows that if $\mathrm{Im}(\omega) \geq 0$ and $|\xi| = 1$, then $d\xi/dt \leq 0$. Therefore, if the initial condition satisfies $|X(0)| \leq 1$, then $|X(t)| \leq 1$ for any $t > 0$ and any $\mathrm{Im}(\omega) \geq 0$. Thus the analytic continuation of $Z_j(t,\omega)$ to the upper half plane is estimated as

$$|Z_j(t,\omega)| \leq \int_{\mathbf{R}} |X(t)|^j h(\theta)d\theta \leq 1, \quad j = 1, 2, \cdots,$$

which means that $Z_j(t,\omega) \in V_{1,0}$ for every $t \geq 0$. ■

Eq.(5.33) gives the spectral decomposition for an element $|\overline{\psi}\rangle \in \overline{i(\mathrm{Exp}_+)}$. Now the spectral decomposition for $\xi \in \overline{i(V_{C,\alpha})}$ is obtained as follows. For $\xi \in \overline{i(V_{C,\alpha})}$, there exists a sequence $\{|\overline{\psi}_j\rangle\}_{j=1}^{\infty} \subset i(V_{C,\alpha})$ such that $|\overline{\psi}_j\rangle \to \xi$. Then, from the proof of Thm.5.10 (iii), we obtain

$$\begin{aligned}
(e^{T_1 t})^{\times}\xi &= \lim_{j\to\infty}(e^{T_1 t})^{\times}|\overline{\psi}_j\rangle \\
&= \lim_{j\to\infty}\sum_{n=0}^{\infty}\frac{K}{2\overline{D}_n}e^{\overline{\lambda}_n t}\overline{\langle\psi_j\,|\,\mu_n\rangle}\cdot|\,\mu_n\rangle \\
&= \sum_{n=0}^{\infty}\frac{K}{2\overline{D}_n}e^{\overline{\lambda}_n t}\lim_{j\to\infty}\overline{\langle\psi_j\,|\,\mu_n\rangle}\cdot|\,\mu_n\rangle,
\end{aligned} \tag{5.46}$$

and

$$\xi = \sum_{n=0}^{M}\frac{K}{2\overline{D}_n}\lim_{j\to\infty}\overline{\langle\psi_j\,|\,\mu_n\rangle}\cdot|\,\mu_n\rangle + \lim_{j\to\infty}\mathcal{R}_M[\psi_j]. \tag{5.47}$$

From Eq.(5.47), it turns out that the projection $\Pi_n : \mathrm{Exp}'_+ \to \mathrm{Exp}'_+$ is given by

$$\Pi_n\xi = \frac{K}{2\overline{D}_n}\lim_{j\to\infty}\overline{\langle\psi_j\,|\,\mu_n\rangle}\cdot|\,\mu_n\rangle, \quad \psi_j \in V_{C,\alpha},\ |\overline{\psi}_j\rangle \to \xi, \tag{5.48}$$

for any $\xi \in \overline{i(V_{C,\alpha})}$.

Define the *generalized center subspace* of T_1 to be

$$\mathbf{E}_c = \mathrm{span}\{\mu_n\,|\,\lambda_n \in \sqrt{-1}\mathbf{R}\} \subset \mathrm{Exp}'_+. \tag{5.49}$$

Proposition 5.11. $\mathbf{E}_c$ is a finite dimensional vector space.

Proof. We have to prove that the number of roots of Eq.(4.11) on the imaginary axis is finite. However, it is obvious because $D(\lambda)$ is analytic and tends to zero as $|\lambda| \to \infty$ along the imaginary axis. ■

Now we suppose that $K = K_c$. Then, the operator T_1 has resonance poles $\lambda_0, \cdots, \lambda_M$ on the imaginary axis. Let $\mu_0, \cdots, \mu_M$ be corresponding generalized eigenfunctions; that is, $\mathbf{E}_c = \mathrm{span}\{\mu_0, \cdots, \mu_M\}$. Let $\mathbf{E}_c^{\perp}$ be a complementary subspace of $\mathbf{E}_c$ in Exp'_+ including $\mu_{M+1}, \mu_{M+2}, \cdots$. In this case, Eq.(5.33) gives the decomposition with respect to the direct sum $\mathrm{Exp}'_+ = \mathbf{E}_c \oplus \mathbf{E}_c^{\perp}$. Let $\Pi_c : \mathrm{Exp}'_+ \to \mathbf{E}_c$ be the projection to $\mathbf{E}_c$ with respect to the direct sum $\mathrm{Exp}'_+ = \mathbf{E}_c \oplus \mathbf{E}_c^{\perp}$. Although $\mathbf{E}_c^{\perp}$ may not be unique, $\Pi_c | \overline{\psi} \rangle$ is uniquely determined for $| \overline{\psi} \rangle \in i(\mathrm{Exp}_+)$ because of Thm.5.8 (iii). The complementary subspace $\mathbf{E}_c^{\perp}$ including $\mu_{M+1}, \mu_{M+2}, \cdots$ is called the *stable subspace*. Eq.(5.30) shows that $\Pi_s (e^{T_1 t})^{\times} | \overline{\psi} \rangle$ decays exponentially as $t \to \infty$, where $\Pi_s = id - \Pi_c$ is the projection to $\mathbf{E}_c^{\perp}$.

Theorem 5.12. Let $\mu_0, \cdots, \mu_n$ be generalized eigenfunctions associated with the resonance poles on the imaginary axis. For any $| \overline{\psi} \rangle \in i(\mathrm{Exp}_+)$, the projection to the center subspace Π_c satisfies

(i)

$$\begin{aligned} \Pi_c | \overline{\psi} \rangle &= \Pi_0 | \overline{\psi} \rangle + \cdots + \Pi_n | \overline{\psi} \rangle \\ &= \frac{K}{2\overline{D}_0} \langle \overline{\psi \,|\, \mu_0} \rangle \cdot | \mu_0 \rangle + \cdots + \frac{K}{2\overline{D}_n} \langle \overline{\psi \,|\, \mu_n} \rangle \cdot | \mu_n \rangle. \end{aligned} \tag{5.50}$$

(ii)

$$\Pi_c T_1^{\times} | \overline{\psi} \rangle = T_1^{\times} \Pi_c | \overline{\psi} \rangle, \tag{5.51}$$

$$\Pi_c (e^{T_1 t})^{\times} | \overline{\psi} \rangle = (e^{T_1 t})^{\times} \Pi_c | \overline{\psi} \rangle. \tag{5.52}$$

(iii)

$$\Pi_c^2 = \Pi_c. \tag{5.53}$$

(iv) Π_c is continuous on $i(V_{C,\alpha})$, and Eq.(5.50) is continuously extended to any $\xi \in \overline{i(V_{C,\alpha})}$.

Proof. (i) and (iii) follows from the definition. (ii) is verified by using Eq.(5.7) and equalities $T_1^{\times} \mu_n = \overline{\lambda}_n \mu_n$, $(e^{T_1 t})^{\times} \mu_n = e^{\overline{\lambda}_n t} \mu_n$. (iv) follows from Thm.5.10. ■

For Theorem 5.12, it is sufficient to assume that resonance poles on the imaginary axis are poles of first order instead of the assumption (A4). If resonance poles on the left half plane are poles of higher order, the projection to the complement $\mathbf{E}_c^{\perp}$ may become complex, however, the projection to $\mathbf{E}_c$ is still given as above. For the bifurcation theory, it is sufficient to assume this fact.

5.3 Spectral theory on $F_p \subset L^2(\mathbf{R}, g(\omega) d\omega) \subset F'_p$

In the previous section, we have developed the spectral theory on Exp'_+, and the order parameter is completely resolved: it is decomposed into a countable set of exponential functions of the form $e^{\lambda_n t}$. However, for the bifurcation theory, it is sufficient to construct more coarse decompositions. To develop the center manifold theory, it is sufficient to express the semigroup as the sum of a neutrally stable term and an exponentially decaying term. To do so, we expect that it is sufficient to investigate resonance poles near the imaginary axis. For this purpose, let us introduce a space F_p, which consists of certain holomorphic functions defined near the real axis, and study the spectral theory of the semigroup $e^{T_1 t}$ on F_p. On this space, we can remove the technical assumption $K \notin \mathcal{K}_1$ because we do not use resonance poles near infinity.

For a fixed positive number $p > 0$, let $F_p(\beta, n)$ be the set of holomorphic functions on the region $\{z \in \mathbf{C} \,|\, -1/n \leq \mathrm{Im}(z) \leq p + 1/n\}$ such that the norm

$$||\phi||_{\beta,n} := \sup_{-1/n \leq \mathrm{Im}(z) \leq p+1/n} |\phi(z)| e^{-\beta |z|} \tag{5.54}$$

is finite. With this norm, $F_p(\beta, n)$ is a Banach space. Let $F_p(\beta)$ be their inductive limit with respect to $n = 1, 2, \cdots$

$$F_p(\beta) := \varinjlim_{n \geq 1} F_p(\beta, n) \ (= \bigcup_{n \geq 1} F_p(\beta, n)). \tag{5.55}$$

Next, define F_p to be their inductive limit with respect to $\beta = 0, 1, \cdots$

$$F_p := \varinjlim_{\beta \geq 0} F_p(\beta) \ (= \bigcup_{\beta \geq 0} F_p(\beta)). \tag{5.56}$$

Obviously $\mathrm{Exp}_+(\beta) \subset F_p(\beta)$ and $\mathrm{Exp}_+ \subset F_p \subset L^2(\mathbf{R}, g(\omega)d\omega)$. The analysis of the space F_p and the operator T_1 on F_p is done in a similar way to that of Exp_+. By the same way as the proof of Thm.5.3, we can show that $F_p(\beta)$ and F_p are complete Montel spaces. The topology on F_p is stronger than that of $L^2(\mathbf{R}, g(\omega)d\omega)$, and F_p is a dense subspace of $L^2(\mathbf{R}, g(\omega)d\omega)$. Thus the rigged Hilbert space $F_p \subset L^2(\mathbf{R}, g(\omega)d\omega) \subset F_p'$ is well defined. On the dual space F_p', three topologies are defined; the weak dual topology, the projective topology and the strong dual topology. Since F_p is Montel, on any bounded set of F_p', these topologies coincide with one another. In particular, a weakly convergent series on F_p' also converges with respect to the projective topology and the strong dual topology. The projective topology is metrizable with the distance d defined by Eq.(5.42) (we use the same notation as before).

As Prop.5.5, we can verify that T_1 is continuous as an operator on F_p, and it generates a semigroup $e^{T_1 t} : F_p \to F_p$. Unlike the semigroup on Exp_+, $e^{T_1 t}$ on F_p is not a holomorphic semigroup. However, we do not need this property because the spectral decomposition of $e^{T_1 t}$ on F_p shown below holds for every $t > 0$, although the decomposition (4.18) on Exp_+ holds for $t > t_0$.

The resonance poles λ_n and the generalized eigenfunctions μ_n are defined in the same way as in Sec.5.2, although only the resonance poles which lie on the strip $\{z \in \mathbf{C} \,|\, -p \leq \mathrm{Re}(z) \leq 0\}$ will be considered. It is easy to verify that μ_n is continuous on F_p, and thus $\mu_n \in F_p' \subset \mathrm{Exp}_+'$. Prop.5.6 and 5.7 also hold on F_p.

To develop the spectral theory on F_p, we start with the Laplace inversion formula like as Thm.4.4.

Proposition 5.13. Suppose that $0 < K \leq K_c$ and $g(\omega)$ satisfies the assumptions (A1) to (A3). For a positive number $0 < a \leq p$, suppose that there are no resonance poles on the line $\mathrm{Re}(z) = -a$. Then, for any $\phi, \psi \in F_p$, the equality

$$(e^{T_1 t}\phi, \overline{\psi}) = \sum_{j=0}^{n} R_j[t, \phi, \psi] e^{\lambda_j t} + \frac{e^{-at}}{2\pi} \mathrm{p.v.} \int_{-\infty}^{\infty} e^{\sqrt{-1}yt} F_1(-a + \sqrt{-1}y) dy \tag{5.57}$$



Fig. 10: The contour for the Laplace inversion formula.

holds for $t > 0$, where $\lambda_0, \cdots, \lambda_n$ are resonance poles of T_1 in the strip $\{z \,|\, -a < \mathrm{Re}(z) \leq 0\}$, $R_j[t, \phi, \psi]e^{\lambda_j t}$ are corresponding residues of the function $F_1(\lambda)e^{\lambda t}$, and where $F_1(\lambda)$ is defined by Eq.(4.5).

The contour used in the proof of this theorem does not pass close to infinity in the domain $\pi/2 + \delta < \arg(z) < 3\pi/2 - \delta$. Thus we need not introduce the set $\mathcal{K}$ of asymptotic values, and the proof is simpler than that of Thm.4.4.

Proof. The proof is similar to that of Thm.4.4. Let $d > 0$ be a sufficiently small number and $r > 0$ a sufficiently large number. Take paths C_1 to C_6 to be

$$\begin{aligned} C_1 &= \{d + \sqrt{-1}y \,|\, -r \leq y \leq r\}, \\ C_2 &= \{x + \sqrt{-1}r \,|\, 0 \leq x \leq d\}, \\ C_3 &= \{x + \sqrt{-1}r \,|\, -a \leq x \leq 0\}, \\ C_4 &= \{-a + \sqrt{-1}y \,|\, -r \leq y \leq r\}, \end{aligned}$$

and C_5 and C_6 are defined in a similar way to C_3 and C_2, respectively, see Fig.10.

Note that if r is sufficiently large, all resonance poles in the strip $\{z \,|\, -a < \mathrm{Re}(z) \leq 0\}$ are included in the closed curve $C = \sum_{i=1}^{6} C_i$ because there are no resonance poles on the area (4.28). In particular, the number of resonance poles in the strip is finite.

By the residue theorem, we have

$$2\pi\sqrt{-1} \sum_{j=0}^{n} R_j[t, \phi, \psi]e^{\lambda_j t} = \int_{C_6+C_1+C_2} e^{\lambda t} F_0(\lambda) d\lambda + \int_{C_3+C_4+C_5} e^{\lambda t} F_1(\lambda) d\lambda. \tag{5.58}$$

By using $|(\lambda - \sqrt{-1}\mathcal{M})^{-1}| \sim O(1/|\lambda|)$ and the assumption (A3), it is easy to verify that the integrals along C_2, C_3, C_5 and C_6 tend to zero as $r \to \infty$. Thus we obtain

$$\begin{aligned}(e^{T_1 t}\phi, \overline{\psi}) &= \sum_{j=0}^{n} R_j[t, \phi, \psi]e^{\lambda_j t} + \lim_{r\to\infty} \frac{1}{2\pi\sqrt{-1}} \int_{-a-\sqrt{-1}r}^{-a+\sqrt{-1}r} e^{\lambda t} F_1(\lambda) d\lambda \\ &= \sum_{j=0}^{n} R_j[t, \phi, \psi]e^{\lambda_j t} + \lim_{r\to\infty} \frac{e^{-at}}{2\pi} \int_{-r}^{r} e^{\sqrt{-1}yt} F_1(-a+\sqrt{-1}y) dy. \qquad (5.59)\end{aligned}$$

We have to show that the principle value of the integral in the right hand side exists. It is easy to verify that

$$((\lambda - \sqrt{-1}\mathcal{M})^{-1}\phi, \overline{\psi}) = \frac{1}{\lambda}\int_{\mathbf{R}} \phi(\omega)\psi(\omega)g(\omega)d\omega + O(\frac{1}{|\lambda|^2})$$

as $|\lambda| \to \infty$ outside the imaginary axis. Put $A := \int_{\mathbf{R}} \phi(\omega)\psi(\omega)g(\omega)d\omega$. Then, the definition of $F_1(\lambda)$ provides

$$F_1(-a+\sqrt{-1}y) \sim \frac{A}{\sqrt{-1}y} + O(1/y^2) \qquad (5.60)$$

as $y \to \pm\infty$. Therefore, for a sufficiently large $M > 0$, the integral is estimated as

$$\begin{aligned}&\int_{-\infty}^{\infty} e^{\sqrt{-1}yt} F_1(-a+\sqrt{-1}y) dy \\ = &\int_{-M}^{M} e^{\sqrt{-1}yt} F_1(-a+\sqrt{-1}y) dy + \int_{M}^{\infty} e^{\sqrt{-1}yt} \frac{A}{\sqrt{-1}y} dy + \int_{-\infty}^{-M} e^{\sqrt{-1}yt} \frac{A}{\sqrt{-1}y} dy + O(1/M) \\ = &\int_{-M}^{M} e^{\sqrt{-1}yt} F_1(-a+\sqrt{-1}y) dy + 2A\int_{M}^{\infty} \frac{\sin(yt)}{y} dy + O(1/M).\end{aligned}$$

It is well known that the integral $\int_{M}^{\infty} \sin(yt)/y dy$ exists for $t > 0$. This proves Prop.5.13. $\blacksquare$

Since $F_1(-a+\sqrt{-1}y)$ is smooth in y, the integral in Eq.(5.57) tends to zero as $t \to \infty$ by virtue of the Riemann-Lebesgue lemma. When $0 < K < K_c$, resonance poles $\lambda_0, \cdots, \lambda_n$ satisfy $\mathrm{Re}(\lambda_j) < 0$. This proves that $(e^{T_1 t}\phi, \overline{\psi})$ converges to zero as $t \to \infty$. In particular, the order parameter $\eta(t) = (e^{T_1 t}\phi, P_0)$ for the linearized system (3.4) tends to zero as $t \to \infty$. This fact was proved in Thm.4.4 under the assumption that $K \notin \mathcal{K}_1$, and now the assumption is removed.

By using the canonical inclusion $i : F_p \to F_p'$, the left hand side of Eq.(5.57) is rewritten as $\langle \phi \,|\, (e^{T_1 t})^{\times}\overline{\psi} \rangle$. On the other hand, by using the functional $\mu(\lambda)$ defined in (5.16), $F_1(\lambda)$ given by (4.5) is rewritten as

$$F_1(\lambda) = \langle \phi \,|\, \overline{\psi}\cdot\mu(\lambda) \rangle + \frac{K}{2}\frac{1}{G(\lambda)} \langle \phi \,|\, \mu(\lambda) \rangle \langle \psi \,|\, \mu(\lambda) \rangle, \qquad (5.61)$$

where the function $G(\lambda)$ is defined to be

$$G(\lambda) = 1 - \frac{K}{2}D(\lambda) - \pi K g(-\sqrt{-1}\lambda), \qquad (5.62)$$

and the functional $\overline{\psi} \cdot \mu(\lambda)$ is defined to be

$$\langle \phi \,|\, \overline{\psi} \cdot \mu(\lambda) \rangle = \int_{\mathbf{R}} \frac{1}{\lambda - \sqrt{-1}\omega} \phi(\omega)\psi(\omega)g(\omega)d\omega + 2\pi\phi(-\sqrt{-1}\lambda)\psi(-\sqrt{-1}\lambda)g(-\sqrt{-1}\lambda). \tag{5.63}$$

As a result, we obtain the spectral decomposition of $(e^{T_1 t})^{\times}$ on F_p.

Theorem 5.14. For Eq.(5.57), assume further that resonance poles $\lambda_0, \cdots, \lambda_n$ are poles of first order. Then, the equality

$$\begin{aligned} (e^{T_1 t})^{\times} | \overline{\psi} \rangle &= \sum_{j=0}^{n} \frac{K}{2\overline{D}_j} e^{\overline{\lambda}_j t} \overline{\langle \psi \,|\, \mu_j \rangle} \cdot | \mu_j \rangle + \frac{e^{-at}}{2\pi} \int_{-\infty}^{\infty} e^{-\sqrt{-1}yt} | \overline{\psi} \cdot \mu(-a + \sqrt{-1}y) \rangle dy \\ &\quad + \frac{Ke^{-at}}{4\pi} \int_{-\infty}^{\infty} e^{-\sqrt{-1}yt} \frac{1}{\overline{G(-a + \sqrt{-1}y)}} \overline{\langle \psi \,|\, \mu(-a + \sqrt{-1}y) \rangle} \cdot | \mu(-a + \sqrt{-1}y) \rangle dy \\ &= \sum_{j=0}^{n} \frac{K}{2\overline{D}_j} e^{\overline{\lambda}_j t} \overline{\langle \psi \,|\, \mu_j \rangle} \cdot | \mu_j \rangle + \frac{1}{\overline{2\pi\sqrt{-1}}} \int_{-a-\sqrt{-1}\infty}^{-a+\sqrt{-1}\infty} e^{\overline{\lambda} t} | \overline{\psi} \cdot \mu(\lambda) \rangle d\lambda \\ &\qquad + \frac{K}{\overline{4\pi\sqrt{-1}}} \int_{-a-\sqrt{-1}\infty}^{-a+\sqrt{-1}\infty} e^{\overline{\lambda} t} \frac{1}{\overline{G(\lambda)}} \overline{\langle \psi \,|\, \mu(\lambda) \rangle} \cdot | \mu(\lambda) \rangle d\lambda \end{aligned} \tag{5.64}$$

holds for any $\psi \in F_p$ and $t > 0$. In particular, since $(e^{T_1 t})^{\times} | \overline{\psi} \rangle$ is continuous at $t = 0$, taking the limit yields

$$\begin{aligned} | \overline{\psi} \rangle &= \sum_{j=0}^{n} \frac{K}{2\overline{D}_j} \overline{\langle \psi \,|\, \mu_j \rangle} \cdot | \mu_j \rangle + \frac{1}{\overline{2\pi\sqrt{-1}}} \lim_{t \to +0} \int_{-a-\sqrt{-1}\infty}^{-a+\sqrt{-1}\infty} e^{\overline{\lambda} t} | \overline{\psi} \cdot \mu(\lambda) \rangle d\lambda \\ &\qquad + \frac{K}{\overline{4\pi\sqrt{-1}}} \lim_{t \to +0} \int_{-a-\sqrt{-1}\infty}^{-a+\sqrt{-1}\infty} e^{\overline{\lambda} t} \frac{1}{\overline{G(\lambda)}} \overline{\langle \psi \,|\, \mu(\lambda) \rangle} \cdot | \mu(\lambda) \rangle d\lambda, \end{aligned} \tag{5.65}$$

which gives the spectral decomposition of $| \overline{\psi} \rangle \in i(F_p)$.

If $\lambda_0, \cdots, \lambda_n$ are poles of higher order, the first term in the right hand side becomes more complex involving polynomials in t. In this paper, we avoid such a situation for simplicity. Since Eq.(5.64) comes from Eq.(5.57), the integrals in the right hand side of Eq.(5.64) converge with respect to the weak dual topology on F_p'. However, since F_p is Montel, they also converge with respect to the strong dual and the projective topologies.

A similar result to Thm.5.8 is stated as follows.

Theorem 5.15. (i) A system of generalized functions

$$\mathcal{V} := \{ \mu_j, \, \mu(\lambda) \,|\, j = 0, \cdots, n, \, \lambda = -a + \sqrt{-1}y, y \in \mathbf{R} \} \tag{5.66}$$

is complete in the sense that if $\langle \psi \,|\, \mu_j \rangle = \langle \psi \,|\, \mu(\lambda) \rangle = 0$ for $j = 0, \cdots, n$ and $\lambda = -a + \sqrt{-1}y, y \in \mathbf{R}$, then $\psi = 0$.
(ii) Suppose that a functional $\xi(\lambda)$ parameterized by $\lambda \in \mathbf{C}$ satisfies

$$(e^{T_1 t})^{\times} \int_{-a-\sqrt{-1}\infty}^{-a+\sqrt{-1}\infty} \xi(\lambda) d\lambda = \int_{-a-\sqrt{-1}\infty}^{-a+\sqrt{-1}\infty} e^{\overline{\lambda} t} \xi(\lambda) d\lambda \in F_p'. \tag{5.67}$$

If $\sum_{j=0}^{n} a_j|\mu_j\rangle + \int_{-a-\sqrt{-1}\infty}^{-a+\sqrt{-1}\infty} \xi(\lambda)d\lambda = 0$ for $a_j \in \mathbf{C}$, then $a_j = 0\,(j = 0,\cdots,n)$ and $\int_{-a-\sqrt{-1}\infty}^{-a+\sqrt{-1}\infty} \xi(\lambda)d\lambda = 0$.
(iii) Let V be a complementary subspace of span$\{\mu_0,\cdots,\mu_n\}$ in F_p', which includes any functionals of the form $\int_{-a-\sqrt{-1}\infty}^{-a+\sqrt{-1}\infty} \xi(\lambda)d\lambda$ satisfying Eq.(5.67). Then, any $|\overline{\psi}\rangle \in i(F_p)$ is uniquely decomposed with respect to the direct sum span$\{\mu_0,\cdots,\mu_n\} \oplus V$ as Eq.(5.65), and this decomposition is independent of the choice of V.

Proof. (i) If $\langle\psi\,|\,\mu_j\rangle = \langle\psi\,|\,\mu(\lambda)\rangle = 0$, we obtain

$$\begin{aligned}\langle P_0\,|\,(e^{T_1 t})^{\times}\overline{\psi}\rangle &= \frac{1}{2\pi\sqrt{-1}}\int_{-a-\sqrt{-1}\infty}^{-a+\sqrt{-1}\infty} e^{\lambda t}\langle P_0\,|\,\overline{\psi}\cdot\mu(\lambda)\rangle d\lambda \\ &= \frac{1}{2\pi\sqrt{-1}}\int_{-a-\sqrt{-1}\infty}^{-a+\sqrt{-1}\infty} e^{\lambda t}\langle\psi\,|\,\mu(\lambda)\rangle d\lambda = 0.\end{aligned}$$

Differentiating in t and taking the limit $t \to 0$ yield $\langle T_1^n P_0\,|\,\overline{\psi}\rangle = 0$ for $n = 0,1,2,\cdots$. Since $T_1^n P_0$ is a polynomial of degree n in ω, we can prove by induction that $\langle\omega^n\,|\,\overline{\psi}\rangle = 0$ for $n = 0,1,2,\cdots$. Since the set of polynomials is dense in F_p, $\langle\phi\,|\,\overline{\psi}\rangle = 0$ for any $\phi \in F_p$. This proves $\psi = 0$. Parts (ii) and (iii) are proved in the same way as the proof of Thm.5.8. ■

Because of Thm.5.15, we can define the projection $\Pi_j : i(F_p) \to \text{span}\{\mu_j\}$ to the generalized eigenspace by Eq.(5.37). Now we suppose that $K = K_c$. Then, the operator T_1 has resonance poles on the imaginary axis, say $\lambda_0,\cdots,\lambda_n$. Take a positive number a so that all other resonance poles satisfy $\text{Re}(\lambda_j) < -a$, $j = n+1, n+2,\cdots$. In this case, Eq.(5.65) gives the decomposition with respect to the direct sum $F_p' = \mathbf{E}_c \oplus \mathbf{E}_c^{\perp}$. The projection Π_c to the generalized center subspace $\mathbf{E}_c$ is given by $\Pi_c = \Pi_0 + \cdots + \Pi_n$ as before. The complementary subspace $\mathbf{E}_c^{\perp}$ satisfying the condition given in Thm.5.15 (iii) is called the stable subspace. Eq.(5.64) shows that $\Pi_s(e^{T_1 t})^{\times}|\overline{\psi}\rangle$ decays faster than e^{-at}, where $\Pi_s = id - \Pi_c$ is the projection to $\mathbf{E}_c^{\perp}$.

Unfortunately, the projection Π_c to the generalized center subspace $\mathbf{E}_c$ is not continuous on $i(F_p)$, however, we can show that it is continuous on a subset of $i(F_p)$ like as Thm.5.10 (iii). Define a subset $V_p(C,\alpha) \subset F_p(\alpha)$ to be

$$V_p(C,\alpha) = \{\phi \in F_p(\alpha)\,|\,|\phi(z)|e^{-\alpha|z|} \leq C \text{ when } 0 \leq \text{Im}(z) \leq p\}. \tag{5.68}$$

Note that $V_{C,\alpha}$ defined by (5.39) is included in $V_p(C,\alpha)$. Then, we can prove that $\Pi_c : i(V_p(C,\alpha)) \to \mathbf{E}_c$ is continuous with respect to the projective topology on F_p. Indeed, take a sequence $\{\psi_m\}_{m=1}^{\infty} \subset V_p(C,\alpha)$ such that $i(\psi_m) = |\overline{\psi_m}\rangle \to 0$ as $m \to \infty$ with respect to the weak dual topology. Consider the formula (5.58) for each ψ_m. We can prove that the integrals along C_2 to C_6 converge as $r \to \infty$ uniformly in m. This fact yields

$$\begin{aligned} 0 &= \lim_{m\to\infty} \langle \phi \,|\, (e^{T_1 t})^{\times} \overline{\psi}_m \rangle \\ &= \lim_{m\to\infty} \sum_{j=0}^{n} \frac{K}{2D_j} e^{\lambda_j t} \langle \phi \,|\, \mu_j \rangle \langle \psi_m \,|\, \mu_j \rangle + \lim_{m\to\infty} \lim_{r\to\infty} \frac{e^{-at}}{2\pi} \int_{-r}^{r} e^{\sqrt{-1}yt} F_1(-a + \sqrt{-1}y) dy \\ &= \sum_{j=0}^{n} \frac{K}{2D_j} e^{\lambda_j t} \langle \phi \,|\, \mu_j \rangle \lim_{m\to\infty} \langle \psi_m \,|\, \mu_j \rangle + \lim_{r\to\infty} \frac{e^{-at}}{2\pi} \int_{-r}^{r} e^{\sqrt{-1}yt} \lim_{m\to\infty} F_1(-a + \sqrt{-1}y) dy. \end{aligned}$$

Taking the limit $t \to \infty$ proves the equality (5.44) for $j = 0, \cdots, n$, which means that $\Pi_0, \cdots, \Pi_n$ are continuous on $i(V_p(C, \alpha))$. The detailed proof is the same as that of Thm.5.10 (iii) and left to the reader. This also shows that the spectral decomposition holds for $\xi \in \overline{i(V_p(C, \alpha))}$ like as Eq.(5.47). These facts are summarized as follows (see also Thm.5.12).

Theorem 5.16. Let $\mu_0, \cdots, \mu_n$ be generalized eigenfunctions associated with the resonance poles on the imaginary axis. For any $|\overline{\psi}\rangle \in i(F_p)$, the projection Π_c to the center subspace satisfies

(i)

$$\begin{aligned} \Pi_c |\overline{\psi}\rangle &= \Pi_0 |\overline{\psi}\rangle + \cdots + \Pi_n |\overline{\psi}\rangle \\ &= \frac{K}{2\overline{D}_0} \overline{\langle \psi \,|\, \mu_0 \rangle} \cdot |\mu_0\rangle + \cdots + \frac{K}{2\overline{D}_n} \overline{\langle \psi \,|\, \mu_n \rangle} \cdot |\mu_n\rangle. \end{aligned} \tag{5.69}$$

(ii)

$$\Pi_c T_1^{\times} |\overline{\psi}\rangle = T_1^{\times} \Pi_c |\overline{\psi}\rangle, \tag{5.70}$$

$$\Pi_c (e^{T_1 t})^{\times} |\overline{\psi}\rangle = (e^{T_1 t})^{\times} \Pi_c |\overline{\psi}\rangle. \tag{5.71}$$

(iii)

$$\Pi_c^2 = \Pi_c. \tag{5.72}$$

(iv) Π_c is continuous on $i(V_p(C, \alpha))$ with respect to the projective topology. Thus Eq.(5.69) is continuously extended to the closure $\overline{i(V_p(C, \alpha))}$.

We have shown that Π_c is continuous on $i(V_p(C, \alpha))$. In general, there are no constants M such that the inequality $||\Pi_c \xi||^*_{\beta,n} \leq M ||\xi||^*_{\beta,n}$ holds for $\xi \in i(V_p(C, \alpha))$ because $i(V_p(C, \alpha))$ is not closed under the multiplication by a scalar. To obtain an estimate of $||\Pi_c \xi||^*_{\beta,n}$, we need an additional assumption for $g(\omega)$:

(A5) For fixed $p > 0$, $g(\omega + \sqrt{-1}p)/g(\omega) \in F_p$.

For example, the Gaussian distribution satisfies this assumption for any $p > 0$.

Next, for $C \geq 1$ and $\alpha \geq 0$, we define a subset $W_p(C, \alpha)$ of F_p to be

$$W_p(C, \alpha) := \{ \psi \in F_p(\alpha) \,|\, |\psi(z + \sqrt{-1}p)/\psi(z)| e^{-\alpha |z|} \leq C, \ \text{when } 0 \leq \mathrm{Im}(z) \leq p \}. \tag{5.73}$$

In other words, $\psi \in W_p(C, \alpha)$ if and only if $\psi(z + \sqrt{-1}p)/\psi(z) \in V_p(C, \alpha)$. Let $\overline{i(W_p(C, \alpha))} \subset F'_p$ be a closure of $i(W_p(C, \alpha)) = \{ i(\psi) \,|\, \psi \in W_p(C, \alpha) \}$ with respect to the weak dual topology. If $\xi \in \overline{i(W_p(C, \alpha))}$, then $k\xi \in \overline{i(W_p(C, \alpha))}$ for any $k \in \mathbf{C}$.

For elements in $\overline{i(W_p(C,\alpha))}$, let us estimate the norm of the projection Π_j.

Lemma 5.17. Suppose that $g(\omega)$ satisfies (A1) to (A3) and (A5).
(i) For each $\xi \in i(F_p)$, $||\xi||^*_{\beta,n}$ is bounded as $n \to \infty$.
(ii) For every $\beta = 0, 1, \cdots$ and $n = 1, 2, \cdots$, there exists a positive number $Q(\beta)$ such that the inequalities

$$||\xi||^*_{0,1} \leq ||\xi||^*_{\beta,n}, \quad ||\xi||^*_{\beta,n} \leq Q(\beta)||\xi||^*_{0,1} \tag{5.74}$$

hold for $\xi \in i(F_p)$ (this means that norms $||\cdot||^*_{\beta,n}$ are comparable [17]).
(iii) For $\mu(\lambda)$ defined by Eq.(5.16), the linear mapping $|\overline{\psi}\rangle \mapsto \langle \psi \,|\, \mu(\lambda)\rangle$ from $i(W_p(C,\alpha))$ into **C** is continuous with respect to the projective topology when $-p < \mathrm{Re}(\lambda) \leq 0$. In particular, if a resonance pole λ_j satisfies $-p < \mathrm{Re}(\lambda_j) \leq 0$, the corresponding projection Π_j is continuous on $i(W_p(C,\alpha))$.
(iv) For every $\beta = 0, 1, \cdots$ and $n = 1, 2, \cdots$, there exists a positive number $D_{C,\alpha,\beta,j}$ such that the inequality

$$||\Pi_j\xi||^*_{\beta,n} \leq D_{C,\alpha,\beta,j}||\xi||^*_{\beta,n} \tag{5.75}$$

holds for $\xi \in \overline{i(W_p(C,\alpha))}$.

In Thm.5.10, we have proved that Π_j is continuous on $i(V_p(C,\alpha))$. This lemma means that under the additional assumption (A5), Π_j proves to be continuous on $i(W_p(C,\alpha))$. Since the set $i(W_p(C,\alpha))$ is closed under the multiplication by a scalar, we can obtain the inequality above.

Proof. (i) $||\xi||^*_{\beta,n}$ has an upper bound

$$||\xi||^*_{\beta,n} := \sup_{||\phi||_{\beta,n}=1} \left| \int_{\mathbf{R}} \phi(\omega)\xi(\omega)g(\omega)d\omega \right| \leq \int_{\mathbf{R}} e^{\beta|\omega|}|\xi(\omega)|g(\omega)d\omega,$$

which is independent of $n = 1, 2, \cdots$.

(ii) The inequality $||\xi||^*_{0,1} \leq ||\xi||^*_{\beta,n}$ follows from the definition. It is easy to verify that the inclusion $F_p(0,1) \to F_p(\beta,n)$ is continuous. Thus its dual operator from $F_p(\beta,n)'$ into $F_p(0,1)'$ is continuous. This shows that there exists a positive number $Q(\beta)$ such that $||\xi||^*_{\beta,n} \leq Q(\beta)||\xi||^*_{0,1}$. Since the norm $||\cdot||^*_{\beta,n}$ is bounded as $n \to \infty$, we can take $Q(\beta)$ not to depend on $n = 1, 2, \cdots$.

(iii) Let $\{|\overline{\psi_m}\rangle\}_{m=1}^{\infty} \subset i(W_p(C,\alpha))$ be a sequence converging to zero as $m \to \infty$ with respect to the projective topology. By the definition of the projective topology, we have $||\,|\overline{\psi_m}\rangle||^*_{\beta,n} \to 0$ for every β and n. This means that $\langle f \,|\, \overline{\psi_m}\rangle \to 0$ uniformly in $f \in F_p(\beta,n)$ satisfying $||f||_{\beta,n} \leq C$ for each $C > 0$. Due to the part (i) of the lemma, $||\,|\overline{\psi_m}\rangle||^*_{\beta,n} \to 0$ uniformly in $n = 1, 2, \cdots$, which shows that $\langle f \,|\, \overline{\psi_m}\rangle \to 0$ uniformly in $f \in V_p(C,\beta)$. For a positive number $p > 0$ satisfying $-p < \mathrm{Re}(\lambda)$, $\langle \psi_m \,|\, \mu(\lambda)\rangle$ is given by

$$\begin{aligned}
\langle \psi_m \,|\, \mu(\lambda)\rangle &= \int_{\mathbf{R}} \frac{1}{\lambda - \sqrt{-1}\omega}\psi_m(\omega)g(\omega)d\omega + 2\pi\psi_m(-\sqrt{-1}\lambda)g(-\sqrt{-1}\lambda) \\
&= \int_{\mathbf{R}} \frac{1}{\lambda - \sqrt{-1}(\omega + \sqrt{-1}p)}\psi_m(\omega + \sqrt{-1}p)g(\omega + \sqrt{-1}p)d\omega \\
&= \int_{\mathbf{R}} \frac{1}{\lambda - \sqrt{-1}(\omega + \sqrt{-1}p)} \frac{\psi_m(\omega + \sqrt{-1}p)g(\omega + \sqrt{-1}p)}{\psi_m(\omega)g(\omega)}\psi_m(\omega)g(\omega)d\omega,
\end{aligned}$$

where we used the residue theorem. Putting

$$f_m(\omega) := \frac{1}{\lambda - \sqrt{-1}(\omega + \sqrt{-1}p)} \frac{\psi_m(\omega + \sqrt{-1}p) g(\omega + \sqrt{-1}p)}{\psi_m(\omega) g(\omega)}$$

provides $\langle \psi_m \,|\, \mu(\lambda) \rangle = \langle f_m \,|\, \overline{\psi_m} \rangle$. By the assumptions (A5) and $\psi_m \in W_p(C, \alpha)$, there exist $C' \geq 1$ and $\alpha' \geq 0$ such that $f_m \in V_p(C', \alpha')$ for every m. Therefore, $\langle \psi_m \,|\, \mu(\lambda) \rangle \to 0$ as $m \to 0$. Since the projective topology is metrizable, this implies that the mapping $|\overline{\psi}\rangle \mapsto \langle \psi \,|\, \mu(\lambda) \rangle$ is continuous.

(iv) Since Π_j is continuous on $i(W_p(C, \alpha))$ with respect to the metric d, for any $\varepsilon > 0$, there exists a number $\delta_{C,\alpha,j} > 0$ such that if $d(\xi, 0) \leq \delta_{C,\alpha,j}$, then $d(\Pi_j \xi, 0) \leq \varepsilon$ for $\xi \in i(W_p(C, \alpha))$. For $\varepsilon > 0$, take $\xi \in i(W_p(C, \alpha))$ and numbers β, n and $\hat{\delta}_{C,\alpha,\beta,j}$ such that $||\xi||^*_{\beta,n} \leq \hat{\delta}_{C,\alpha,\beta,j}$. We can take β and n sufficiently large and $\hat{\delta}_{C,\alpha,\beta,j}$ sufficiently small so that $d(\xi, 0) \leq \delta_{C,\alpha,j}$ holds, which implies $d(\Pi_j \xi, 0) \leq \varepsilon$. By the definition of d, this yields

$$||\Pi_j \xi||^*_{\beta,n} \leq \frac{2^n \kappa}{1 - 2^n \kappa}, \quad \kappa := \frac{2^\beta \varepsilon}{1 - 2^\beta \varepsilon}.$$

If $\xi \in i(W_p(C, \alpha))$, then $\xi / ||\xi||^*_{\beta,n} \in i(W_p(C, \alpha))$. Thus we obtain

$$\left|\left| \Pi_j \frac{\hat{\delta}_{C,\alpha,\beta,j} \xi}{||\xi||^*_{\beta,n}} \right|\right|^*_{\beta,n} \leq \frac{2^n \kappa}{1 - 2^n \kappa}, \tag{5.76}$$

for $\xi \in i(W_p(C, \alpha))$, which yields Eq.(5.75) by putting $D_{C,\alpha,\beta,j} = 2^n \kappa / (1 - 2^n \kappa) / \hat{\delta}_{C,\alpha,\beta,j}$. Because of the part (ii) of the lemma, the same inequality holds for any β and n with a suitable constant $D_{C,\alpha,\beta,j}$. Since the norm $|| \cdot ||^*_{\beta,n}$ is bounded as $n \to \infty$, we can take $D_{C,\alpha,\beta,j}$ not to depend on $n = 1, 2, \cdots$. The result is continuously extended to the closure $\overline{i(W_p(C, \alpha))}$. ■

Since the projection Π_j and the mapping $|\overline{\psi}\rangle \mapsto \langle \psi \,|\, \mu(\lambda) \rangle$ is continuous on $i(W_p(C, \alpha))$, the spectral decomposition (5.65) also holds for $\xi \in \overline{i(W_p(C, \alpha))}$. In particular, let $\{\psi_m\}_{m=1}^{\infty} \subset W_p(C, \alpha)$ be a sequence such that $|\overline{\psi_m}\rangle \to \xi \in \overline{i(W_p(C, \alpha))}$. Then, the projection of ξ is given by

$$\Pi_j \xi = \frac{K}{2\overline{D}_j} \lim_{m \to \infty} \overline{\langle \psi_m \,|\, \mu_j \rangle} \cdot |\, \mu_j \rangle. \tag{5.77}$$

Next purpose is to estimate norms of semigroups. At first, we suppose that $K < K_c$. In this case, there are no resonance poles on the imaginary axis and thus $\Pi_c = 0$.

Proposition 5.18. Suppose that $0 < K < K_c$ and $g(\omega)$ satisfies (A1) to (A3) and (A5). Take the constant p, which is used to define F_p and $W_p(C, \alpha)$, sufficiently small. For every $\beta = 0, 1, \cdots$ and $n = 1, 2, \cdots$, there exist positive numbers $M_{C,\alpha,\beta}$ and a such that the inequality

$$||(e^{T_j t})^{\times} \xi||^*_{\beta,n} \leq M_{C,\alpha,\beta} e^{-jat} ||\xi||^*_{\beta,n}, \quad j = 1, 2, \cdots \tag{5.78}$$

holds for $\xi \in \overline{i(W_p(C, \alpha))}$.

Proof. At first, we show the proposition for $j = 1$. Because of the assumption (A3), there are no resonance poles of T_1 on the area

$$\{z \in \mathbf{C} \,|\, |\arg(z) - \pi/2| \leq \delta,\ |\arg(z) - 3\pi/2| \leq \delta,\ |z| \geq r\},$$

if $r > 0$ is sufficiently large. Further, the left hand side of Eq.(4.11) is analytic. This shows that resonance poles can not accumulate on the imaginary axis. Therefore, there exist positive constants a and a' such that all resonance poles satisfy $\mathrm{Re}(\lambda_n) < -a' < -a$. For the definition of F_p and $W_p(C, \alpha)$, we choose the constant p so that $\mathrm{Re}(\lambda_n) < -p < -a'$. Then,

$$\frac{\|(e^{T_1 t})^\times \xi\|^*_{\beta,n}}{\|\xi\|^*_{\beta,n}} = e^{-at} \frac{\|(e^{(T_1+a)t})^\times \xi\|^*_{\beta,n}}{\|\xi\|^*_{\beta,n}},$$

and $(e^{(T_1+a)t})^\times \xi / \|\xi\|^*_{\beta,n}$ is given by

$$\begin{aligned}
(e^{(T_1+a)t})^\times \frac{\xi}{\|\xi\|^*_{\beta,n}} \quad &= \quad \frac{1}{2\pi\sqrt{-1}} \int^{-a'+\sqrt{-1}\infty}_{-a'-\sqrt{-1}\infty} e^{(\overline{\lambda}+a)t} \Big| \frac{\overline{\xi}}{\|\xi\|^*_{\beta,n}} \cdot \mu(\lambda) \Big\rangle d\lambda \\
&\quad + \frac{K}{4\pi\sqrt{-1}} \int^{-a'+\sqrt{-1}\infty}_{-a'-\sqrt{-1}\infty} e^{(\overline{\lambda}+a)t} \frac{1}{\overline{G(\lambda)}} \Big\langle \overline{\frac{\xi}{\|\xi\|^*_{\beta,n}} \,|\, \mu(\lambda)} \Big\rangle \cdot |\mu(\lambda)\rangle d\lambda,
\end{aligned}$$

for $\xi \in i(F_p)$. Since $-a' < -a$, this quantity tends to zero as $t \to \infty$ for each $\xi \in i(F_p)$. Because of Lemma 5.17, the right hand side is bounded with respect to the norm $\|\cdot\|^*_{\beta,n}$ uniformly in $\xi \in i(W_p(C, \alpha))$ and $t > 0$. This proves that there exists a positive constant $L_{C,\alpha,\beta}$ such that $\|(e^{T_1 t})^\times \xi\|^*_{\beta,n} \leq e^{-at} L_{C,\alpha,\beta} \|\xi\|^*_{\beta,n}$ for $\xi \in i(W_p(C, \alpha))$. Since the norm $\|\cdot\|^*_{\beta,n}$ is bounded as $n \to \infty$, we can take $L_{C,\alpha,\beta}$ not to depend on $n = 1, 2, \cdots$. The result is continuously extended to the closure $\overline{i(W_p(C, \alpha))}$.

Next, let us consider T_j for $j = 2, 3, \cdots$. Let $z_1, z_2, \cdots$ be poles of $g(\omega)$ on the upper half plane. Take $p > 0$ so small that $p < \mathrm{Im}(z_n)$ for every n. Then, Cauchy's theorem with the aid of the assumption (A3) proves that

$$\begin{aligned}
\langle \phi \,|\, (e^{T_j t})^\times \overline{\psi} \rangle \quad &= \quad (e^{j\sqrt{-1}\omega t} \phi, \overline{\psi}) \\
&= \quad \int_{\mathbf{R}} e^{j\sqrt{-1}\omega t} \phi(\omega) \psi(\omega) g(\omega) d\omega \\
&= \quad \int^{\sqrt{-1}p+\infty}_{\sqrt{-1}p-\infty} e^{j\sqrt{-1}\omega t} \phi(\omega) \psi(\omega) g(\omega) d\omega \\
&= \quad e^{-jpt} \int^{\infty}_{-\infty} e^{j\sqrt{-1}\omega t} \phi(\omega + \sqrt{-1}p) \psi(\omega + \sqrt{-1}p) g(\omega + \sqrt{-1}p) d\omega,
\end{aligned}$$

for any $\phi, \psi \in F_p$. Since $g(\omega + \sqrt{-1}p)$ decays faster than any exponential functions as $\omega \to \pm\infty$, the integral in the right hand side above exists. Now take a sequence $\{|\overline{\psi}_m\rangle\}^\infty_{m=1} \subset i(W_p(C, \alpha))$ converging to zero as $m \to \infty$ with respect to the projective topology. Then,

$$\begin{aligned}\langle \phi \,|\, (e^{(T_j+jp)t})^{\times}\overline{\psi_m}\rangle &= \int_{-\infty}^{\infty} e^{j\sqrt{-1}\omega t}\phi(\omega+\sqrt{-1}p)\psi_m(\omega+\sqrt{-1}p)g(\omega+\sqrt{-1}p)d\omega \\ &= \int_{-\infty}^{\infty} e^{j\sqrt{-1}\omega t}\phi(\omega+\sqrt{-1}p)\frac{\psi_m(\omega+\sqrt{-1}p)g(\omega+\sqrt{-1}p)}{\psi_m(\omega)g(\omega)}\psi_m(\omega)g(\omega)d\omega.\end{aligned}$$

Because of the assumptions, there exist $C' \geq 1$ and $\alpha' \geq 0$ such that the functions

$$e^{j\sqrt{-1}\omega t}\phi(\omega+\sqrt{-1}p)\frac{\psi_m(\omega+\sqrt{-1}p)g(\omega+\sqrt{-1}p)}{\psi_m(\omega)g(\omega)}$$

are included in $V_p(C', \alpha')$ for every m, j and $t > 0$. This proves that $\langle \phi \,|\, (e^{(T_j+jp)t})^{\times}\overline{\psi_m}\rangle$ tends to zero as $m \to \infty$ uniformly in $j = 2, 3, \cdots$ and $t > 0$. Therefore, the operator $(e^{(T_j+jp)t})^{\times}$ is continuous on $i(W_p(C, \alpha))$ uniformly in $j = 2, 3, \cdots$ and $t > 0$. By the same way as the proof of Lemma 5.17 (iii), we can show that there exists a positive number $N_{C,\alpha,\beta}$ such that $||(e^{(T_j+jp)t})^{\times}\xi||^*_{\beta,n} \leq N_{C,\alpha,\beta}||\xi||^*_{\beta,n}$ for $\xi \in \overline{i(W_p(C, \alpha))}$. Choose the constant $p > 0$ so that it satisfies both conditions $\mathrm{Re}(\lambda_n) < -p < -a'$ and $p < \mathrm{Im}(z_n)$ above. Then, putting $M_{C,\alpha,\beta} = \max\{L_{C,\alpha,\beta}, N_{C,\alpha,\beta}\}$ yields the desired result. ■

Next, we suppose that $K = K_c$. In this case, there exist resonance poles on the imaginary axis and $\Pi_c \neq 0$. Then, we can prove the following proposition.

Proposition 5.19. Suppose that $K = K_c$, $g(\omega)$ satisfies the assumptions (A1) to (A3) and (A5), and that resonance poles on the imaginary axis are poles of first order. Take the constant $p > 0$ sufficiently small. Then, for every $\beta = 0, 1, \cdots$ and $n = 1, 2, \cdots$, there exist positive constants $L_{C,\alpha,\beta}$, $M_{C,\alpha,\beta}$, $N_{C,\alpha,\beta}$ and a such that the inequalities

$$||(e^{T_1 t})^{\times}\Pi_c\xi||^*_{\beta,n} \leq L_{C,\alpha,\beta}||\xi||^*_{\beta,n}, \tag{5.79}$$

$$||(e^{T_1 t})^{\times}\Pi_s\xi||^*_{\beta,n} \leq M_{C,\alpha,\beta}e^{-at}||\xi||^*_{\beta,n}, \tag{5.80}$$

hold for $\xi \in \overline{i(W_p(C, \alpha))}$, and the inequality

$$||(e^{T_j t})^{\times}\xi||^*_{\beta,n} \leq N_{C,\alpha,\beta}e^{-jat}||\xi||^*_{\beta,n}, \quad j = 2, 3, \cdots, \tag{5.81}$$

holds for $\xi \in \overline{i(W_p(C, \alpha))}$.

Proof. Let $\lambda_0, \cdots, \lambda_n$ be resonance poles on the imaginary axis. For $\xi \in i(W_p(C, \alpha))$, $(e^{T_1 t})^{\times}\Pi_c\xi$ is calculated as

$$(e^{T_1 t})^{\times}\Pi_c\xi = \sum_{j=0}^{n}\frac{K}{2\overline{D}_j}\overline{\langle \xi \,|\, \mu_j\rangle}\cdot(e^{T_1 t})^{\times}|\,\mu_j\rangle = \sum_{j=0}^{n}\frac{K}{2\overline{D}_j}\overline{\langle \xi \,|\, \mu_j\rangle}\cdot e^{\overline{\lambda}_j t}|\,\mu_j\rangle.$$

Since $\lambda_j \in \sqrt{-1}\mathbf{R}$, we obtain

$$||\,(e^{T_1 t})^{\times}\Pi_c\xi\,||^*_{\beta,n} \leq \sum_{j=0}^{n}||\,\frac{K}{2\overline{D}_j}\overline{\langle \xi \,|\, \mu_j\rangle}\cdot|\,\mu_j\rangle\,||^*_{\beta,n} = \sum_{j=0}^{n}||\,\Pi_j\xi\,||^*_{\beta,n} \leq \sum_{j=0}^{n}D_{C,\alpha,\beta,j}||\xi||^*_{\beta,n}.$$

This provides Eq.(5.79) for $\xi \in i(W_p(C,\alpha))$ with $L_{C,\alpha,\beta} = \sum_{j=0}^{n} D_{C,\alpha,\beta,j}$. The result is continuously extended for $\xi \in \overline{i(W_p(C,\alpha))}$. The proofs of Eq.(5.80) and (5.81) are the same as that of Prop 5.18 with the aid of the fact that $\Pi_s = id - \Pi_c$ is continuous on $i(W_p(C,\alpha))$. ■

Note that the constant $a > 0$ in Prop.5.19, which determines the decay rate of $(e^{T_1 t})^\times$, is chosen so that all resonance poles λ_j on the left half plane satisfy $\mathrm{Re}(\lambda_j) < -a$, and all poles z_j of $g(\omega)$ on the upper half plane satisfy $\mathrm{Im}(z_j) > a$. If resonance poles on the imaginary axis are poles of higher order, $\|(e^{T_1 t})^\times \Pi_c \xi\|^*_{\beta,n}$ may diverge as $t \to \infty$ with a polynomial rate.

The next proposition is used to show that the inequalities (5.78),(5.79),(5.80),(5.81) are applicable to solutions of the Kuramoto model.

Proposition 5.20. (i) For any $C \geq 1$, $i(V_{2p}(C,0)) \subset \overline{i(W_p(3,0))}$.
(ii) Put $V = \bigcup_{C\geq 1} V_{2p}(C,0)$. Then, the generalized center subspace $\mathbf{E}_c$ is included in $\overline{i(V)}$;

$$\mathbf{E}_c \subset \overline{i(V)} \subset \overline{i(W_p(3,0))} \subset F_p'. \tag{5.82}$$

Proof. (i) If a function $\psi \in V_{2p}(C,0)$ has zeros on the region $0 \leq \mathrm{Im}(z) \leq p$, $\psi \notin W_p(C,\alpha)$ for any C and α. To prove that $|\overline{\psi}\rangle = i(\psi) \in \overline{i(W_p(3,0))}$, let us perturb the function $\psi \in V_{2p}(C,0)$. For $n = 1,2,\cdots$, put

$$\tilde{\psi}(\omega) = \psi(\omega) + 2Ce^{\sqrt{-1}n\omega + np}. \tag{5.83}$$

Since $\psi \in V_{2p}(C,0)$, $\psi(\omega + \sqrt{-1}p) \in V_p(C,0)$. For $0 \leq \mathrm{Im}(\omega) \leq p$, we have

$$\left| \frac{\tilde{\psi}(\omega + \sqrt{-1}p)}{\tilde{\psi}(\omega)} \right| = \left| \frac{\psi(\omega + \sqrt{-1}p) + 2Ce^{\sqrt{-1}n\omega}}{\psi(\omega) + 2Ce^{\sqrt{-1}n\omega + np}} \right| \leq \frac{C + 2Ce^{-n\mathrm{Im}(\omega)}}{2Ce^{n(p-\mathrm{Im}(\omega))} - C} \leq 3,$$

which implies $\tilde{\psi} \in W_p(3,0)$. It is easy to verify that $2Ce^{\sqrt{-1}n\omega + np} \to 0$ as $n \to \infty$ with respect to the weak dual topology. Therefore, $i(\tilde{\psi}) \to i(\psi) \in \overline{i(W_p(3,0))}$ as $n \to \infty$ for any $C \geq 1$.

(ii) Put $v_\lambda(\omega) = 1/(\lambda - \sqrt{-1}\omega)$. Let $\lambda_j = \sqrt{-1}y_j$ be a resonance pole on the imaginary axis. By the definition, the corresponding generalized eigenfunction μ_j is given by

$$\mu_j = \lim_{x\to+0} i(v_{x+\sqrt{-1}y_j}(\omega)) = \lim_{x\to+0} |\,\overline{v_{x+\sqrt{-1}y_j}(\omega)}\rangle, \tag{5.84}$$

where the limit is taken with respect to the weak dual topology. It is easy to verify that $v_\lambda(\omega) \in V$ for $\mathrm{Re}(\lambda) > 0$. This implies that $\mu_j \in \overline{i(V)}$ and thus the generalized center subspace $\mathbf{E}_c$ is included in $\overline{i(V)}$. ■

Note that $\overline{i(V)}$ is a closed subspace of F_p'. If $\psi \in V$, $i(\psi) = |\overline{\psi}\rangle$ satisfies inequalities (5.78), (5.79), (5.80),(5.81), in which the constants depend only on β. Since $\mathbf{E}_c \subset \overline{i(V)}$, the generalized eigenfunctions in $\mathbf{E}_c$ also satisfy the inequalities with the same constants. The set $\overline{i(V)}$ has all properties for developing a bifurcation theory: it is a metric space including all solutions of the Kuramoto model and the generalized center subspace. The projection Π_c is continuous on $\overline{i(V)}$. The semigroup $(e^{T_1 t})^\times$ admits the spectral decomposition on it, and norms of the semigroups $(e^{T_j t})^\times$ satisfy the appropriate inequalities. By using these properties, we will prove the existence of center manifolds in Section 7.

Remark. Since the generalized eigenfunction μ_j is given by Eq.(5.84) and since $\mathbf{E}_c \subset \overline{i(V)}$, Eq.(5.77) provides

$$\Pi_j \mu_j = \frac{K}{2\overline{D}_j} \lim_{x\to+0} \overline{\langle v_{x+\sqrt{-1}y_j}(\omega)\,|\,\mu_j\rangle} \cdot |\,\mu_j\rangle. \tag{5.85}$$

Since $\Pi_j \mu_i = \delta_{ij}\mu_i$, we obtain

$$\lim_{x\to+0} \langle v_{x+\sqrt{-1}y_j}(\omega)\,|\,\mu_i\rangle = \frac{2D_i}{K}\delta_{ij} = \begin{cases} \dfrac{2D_i}{K} & (i=j), \\ 0 & (i\neq j), \end{cases} \tag{5.86}$$

for $i, j = 0, 1, \cdots, n$. This is formally written as $\langle \mu_j\,|\,\mu_i\rangle = \frac{2D_i}{K}\delta_{ij}$, which means that a system of generalized eigenfunctions $\{\mu_0, \mu_1, \cdots, \mu_n\}$ forms a complete "orthogonal" basis of $\mathbf{E}_c$.

5.4 Spectral theory on $(H_+, L^2(\mathbf{R}, g(\omega)d\omega), H_+')$

In this subsection, we suppose that $g(\omega)$ is a rational function. Let H_+ be a Banach space of bounded holomorphic functions on the real axis and the upper half plane (see Sec.4.3). In this case, H_+ is not a dense subspace of $L^2(\mathbf{R}, g(\omega)d\omega)$, and thus the triplet $(H_+, L^2(\mathbf{R}, g(\omega)d\omega), H_+')$ is a degenerate rigged Hilbert space. Indeed, recall that a triplet is degenerate if and only if the canonical inclusion is not injective.

Proposition 5.21. The canonical inclusion $i : H_+ \to H_+'$ is a finite dimensional operator ; that is, $i(H_+) \subset H_+'$ is a finite dimensional vector space.

Proof. By the definition,

$$i(\psi)(\phi) = \langle \phi\,|\,\overline{\psi}\rangle = (\phi, \overline{\psi}) = \int_{\mathbf{R}} \phi(\omega)\psi(\omega)g(\omega)d\omega,$$

for $\phi, \psi \in H_+$. Let $z_1, \cdots, z_n$ be poles of $g(\omega)$ on the upper half plane. By the residue theorem, we obtain

$$\begin{aligned} & \int_{\mathbf{R}} \phi(\omega)\psi(\omega)g(\omega)d\omega + \int_0^\pi \phi(re^{\sqrt{-1}\theta})\psi(re^{\sqrt{-1}\theta})g(re^{\sqrt{-1}\theta})\sqrt{-1}re^{\sqrt{-1}\theta}d\theta \\ = \;& 2\pi\sqrt{-1}\sum_{j=1}^n \mathrm{Res}(z_j), \end{aligned}$$

where $r > 0$ is a sufficiently large number and $\mathrm{Res}(z_j)$ denotes the residue of $\phi(\omega)\psi(\omega)g(\omega)$ at z_j. Since $g(\omega)$ is a rational function which is integrable on the real axis, the degree of the denominator is at least two greater than the degree of the numerator : $g(\omega) \sim O(1/|\omega|^2)$ as $|\omega| \to \infty$. Since $\phi, \psi \in H_+$ is bounded on the upper half plane, we obtain

$$\langle \phi \,|\, \overline{\psi} \rangle = 2\pi \sqrt{-1} \sum_{j=1}^{n} \mathrm{Res}(z_j), \tag{5.87}$$

as $r \to \infty$. This means that the action of $|\,\overline{\psi}\rangle \in i(H_+)$ on H_+ is determined by the values of $\psi(\omega)$ and its derivatives at $z_1, \cdots, z_n$. In particular, if the denominator of g is of degree M, then $i(H_+) \simeq \mathbf{C}^M$. ■

Since $i(H_+)$ is of finite dimensional, the semigroup $(e^{T_1 t})^\times$ restricted to $i(H_+)$ is a finite dimensional operator. This is the reason that Eq.(4.30) consists of a finite sum. In what follows, we suppose the assumption (A4). Then Eq.(4.30) is rewritten as

$$(e^{T_1 t}\phi, \overline{\psi}) = \langle \phi \,|\, (e^{T_1 t})^\times \overline{\psi} \rangle = \sum_{n=0}^{M} \frac{K}{2D_n} e^{\lambda_n t} \langle \phi \,|\, \mu_n \rangle \langle \psi \,|\, \mu_n \rangle, \tag{5.88}$$

where definitions of D_n and $|\,\mu_n\rangle$ are the same as those in previous sections. Now we have obtained the following theorem.

Theorem 5.22. Suppose that $g(\omega)$ is a rational function with the assumption (A4). For any $\psi \in H_+$, the equalities

$$(e^{T_1 t})^\times |\,\overline{\psi}\rangle = \sum_{n=0}^{M} \frac{K}{2\overline{D}_n} e^{\overline{\lambda}_n t} \overline{\langle \psi \,|\, \mu_n \rangle} \cdot |\,\mu_n\rangle, \tag{5.89}$$

$$|\,\overline{\psi}\rangle = \sum_{n=0}^{M} \frac{K}{2\overline{D}_n} \overline{\langle \psi \,|\, \mu_n \rangle} \cdot |\,\mu_n\rangle, \tag{5.90}$$

hold. In particular, a system of generalized eigenfunctions $\{\mu_n\}_{n=0}^{M}$ forms a base of $i(H_+)$.

The projection $\Pi_n : i(H_+) \to \mathrm{span}\{\mu_n\} \subset H'_+$ is defined to be

$$\Pi_n |\,\overline{\psi}\rangle = \frac{K}{2\overline{D}_n} \overline{\langle \psi \,|\, \mu_n \rangle} \cdot |\,\mu_n\rangle, \quad n = 0, \cdots, M \tag{5.91}$$

as before. Since $i(H_+)$ is a finite dimensional vector space, Π_n is continuous on the whole space. Note that solutions $Z_1, Z_2, \cdots$ of the Kuramoto model are included in H_+ (we have proved that $Z_j \in V_{1,0}$ in Thm.5.10 (iv)). Thus the bifurcation problem of the Kuramoto model is reduced to the bifurcation theory on a finite dimensional space, and the usual center manifold theory is applicable.

6 Nonlinear stability

Before going to the bifurcation theory, let us consider the nonlinear stability of the de-synchronous state. In Sec.4 and Sec.5.3, we proved that the order parameter $\eta(t) \equiv 0$ is linearly stable when $0 < K < K_c$; that is, the asymptotic stability of $\eta(t) \equiv 0$ is proved for the linearized system (3.4). For a system on an infinite dimensional space, in general, the linear stability does not imply the nonlinear stability. Infinitesimally small nonlinear terms may change the stability of fixed points. In this section, we show that the de-synchronous state $Z_j(t) \equiv 0\,(j = 1, 2, \cdots)$ (which corresponds to $\rho_t \equiv 1/2\pi$) is locally stable with respect to a suitable topology when $0 < K < K_c$. In particular, the order parameter proves to decay to zero as $t \to \infty$ without neglecting the nonlinear terms.

Recall that the continuous model (2.1) is rewritten as Eqs.(3.2),(3.3) by putting $Z_j(t, \omega) = \int_0^{2\pi} e^{\sqrt{-1}j\theta}\rho_t(\theta, \omega)d\theta$ with the initial condition

$$Z_j(0, \omega) = \int_0^{2\pi} e^{\sqrt{-1}j\theta} h(\theta)d\theta := h_j \in \mathbf{C}. \tag{6.1}$$

We need not suppose that $h(\theta)$ is a usual function. It may be a probability measure on S^1. By using the canonical inclusion, Eqs.(3.2),(3.3) are rewritten as a system of evolution equations on $\prod_{j=1}^{\infty} F'_p$ of the form

$$\begin{cases} \dfrac{d}{dt}|\,\overline{Z}_1\rangle = T_1^{\times}|\,\overline{Z}_1\rangle - \dfrac{K}{2}\langle P_0\,|\,\overline{Z}_1\rangle|\,\overline{Z}_2\rangle, \\ \dfrac{d}{dt}|\,\overline{Z}_j\rangle = T_j^{\times}|\,\overline{Z}_j\rangle + \dfrac{jK}{2}\left(\overline{\langle P_0\,|\,\overline{Z}_1\rangle}|\,\overline{Z}_{j-1}\rangle - \langle P_0\,|\,\overline{Z}_1\rangle|\,\overline{Z}_{j+1}\rangle\right), \quad j = 2, 3, \cdots, \\ |\,\overline{Z_j(0,\,\cdot\,)}\rangle = \overline{h}_j|\,P_0\rangle, \end{cases} \tag{6.2}$$

where linear operators T_j are defined to be

$$T_1\phi(\omega) = (\sqrt{-1}\mathcal{M} + \frac{K}{2}\mathcal{P})\phi(\omega) = \sqrt{-1}\omega\phi(\omega) + \frac{K}{2}\langle P_0\,|\,\overline{\phi}\rangle P_0(\omega), \tag{6.3}$$

and

$$T_j\phi(\omega) = \sqrt{-1}j\mathcal{M}\phi(\omega) = \sqrt{-1}j\omega\phi(\omega), \tag{6.4}$$

for $j = 2, 3, \cdots$. $T_j^{\times}$ are their dual operators. The main theorem in this section is stated as follows.

Theorem 6.1 (local stability of the de-synchronous state). Suppose that $0 < K < K_c$ and $g(\omega)$ satisfies the assumptions (A1) to (A3) and (A5). Then, there exists a positive constant δ_β such that if the initial condition $h(\theta)$ of the initial value problem (2.1) satisfies

$$|h_j| = \left|\int_0^{2\pi} e^{j\sqrt{-1}\theta} h(\theta)d\theta\right| \leq \delta_\beta, \quad j = 1, 2, \cdots, \tag{6.5}$$

then the quantities

$$(Z_j, \overline{\phi}) = \int_0^{2\pi}\int_{\mathbf{R}} e^{\sqrt{-1}j\theta}\phi(\omega)\rho_t(\theta, \omega)d\omega d\theta$$

tend to zero as $t \to \infty$ for every $\phi \in F_p(\beta)$ uniformly in $j = 1, 2, \cdots$. In particular, the order parameter $\eta(t) = (Z_1, P_0)$ tends to zero as $t \to \infty$.

This theorem means that the trivial solution $Z_j \equiv 0$ of (6.2) is locally stable with respect to the weak dual topology on F'_p. In general, $\delta_\beta \to 0$ as $\beta \to \infty$. One of the reasons is that the norm $\|\cdot\|^*_{\beta,n}$ goes to infinity as $\beta \to \infty$. For the case $g(\omega)$ is a rational function, we can show the same statement : $(Z_j, \overline{\phi})$ tends to zero as $t \to \infty$ for every $\phi \in H_+$ if the initial condition satisfies (6.5), in which δ_β is independent of β.

Proof of Thm.6.1. Since we have Prop.5.18, the proof is done in a similar manner to the proof of the local stability of fixed points of finite dimensional systems. Eq.(6.2) provides

$$\begin{cases} |\,\overline{Z_1(t,\cdot)}\rangle = (e^{T_1(t-t_0)})^\times |\,\overline{Z_1(t_0,\cdot)}\rangle - \dfrac{K}{2}\displaystyle\int^t_{t_0} \langle P_0\,|\,\overline{Z_1(s,\cdot)}\rangle (e^{T_1(t-s)})^\times |\,\overline{Z_2(s,\cdot)}\rangle ds, \\ |\,\overline{Z_j(t,\cdot)}\rangle = (e^{T_j(t-t_0)})^\times |\,\overline{Z_j(t_0,\cdot)}\rangle + \dfrac{jK}{2}\displaystyle\int^t_{t_0} \Big(\langle \overline{P_0\,|\,\overline{Z_1(s,\cdot)}}\rangle (e^{T_j(t-s)})^\times |\,\overline{Z_{j-1}(s,\cdot)}\rangle \\ \qquad\qquad - \langle P_0\,|\,\overline{Z_1(s,\cdot)}\rangle (e^{T_j(t-s)})^\times |\,\overline{Z_{j+1}(s,\cdot)}\rangle \Big) ds, \end{cases} \tag{6.6}$$

for $0 \leq t_0 < t$. Since $|\,\overline{Z}_j\rangle \in i(V_{1,0}) \subset i(V_{2p}(1,0)) \subset i(W_p(3,0))$ for every $t > 0$ and $j = 1, 2, \cdots$, Prop.5.18 is applied to show that there exists $M_{3,0,\beta} = M_\beta > 0$ such that

$$\begin{cases} \|\,|\,\overline{Z_1(t,\cdot)}\rangle\|^*_{\beta,n} \leq M_\beta e^{-a(t-t_0)} \|\,|\,\overline{Z_1(t_0,\cdot)}\rangle\|^*_{\beta,n} + \dfrac{K}{2}\displaystyle\int^t_{t_0} M_\beta e^{-a(t-s)} \|\,|\,\overline{Z_1(s,\cdot)}\rangle\|^*_{\beta,n} \cdot \|\,|\,\overline{Z_2(s,\cdot)}\rangle\|^*_{\beta,n} ds, \\ \|\,|\,\overline{Z_j(t,\cdot)}\rangle\|^*_{\beta,n} \leq M_\beta e^{-ja(t-t_0)} \|\,|\,\overline{Z_j(t_0,\cdot)}\rangle\|^*_{\beta,n} + \dfrac{jK}{2}\displaystyle\int^t_{t_0} M_\beta e^{-ja(t-s)} \|\,|\,\overline{Z_1(s,\cdot)}\rangle\|^*_{\beta,n} \cdot \\ \qquad\qquad \Big(\|\,|\,\overline{Z_{j-1}(s,\cdot)}\rangle\|^*_{\beta,n} + \|\,|\,\overline{Z_{j+1}(s,\cdot)}\rangle\|^*_{\beta,n} \Big) ds. \end{cases} \tag{6.7}$$

Take a small constant $\delta_\beta > 0$ such that $h_j = \int^{2\pi}_0 e^{j\sqrt{-1}\theta} h(\theta) d\theta$ satisfies Eq.(6.5). Let us show that there exist $t_0 > 0$ and $N_\beta > 0$ such that

$$\|\,|\,\overline{Z_j(t,\cdot)}\rangle\|^*_{\beta,n} \leq \delta_\beta N_\beta \tag{6.8}$$

holds for any $t > t_0$ and $j = 1, 2, \cdots$. Indeed, $\|\,|\,\overline{Z_j(t,\cdot)}\rangle\|^*_{\beta,n}$ is calculated as

$$\begin{aligned} \|\,|\,\overline{Z_j(t,\cdot)}\rangle\|^*_{\beta,n} &= \sup_{\|\phi\|_{\beta,n}=1} \Big| \int_{\mathbf{R}} \phi(\omega) Z_j(t,\omega) g(\omega) d\omega \Big| \\ &= \sup_{\|\phi\|_{\beta,n}=1} \Big| \int_{\mathbf{R}} \phi(\omega) \int^{2\pi}_0 e^{\sqrt{-1}jx(t,0;\theta,\omega)} h(\theta) d\theta\, g(\omega) d\omega \Big|. \end{aligned}$$

Let

$$e^{\sqrt{-1}jx(t,0;\theta,\omega)} = \sum^{\infty}_{n=-\infty} a_{jn}(t,\omega) e^{\sqrt{-1}n\theta}$$

be the Fourier expansion of $e^{\sqrt{-1}jx(t,0;\theta,\omega)}$. Then, we obtain

$$\begin{aligned} \|\,|\,\overline{Z_j(t,\cdot)}\rangle\|^*_{\beta,n} &= \sup_{\|\phi\|_{\beta,n}=1} \Big| \sum^{\infty}_{n=-\infty} \int_{\mathbf{R}} \phi(\omega) a_{jn}(t,\omega) h_n g(\omega) d\omega \Big| \\ &\leq \delta_\beta \cdot \sup_{\|\phi\|_{\beta,n}=1} \sum^{\infty}_{n=-\infty} \Big| \int_{\mathbf{R}} \phi(\omega) a_{jn}(t,\omega) g(\omega) d\omega \Big|. \end{aligned}$$

In general, for any C^1 periodic function $f(\theta) = \sum_{n=-\infty}^{\infty} f_n e^{\sqrt{-1}n\theta}$, the inequality

$$\sum_{n=-\infty}^{\infty} |f_n| \leq \max|f(\theta)| + 2\max|f'(\theta)| \tag{6.9}$$

holds, where f_n are Fourier coefficients of f, see Newman [34] for the proof. Applying this to the function $\int_{\mathbf{R}} \phi(\omega)e^{\sqrt{-1}jx(t,0;\theta,\omega)}g(\omega)d\omega$, we obtain

$$\begin{aligned}
&\sum_{n=-\infty}^{\infty}\Big|\int_{\mathbf{R}}\phi(\omega)a_{jn}(t,\omega)g(\omega)d\omega\Big| \\
\leq\ &\max_{\theta}\Big|\int_{\mathbf{R}}\phi(\omega)e^{\sqrt{-1}jx(t,0;\theta,\omega)}g(\omega)d\omega\Big| + 2\max_{\theta}\Big|\int_{\mathbf{R}}\phi(\omega)\frac{\partial}{\partial\theta}e^{\sqrt{-1}jx(t,0;\theta,\omega)}g(\omega)d\omega\Big|.
\end{aligned}$$

Cauchy's theorem with the aid of the assumption (A3) provides

$$\begin{aligned}
&\sum_{n=-\infty}^{\infty}\Big|\int_{\mathbf{R}}\phi(\omega)a_{jn}(t,\omega)g(\omega)d\omega\Big| \leq \int_{\mathbf{R}}|\phi(\omega)|g(\omega)d\omega \\
&+2\max_{\theta}\Big|\int_{\mathbf{R}}\phi(\omega+\sqrt{-1}a)je^{\sqrt{-1}(j-1)x(t,0;\theta,\omega+\sqrt{-1}a)}\frac{\partial}{\partial\theta}e^{\sqrt{-1}x(t,0;\theta,\omega+\sqrt{-1}a)}g(\omega+\sqrt{-1}a)d\omega\Big|,
\end{aligned}$$

where a is a positive number. Let us show that the second term above is bounded uniformly in $j = 1, 2\cdots$ and t. Put $X = X(\omega) = e^{\sqrt{-1}x(t,0;\theta,\omega)}$. From the proof of Thm.5.10 (iv), we can verify that there exist $\varepsilon > 0$ and $t_0 > 0$ such that when $t > t_0$, $|X(\omega+\sqrt{-1}a)| < 1-\varepsilon$ for any $t > t_0$. Therefore, the factor jX^{j-1} included in the second term tends to zero as $j \to \infty$. To estimate the derivative $\partial X(\omega+\sqrt{-1}a)/\partial\theta$, note that it satisfies the equation

$$\frac{d}{dt}\left(\frac{\partial X}{\partial\theta}(\omega+\sqrt{-1}a)\right) = \left(\sqrt{-1}\omega - a - K\overline{\eta(t)}X(\omega+\sqrt{-1}a)\right)\frac{\partial X}{\partial\theta}(\omega+\sqrt{-1}a),$$

which is solved as

$$\frac{\partial X}{\partial\theta}(\omega+\sqrt{-1}a) = X|_{t=t_0}\cdot\exp\left[\int_{t_0}^{t}\left(\sqrt{-1}\omega - a - K\overline{\eta(s)}X(\omega+\sqrt{-1}a)\right)ds\right].$$

Put $X = \xi e^{\sqrt{-1}p}$, $\eta = \zeta e^{\sqrt{-1}q}$ with $\xi, \zeta, p, q \in \mathbf{R}$. Then, ξ satisfies the equation (5.45). By using it, we obtain

$$\begin{aligned}
&\mathrm{Re}\left[\int_{t_0}^{t}\left(\sqrt{-1}\omega - a - K\overline{\eta(s)}X(\omega+\sqrt{-1}a)\right)ds\right] \\
=\ &\int_{t_0}^{t}(-a - K\zeta\xi\cos(p-q))\,ds \\
=\ &\int_{t_0}^{t}\left(-a - \frac{2\xi\dot{\xi} + 2a\xi^2}{1-\xi^2}\right)ds \\
=\ &-a\int_{t_0}^{t}\frac{1+\xi^2}{1-\xi^2}ds + \log\left(\frac{1-\xi(t)^2}{1-\xi(t_0)^2}\right).
\end{aligned}$$

Since $0 < \xi(t) < 1 - \varepsilon$ for $t > t_0$, the second term above is bounded in t and the first term tends to $-\infty$ as $t \to \infty$. This proves that $\partial X/\partial\theta \to 0$ as $t \to \infty$. Therefore, there exists N_β such that Eq.(6.8) holds for $t > t_0$. Since the norm $\| \cdot \|^*_{\beta,n}$ is bounded as $n \to \infty$, we can take N_β not to depend on n. Then, the first equation of (6.7) gives

$$\| \, | \overline{Z_1(t,\cdot)} \rangle \|^*_{\beta,n} \leq \delta_\beta M_\beta N_\beta e^{-a(t-t_0)} + \frac{\delta_\beta K M_\beta N_\beta}{2} \int_{t_0}^{t} e^{-a(t-s)} \| \, | \overline{Z_1(s,\cdot)} \rangle \|^*_{\beta,n} ds, \qquad (6.10)$$

for $t > t_0$. Now the Gronwall inequality proves

$$\| \, | \overline{Z_1(t,\cdot)} \rangle \|^*_{\beta,n} \leq \delta_\beta M_\beta N_\beta e^{(\delta_\beta K M_\beta N_\beta/2 - a)(t-t_0)}. \qquad (6.11)$$

Since M_β and N_β are independent of the choice of δ_β, by taking δ_β sufficiently small, this quantity proves to tend to zero as $t \to \infty$. Substituting it into the second equation of (6.7), we obtain

$$\| \, | \overline{Z_j(t,\cdot)} \rangle \|^*_{\beta,n} \leq \delta_\beta M_\beta N_\beta e^{-ja(t-t_0)} + jK(\delta_\beta M_\beta N_\beta)^2 \int_{t_0}^{t} e^{-ja(t-s)} e^{(\delta_\beta K M_\beta N_\beta/2 - a)(s-t_0)} ds \qquad (6.12)$$

for $j = 2, 3, \cdots$. It is easy to verify that the right hand side tends to zero as $t \to \infty$ uniformly in j.

Now we have proved that if the initial condition satisfies (6.5) for each β, then $\| \, | \overline{Z_j(t,\cdot)} \rangle \|^*_{\beta,n}$ decays to zero as $t \to \infty$ for every j and n. By the definition of the norm $\| \cdot \|^*_{\beta,n}$, this means that $(Z_j(t,\cdot), \overline{\phi}) \to 0$ as $t \to \infty$ for every $\phi \in F_p(\beta)$. ■

7 Bifurcation theory

Now we are in a position to investigate bifurcation of the Kuramoto model by using the center manifold reduction. Our strategy to detect bifurcation is that we use the space of functionals F'_p instead of the spaces of usual functions F_p or $L^2(\mathbf{R}, g(\omega)d\omega)$ because the linear operator T_1 admits the spectral decomposition on F'_p consisting of eigenfunctions, while the spectral decomposition on $L^2(\mathbf{R}, g(\omega)d\omega)$ involves the continuous spectrum on the imaginary axis; that is, a center manifold on $L^2(\mathbf{R}, g(\omega)d\omega)$ is an infinite dimensional manifold. To avoid such a difficulty, we will seek a center manifold on F'_p. At first, we have to prove the existence of center manifolds. Standard results of the existence of center manifolds (see [5, 9, 24, 47]) are not applicable to our system because the space F'_p is not a Banach space and the projection Π_c to the center subspace is continuous only on a subspace of F'_p. Thus in Sec.7.1, the existence theorem of center manifolds for our system and a strategy for proving it are given. The proof of the theorem is given in Sec.7.2 to 7.4. In Sec.7.5, the dynamics on the center manifold is derived and the Kuramoto's conjecture is solved. Readers who are interested in a practical method for obtaining a bifurcation structure can skip Sec.7.1 to 7.4 and go to Sec.7.5. Throughout this section, we suppose the assumptions (A1) to (A3) and (A5) for $g(\omega)$. Existence of center manifolds for the case that $g(\omega)$ is a rational function is trivial because the phase space $i(H_+)$ is a finite dimensional vector space.

7.1 Center manifold theorem

Let $i(\mathcal{F})$ be a certain metric subspace of the product space $\prod_{k=1}^{\infty} F'_p$ with a distance d_∞, and $\overline{i(\mathcal{F})}$ its closure. These spaces and the metric d_∞ will be introduced in Sec.7.2 and Sec.7.3. Let Φ_t be the semiflow on $\overline{i(\mathcal{F})}$ generated by the system (6.2) (for the existence of the semiflow, see Sec.7.3). For the generalized center subspace $\mathbf{E}_c$ of T_1 defined by (5.49), put

$$\hat{\mathbf{E}}_c = \mathbf{E}_c \times \{0\} \times \{0\} \times \cdots \subset \prod_{k=1}^{\infty} F'_p. \tag{7.1}$$

Let $\hat{\mathbf{E}}_c^\perp = \mathbf{E}_c^\perp \times F'_p \times F'_p \times \cdots$ be the complement of $\hat{\mathbf{E}}_c$. The existence theorem of center manifolds is stated as follows.

Theorem 7.1. Suppose that $g(\omega)$ satisfies the assumptions (A1) to (A3) and (A5). There exist a positive number ε_0 and an open set U of the origin in $\overline{i(\mathcal{F})} \subset \prod_{k=1}^{\infty} F'_p$ such that when $|K - K_c| < \varepsilon_0$, the following holds:

(I) (center manifold). There exists a C^1 mapping $\hat{q} : \hat{\mathbf{E}}_c \to \hat{\mathbf{E}}_c^\perp \cap \overline{i(\mathcal{F})}$ such that the dim-$\mathbf{E}_c$ dimensional C^1 manifold defined to be

$$W^c_{loc} = \{y + \hat{q}(y) \,|\, y \in \hat{\mathbf{E}}_c\} \cap U \tag{7.2}$$

is Φ_t-invariant (that is, $\Phi_t(W^c_{loc}) \cap U \subset W^c_{loc}$). This is called the *local center manifold*. The mapping $\hat{q}$ is also C^1 with respect to the parameter $\varepsilon := K - K_c$, and $\hat{q}(y) \sim O(y^2, \varepsilon y, \varepsilon^2)$ as $y, \varepsilon \to 0$

(II) (negative semi-orbit). For every $\xi_0 \in W^c_{loc}$, there exists a function $u : (-\infty, 0] \to \overline{i(\mathcal{F})}$ such that $u(0) = \xi_0$ and $\Phi_t(u(s)) = u(t + s)$ when $t \geq 0$, $s \leq -t$. Such a $u(t)$ is called a *negative semi-orbit* of (6.2). As long as $u(t) \in U$, $u(t) \in W^c_{loc}$. In this case, there exist $C_1 > 0$ and a small number $b > 0$ such that

$$d_\infty(u(t), 0) \leq C_1 e^{bt}. \tag{7.3}$$

(III) (invariant foliation). There exists a family of manifolds $\{M_\xi\}_{\xi \in W^c_{loc}} \subset U$, parameterized by $\xi \in W^c_{loc}$, satisfying that
(i) $M_\xi \cap W^c_{loc} = \{\xi\}$, $\bigcup_{\xi \in W^c_{loc}} M_\xi = U$, and $M_\xi \cap M_{\xi'} = \emptyset$ if $\xi \neq \xi'$.
(ii) when $\Phi_t(\xi) \in U$, $\Phi_t(M_\xi) \cap U \subset M_{\Phi_t(\xi)}$.
(iii)

$$\begin{aligned} M_\xi \;=\; \{u \in \overline{i(\mathcal{F})} \cap U \,\big|\, & \\ & \text{there exist } a > b \text{ and } C_2 > 0 \text{ such that } d_\infty(\Phi_t(u), \Phi_t(\xi)) \leq C_2 e^{-at}\}. \end{aligned}$$

By Prop.5.11, W^c_{loc} is a finite dimensional invariant manifold. Part (III) of the theorem means that W^c_{loc} is attracting with the decay rate e^{-at}, where the constant a is the same as that in Prop.5.19. Further, (III)-(iii) means that the semiflow near W^c_{loc} is eventually well approximated by the semiflow on W^c_{loc} if $t > 0$ is large. In particular, if (6.2) has an attractor N near the origin, N is included in W^c_{loc}. Since the topology induced by the metric coincides with the strong and the weak dual topologies on any bounded set, N is attracting for both of the strong and the weak dual topologies. Due to the spectral decomposition (5.65), any element $|\,\overline{Z}_1\rangle \in i(F_p)$ is decomposed as

$$|\,\overline{Z}_1\rangle = \sum_{j=0}^{n} \alpha_j |\,\mu_j\rangle + Y_1, \quad \alpha_j \in \mathbf{C},\ Y_1 \in \mathbf{E}_c^{\perp}, \tag{7.4}$$

where $|\,\mu_0\rangle, \cdots, |\,\mu_n\rangle \in \mathbf{E}_c$ are generalized eigenfunctions associated with the resonance poles on the imaginary axis. Then, part (I) of the theorem means that if $(|\,\overline{Z}_1\rangle, |\,\overline{Z}_2\rangle, \cdots) \in W^c_{loc}$,

$$Y_1 \sim O(\alpha^2, \varepsilon\alpha, \varepsilon^2), \quad |\,\overline{Z}_k\rangle \sim O(\alpha^2, \varepsilon\alpha, \varepsilon^2),\ k = 2, 3, \cdots, \tag{7.5}$$

as $\varepsilon, \alpha_0, \cdots, \alpha_n \to 0$, where $\alpha = (\alpha_0, \cdots, \alpha_n)$. Substituting Eq.(7.4) into the system (6.2) with the condition (7.5), we can obtain the expression of $\hat{q}(y)$ as a function of $\varepsilon, \alpha_0, \cdots, \alpha_n$. The dynamics on W^c_{loc} is realized by a system of ordinary differential equations of $\alpha_0, \cdots, \alpha_n$:

$$\frac{d\alpha_j}{dt} = f_j(\varepsilon; \alpha_0, \cdots, \alpha_n), \quad j = 0, \cdots, n. \tag{7.6}$$

If (6.2) has an attractor N near the origin, N is an attractor of the system (7.6). In this manner, (6.2) is reduced to a finite dimensional problem. Such a technique to investigate bifurcation is called the *center manifold reduction*. Part (II) implies that the center manifold is characterized by the property that the dynamics on it is sufficiently slow.

Although we prove the existence of W^c_{loc} in $\overline{i(\mathcal{F})}$, from a physical viewpoint, especially we are interested in an initial condition of the form $|\overline{Z_j(0)}\rangle = \overline{h}_j|P_0\rangle$ (which corresponds to the initial condition of the form $\rho_0(\theta, \omega) = h(\theta)$ for the system (2.1)). Then $|\overline{Z_j(t)}\rangle \in i(V_{1,0})$ (Thm.5.10 (iv)). This means that an attractor of (6.2) which is reached from the initial condition $|\overline{Z_j(0)}\rangle = \overline{h}_j|P_0\rangle$ is included in $W^c_{loc} \cap \prod_{k=1}^{\infty} \overline{i(V_{1,0})}$.

Sec.7.2 to 7.4 are devoted to prove Thm.7.1. It is well known that a global center manifold uniquely exists only when a Lipschitz constant of nonlinear terms of a system is sufficiently small. Thus in Sec.7.2, we consider a perturbed system of (6.2) so that its Lipschitz constant becomes sufficiently small, while it coincides with the original system in the vicinity of the origin. Because of the perturbation, a solution may fall out of $i(V_{1,0})$ and go into a larger space. Thus we will introduce the space $\mathcal{F}$, and show that solutions $(Z_1, Z_2, \cdots)$ of the perturbed system are included in $\mathcal{F}$. We will prove in Sec.7.3 and Appendix B that the perturbed system generates a smooth flow to prove that the center manifold is smooth. Once we obtain the existence of a proper phase space, a spectral decomposition of the linear operator, estimates of norms of the semigroups and a smooth flow whose Lipschitz constant of nonlinear terms is sufficiently small, then the existence of the center manifold is proved in usual way with a slight modification. We demonstrate it in Sec.7.4. In Sec.7.5, we perform the center manifold reduction: equations of $\alpha_0, \cdots, \alpha_n$ are obtained and investigated. The order parameter $\eta(t)$ is defined as $\eta(t) = (Z_1, P_0) = \langle P_0\,|\,\overline{Z}_1\rangle$. On the center manifold, it is written as

$$\langle P_0\,|\,\overline{Z}_1\rangle = \sum_{j=0}^{n} \overline{\alpha_j(t)}\langle P_0\,|\,\mu_j\rangle + \langle P_0\,|\,Y_1\rangle = \sum_{j=0}^{n} \frac{2}{K_c}\overline{\alpha_j(t)} + O(\alpha^2, \varepsilon\alpha, \varepsilon^2), \tag{7.7}$$

where we use $\langle P_0 \,|\, \mu_j \rangle = 2/K_c$, which follows from the definition of resonance poles. Therefore, if a bifurcation diagram of (7.6) is obtained, a bifurcation diagram of the order parameter is also obtained. In this way, the Kuramoto's conjecture will be proved in Sec.7.5.

7.2 Phase space of the perturbed system

Recall that the trivial solution $Z_j(t) \equiv 0\,(j = 1, 2, \cdots)$ corresponds to $\rho_t \equiv 1/2\pi$ called the de-synchronous state (Sec.3). Since we are interested in bifurcations from $\rho_t \equiv 1/2\pi$ at $K = K_c$, put $\rho_t = 1/2\pi + \hat{\rho}_t$ and $K = K_c + \varepsilon$. Then, Eq.(2.1) is rewritten as

$$\begin{aligned}
&\frac{\partial \hat{\rho}_t}{\partial t} + \frac{\partial}{\partial \theta}\Big(\omega \hat{\rho}_t + \frac{K_c}{4\pi\sqrt{-1}}(\eta(t)e^{-\sqrt{-1}\theta} - \overline{\eta(t)}e^{\sqrt{-1}\theta}) \\
&+ \frac{\varepsilon}{4\pi\sqrt{-1}}(\eta(t)e^{-\sqrt{-1}\theta} - \overline{\eta(t)}e^{\sqrt{-1}\theta}) + \frac{K}{2\sqrt{-1}}(\eta(t)e^{-\sqrt{-1}\theta} - \overline{\eta(t)}e^{\sqrt{-1}\theta})\hat{\rho}_t\Big) = 0, \quad (7.8)
\end{aligned}$$

where

$$\eta(t) = \int_{\mathbf{R}}\int_0^{2\pi} e^{\sqrt{-1}\theta}\rho_t(\theta, \omega)d\theta d\omega = \int_{\mathbf{R}}\int_0^{2\pi} e^{\sqrt{-1}\theta}\hat{\rho}_t(\theta, \omega)d\theta d\omega.$$

An initial condition $\hat{\rho}_0 = \hat{h}(\theta, \omega)$ is a real-valued measure (signed measure) on S^1 parameterized by $\omega \in \mathbf{R}$ satisfying $\int_0^{2\pi}\hat{h}(\theta, \omega)d\theta d\omega = 0$. The first step to prove the existence of center manifolds is to localize the nonlinear term so that the Lipschitz constant of the nonlinear term becomes sufficiently small. For this purpose, let $\hat{\chi} : [0, \infty) \to [0, 1]$ be a C^∞ function, and consider the perturbed continuous model of the form

$$\begin{aligned}
&\frac{\partial \hat{\rho}_t}{\partial t} + \frac{\partial}{\partial \theta}\Big(\omega \hat{\rho}_t + \frac{K_c}{4\pi\sqrt{-1}}(\eta(t)e^{-\sqrt{-1}\theta} - \overline{\eta(t)}e^{\sqrt{-1}\theta}) \\
&+ \frac{\varepsilon}{4\pi\sqrt{-1}}(\eta(t)e^{-\sqrt{-1}\theta} - \overline{\eta(t)}e^{\sqrt{-1}\theta})\hat{\chi}(t) + \frac{K}{2\sqrt{-1}}(\eta(t)e^{-\sqrt{-1}\theta} - \overline{\eta(t)}e^{\sqrt{-1}\theta})\hat{\chi}(t)\hat{\rho}_t\Big) = 0. \quad (7.9)
\end{aligned}$$

If we put

$$\hat{Z}_j(t, \omega) = \int_0^{2\pi} e^{\sqrt{-1}j\theta}\hat{\rho}_t(\theta, \omega)d\theta, \quad (7.10)$$

Eq.(7.9) yields a system of equations

$$\left\{\begin{aligned}
\frac{d\hat{Z}_1}{dt} &= \sqrt{-1}\omega\hat{Z}_1 + \frac{K_c}{2}\eta(t) + \Big(\frac{\varepsilon}{2}\eta(t) - \frac{K}{2}\overline{\eta(t)}\hat{Z}_2\Big)\hat{\chi}(t), \\
\frac{d\hat{Z}_j}{dt} &= j\sqrt{-1}\omega\hat{Z}_j + \frac{jK}{2}\left(\eta(t)\hat{Z}_{j-1} - \overline{\eta(t)}\hat{Z}_{j+1}\right)\hat{\chi}(t), \quad j = 2, 3, \cdots .
\end{aligned}\right. \quad (7.11)$$

If $\hat{\chi} \equiv 1$, this coincides with the original system (3.2),(3.3). If $\hat{\chi}(t)$ is sufficiently small, this perturbation makes the Lipschitz constant of the nonlinear terms of (3.2),(3.3) sufficiently small (Note that when proving the existence of center manifolds, the bifurcation parameter

ε is regarded as a dependent variable. Thus $\varepsilon\eta(t) = \varepsilon(\hat{Z}_1, P_0)$ is regarded as a nonlinear term). A concrete definition of $\hat{\chi}$ will be specified in Sec.7.3. Note that $\hat{Z}_0 \equiv 0$ because of $\int_0^{2\pi} \hat{h}(\theta, \omega) d\theta d\omega = 0$.

Eq.(7.9) is integrated by using the characteristic curve method. The characteristic curve $x = x(t, s; \theta, \omega)$ is defined as a solution of the equation

$$\frac{dx}{dt} = \omega + \frac{K}{2\sqrt{-1}} \left(\eta(t) e^{-\sqrt{-1}x} - \overline{\eta(t)} e^{\sqrt{-1}x} \right) \hat{\chi}(t), \tag{7.12}$$

satisfying the initial condition $x(s, s; \theta, \omega) = \theta$ at an initial time s. Along the characteristic curve, (7.9) is integrated to yield

$$\begin{aligned} \hat{\rho}_t(\theta, \omega) \;&=\; \hat{h}(x(0, t; \theta, \omega), \omega) \exp\Bigl[\frac{K}{2} \int_0^t \Bigl(\eta(s) e^{-\sqrt{-1}x(s,t;\theta,\omega)} + \overline{\eta(s)} e^{\sqrt{-1}x(s,t;\theta,\omega)} \Bigr) \hat{\chi}(s) ds \Bigr] \\ &\quad + \int_0^t \exp\Bigl[\frac{K}{2} \int_s^t \Bigl(\eta(\tau) e^{-\sqrt{-1}x(\tau,t;\theta,\omega)} + \overline{\eta(\tau)} e^{\sqrt{-1}x(\tau,t;\theta,\omega)} \Bigr) \hat{\chi}(\tau) d\tau \Bigr] \times \\ &\qquad \left(\frac{K_c + \varepsilon\hat{\chi}(s)}{4\pi} \right) \Bigl(\eta(s) e^{-\sqrt{-1}x(s,t;\theta,\omega)} + \overline{\eta(s)} e^{\sqrt{-1}x(s,t;\theta,\omega)} \Bigr) ds. \end{aligned} \tag{7.13}$$

Once $x(t, s; \theta, \omega)$ and $\eta(t)$ are determined, this $\hat{\rho}_t$ gives a weak solution of (7.9). The existence of solutions of (7.11) follows from that of the integro-ODE (7.12) and (7.13), which will be proved in Appendix B (see also Prop.7.3).

In Sec.5, we have proved that a solution of (3.2),(3.3) is included in $V_{1,0}$. This property may break down because of the perturbation $\hat{\chi}$. Thus we define an appropriate phase space for (7.11) and prove the existence of the flow on it. Define a subspace $\mathcal{F}$ of the product space $\prod_{j=1}^{\infty} F_p$ as follows: $(Z_1, Z_2, \cdots) \in \mathcal{F}$ if and only if there exists a signed measure $\hat{h}(\theta, \omega)$ on S^1 parameterized by $\omega \in \mathbf{R}$ such that

(F1)

$$\int_0^{2\pi} \hat{h}(\theta, \omega) d\theta = 0.$$

(F2) For $j = 1, 2, \cdots$,

$$Z_j(\omega) = \int_0^{2\pi} e^{\sqrt{-1}j\theta} \hat{h}(\theta, \omega) d\theta \in V \;(= \bigcup_{C \geq 1} V_{2p}(C, 0)). \tag{7.14}$$

(F3) Define Z_j for $j = -1, -2, \cdots$ by Eq.(7.14). There exist positive constants C and γ such that

$$\sup_{0 \leq \mathrm{Im}(\omega) \leq p} |Z_j(\omega)| \leq C e^{|j|\gamma} \tag{7.15}$$

for all $j \in \mathbf{Z}$.

Recall that the set $V \subset F_p$ was defined in Prop.5.20. Hence, $\mathcal{F}$ is the set of Fourier coefficients $\{Z_j\}_{j=1}^{\infty}$ of signed measures included in $\prod_{j=1}^{\infty} V$ whose growth rate in j is not so fast. Although $Z_j(\omega)$ is assumed to be holomorphic near the region $0 \leq \mathrm{Im}(\omega) \leq p$, $\hat{h}(\theta, \omega)$

need not have an analytic continuation to the complex ω-plane.

Example. If $\hat{h}$ is of the form $\hat{h}(\theta,\omega) = h_1(\theta)h_2(\omega)$ with $h_2(\omega)\in V$, the assumptions (F2) and (F3) are satisfied. In particular, $|Z_j(\omega)|$ is bounded uniformly in j. Recall that from a physical viewpoint, we are interested in an initial condition of the form $\hat{h}(\theta,\omega) = h(\theta)$ (see Eq.(2.1)). Let $\delta(\theta+\omega)$ be the delta function at $\theta = -\omega$. If $\hat{h}(\theta,\omega) = \delta(\theta+\omega) - 1/(2\pi)$, these assumptions are satisfied with $C = 1, \gamma = p$ because $Z_j(\omega) = e^{-\sqrt{-1}j\omega}$ and $|Z_j(\omega)| = e^{j\cdot\mathrm{Im}(\omega)}$. If $\hat{h}(\theta,\omega) = \theta\cos(\theta^2\omega)$, we can verify that $|Z_j(\omega)| \sim O(e^{j^2})$ for $\omega\in\mathbf{C}$ as $j\to\pm\infty$, and thus (F3) is violated.

The existence of solutions of Eq.(7.11) will be prove in Sec.7.3 and Appendix B after $\hat{\chi}$ is specified. In this section, we show that if solutions exist, they are included in $\mathcal{F}$.

Proposition 7.2. For a given function $\hat{\chi} : [0,\infty) \to [0,1]$ and an initial condition in $\mathcal{F}$, suppose that a solution of (7.11) exists and a function $x(t,s;\theta,\omega)$ has an analytic continuation with respect to θ and ω (these facts will be verified in Sec.7.3 and Appendix B). Then,
(i) the solution is included in $\mathcal{F}$ for any $t\geq 0$.
(ii) for each $t\geq 0$, $\beta = 0,1,\cdots$ and $n = 1,2,\cdots$, $\|\,|\overline{Z_j(t,\cdot)}\rangle\|^*_{\beta,n}$ is bounded uniformly in $j\in\mathbf{Z}$, where $|\overline{Z_j(t,\cdot)}\rangle = i(\hat{Z}_j(t,\cdot))$.

Proof. (i) At first, let us verify (F2). To show that $\hat{Z}_j(t,\,\cdot\,)\in V$ if $\hat{Z}_j(0,\,\cdot\,)\in V$, we use the equality

$$\begin{aligned}\int_0^{2\pi} a(\theta,\omega)\hat{\rho}_t(\theta,\omega)d\theta &= \int_0^{2\pi} a(x(t,0;\theta,\omega),\omega)\hat{h}(\theta,\omega)d\theta \\ &+ \int_0^{2\pi}\!\!\int_0^t a(x(t,0;\theta,\omega),\omega)\exp\Big[-\frac{K}{2}\int_0^s\Big(\eta(\tau)e^{-\sqrt{-1}x(\tau,0;\theta,\omega)} + \overline{\eta(\tau)}e^{\sqrt{-1}x(\tau,0;\theta,\omega)}\Big)\hat{\chi}(\tau)d\tau\Big]\times \\ &\qquad\left(\frac{K_c+\varepsilon\hat{\chi}(s)}{4\pi}\right)\Big(\eta(s)e^{-\sqrt{-1}x(s,0;\theta,\omega)} + \overline{\eta(s)}e^{\sqrt{-1}x(s,0;\theta,\omega)}\Big)dsd\theta, \qquad (7.16)\end{aligned}$$

for any measurable function $a(\theta,\omega)$, which is proved by substitution of (7.13). Note that if $\hat{\chi}(t)\equiv 1$, it is reduced to Eq.(2.5). From this, it turns out that $\hat{Z}_j$ is expressed as

$$\begin{aligned}\hat{Z}_j(t,\omega) &= \int_0^{2\pi} e^{\sqrt{-1}jx(t,0;\theta,\omega)}\hat{h}(\theta,\omega)d\theta \\ &+ \int_0^{2\pi}\!\!\int_0^t e^{\sqrt{-1}jx(t,0;\theta,\omega)}\exp\Big[-\frac{K}{2}\int_0^s\Big(\eta(\tau)e^{-\sqrt{-1}x(\tau,0;\theta,\omega)} + \overline{\eta(\tau)}e^{\sqrt{-1}x(\tau,0;\theta,\omega)}\Big)\hat{\chi}(\tau)d\tau\Big]\times \\ &\qquad\left(\frac{K_c+\varepsilon\hat{\chi}(s)}{4\pi}\right)\Big(\eta(s)e^{-\sqrt{-1}x(s,0;\theta,\omega)} + \overline{\eta(s)}e^{\sqrt{-1}x(s,0;\theta,\omega)}\Big)dsd\theta. \qquad (7.17)\end{aligned}$$

At first, let us show that $e^{\pm\sqrt{-1}x(t,0;\theta,\omega)}\in V$. This is proved in the same way as Thm.5.10 (iv). Put $X(t) = e^{\sqrt{-1}x(t,0;\theta,\omega)}$. Then X satisfies the equation

$$\left\{\begin{array}{l}\dfrac{dX}{dt} = \sqrt{-1}\omega X + \dfrac{K}{2}\Big(\eta(t) - \overline{\eta(t)}X^2\Big)\hat{\chi}(t), \\ X(0) = e^{\sqrt{-1}\theta}.\end{array}\right. \qquad (7.18)$$

Putting $X = \xi e^{\sqrt{-1}p}$, $\eta = \zeta e^{\sqrt{-1}q}$ with $\xi, \zeta, p, q \in \mathbf{R}$ yields

$$\frac{d\xi}{dt} = -\mathrm{Im}(\omega)\xi + \frac{K}{2}\zeta(1-\xi^2)\cos(p-q)\hat{\chi}(t). \tag{7.19}$$

This equation implies that if $\mathrm{Im}(\omega) \geq 0$ and $\xi = 1$, then $d\xi/dt \leq 0$. Since $|X(0)| = 1$, we obtain $|X(t)| \leq 1$ for any $t \geq 0$ and any $\mathrm{Im}(\omega) \geq 0$. This proves that $X(t)$ is bounded on the real axis and the upper half plane: $X(t) = e^{\sqrt{-1}x(t,0;\theta,\omega)} \in V_{2p}(1,0) \subset V$.

Next thing to do is to investigate $Y(t) = e^{-\sqrt{-1}x(t,0;\theta,\omega)}$, which satisfies

$$\begin{cases} \dfrac{dY}{dt} = -\sqrt{-1}\omega Y - \dfrac{K}{2}\Big(\eta(t)Y^2 - \overline{\eta(t)}\Big)\hat{\chi}(t), \\ Y(0) = e^{-\sqrt{-1}\theta}. \end{cases} \tag{7.20}$$

Putting $Y = \xi e^{\sqrt{-1}p}$, $\eta = \zeta e^{\sqrt{-1}q}$ with $\xi, \zeta, p, q \in \mathbf{R}$ yields

$$\begin{cases} \dfrac{d\xi}{dt} = \mathrm{Im}(\omega)\xi + \dfrac{K}{2}\zeta(1-\xi^2)\cos(p+q)\hat{\chi}(t), \\ \dfrac{dp}{dt} = -\mathrm{Re}(\omega) - \dfrac{K}{2}\zeta(\xi + \dfrac{1}{\xi})\sin(p+q)\hat{\chi}(t). \end{cases} \tag{7.21}$$

When $|\mathrm{Re}(\omega)|$ is sufficiently large, the averaging method is applicable to construct an approximate solution. Eq.(7.21) is averaged with respect to p to provide the averaging equation $d\xi/dt = \mathrm{Im}(\omega)\xi$, which is solved as $\xi(t) = e^{\mathrm{Im}(\omega)t}\xi(0)$. Therefore, a solution of Eq.(7.21) is given as

$$\xi(t) = e^{\mathrm{Im}(\omega)t} + O\Big(\frac{1}{|\mathrm{Re}(\omega)|}\Big), \tag{7.22}$$

as $|\mathrm{Re}(\omega)| \to \infty$. See Sanders and Verhulst [39] for the averaging method. This implies that Y is bounded as $\mathrm{Re}(\omega) \to \pm\infty$ for each $\mathrm{Im}(\omega)$ and t. Thus $e^{-\sqrt{-1}x(t,0;\theta,\omega)} \in V$. Therefore, the second term in the right hand side of Eq.(7.17) is in V; the second term is bounded as $\mathrm{Re}(\omega) \to \pm\infty$ for each j and t.

Next, we show that the first term in the right hand side of Eq.(7.17) is in V. Let

$$e^{\sqrt{-1}jx(t,0;\theta,\omega)} = \sum_{n=-\infty}^{\infty} a_{jn}(t,\omega)e^{\sqrt{-1}n\theta} \tag{7.23}$$

be a Fourier expansion of $e^{\sqrt{-1}jx(t,0;\theta,\omega)}$. Then,

$$\begin{aligned} \int_0^{2\pi} e^{\sqrt{-1}jx(t,0;\theta,\omega)}\hat{h}(\theta,\omega)d\theta &= \sum_{n=-\infty}^{\infty} a_{jn}(t,\omega)\int_0^{2\pi} e^{\sqrt{-1}n\theta}\hat{h}(\theta,\omega)d\theta \\ &= \sum_{n=-\infty}^{\infty} a_{jn}(t,\omega)\hat{Z}_n(0,\omega). \end{aligned} \tag{7.24}$$

Since the series (7.23) converges uniformly in θ, the right hand side of (7.24) exists for each ω. Since $e^{\sqrt{-1}jx(t,0;\theta,\omega)}$ is holomorphic, so is $a_{jn}(t,\omega)$. By the assumption (F2), $\hat{Z}_n(0,\omega)$

is also holomorphic. Therefore, the right hand side of (7.24) converges to a holomorphic function near the region $0 \leq \mathrm{Im}(\omega) \leq p$. By the assumption (F3), there are positive constants C and γ such that

$$\sup_{0\leq \mathrm{Im}(\omega)\leq p} |\hat{Z}_n(0,\omega)| \leq Ce^{|n|\gamma}. \tag{7.25}$$

This provides the inequality

$$\sup_{0\leq \mathrm{Im}(\omega)\leq p} \Big| \sum_{n=-\infty}^{\infty} a_{jn}(t,\omega)\hat{Z}_n(0,\omega) \Big| \leq \sup_{0\leq \mathrm{Im}(\omega)\leq p} C \sum_{n=-\infty}^{\infty} e^{|n|\gamma} |a_{jn}(t,\omega)|. \tag{7.26}$$

Let us prove that the right hand side exists. Eq.(7.12) shows that $x(t,0;\theta+2\pi,\omega) = x(t,0;\theta,\omega)+2\pi$. With this property, we use Cauchy's theorem to the function $e^{\sqrt{-1}jx(t,0;\theta,\omega)}$ along the path represented in Fig.11(a) to yield

$$\begin{aligned} a_{jn}(t,\omega) &= \frac{1}{2\pi}\int_{C_1} e^{-\sqrt{-1}n\theta} e^{\sqrt{-1}jx(t,0;\theta,\omega)} d\theta \\ &= -\frac{1}{2\pi}\int_{C_2} e^{-\sqrt{-1}n\theta} e^{\sqrt{-1}jx(t,0;\theta,\omega)} d\theta \\ &= \frac{e^{-nr}}{2\pi}\int_0^{2\pi} e^{-\sqrt{-1}n\theta} e^{\sqrt{-1}jx(t,0;\theta-\sqrt{-1}r,\omega)} d\theta, \end{aligned} \tag{7.27}$$

for $n = 0,1,2,\cdots$ and $j = \pm 1, \pm 2, \cdots$, where $r > 0$ can be taken arbitrarily large because $e^{\sqrt{-1}jx(t,0;\theta,\omega)}$ is analytic in $\theta \in \mathbf{C}$. By the same way as above, we can show that $e^{\sqrt{-1}jx(t,0;\theta-\sqrt{-1}r,\omega)}$ is estimated as

$$|e^{\sqrt{-1}jx(t,0;\theta-\sqrt{-1}r,\omega)}| = e^{(-\mathrm{Im}(\omega)t+r)j} + O\Big(\frac{1}{|\mathrm{Re}(\omega)|}\Big),$$

as $|\mathrm{Re}(\omega)| \to \infty$ for each t and $\mathrm{Im}(\omega)$. This provides

$$|a_{jn}(t,\omega)| \leq e^{-nr}\left(e^{(-\mathrm{Im}(\omega)t+r)j} + O\Big(\frac{1}{|\mathrm{Re}(\omega)|}\Big)\right),$$

for $n = 0,1,2,\cdots$. When $n < 0$, we take a path represented as Fig.11(b), which yields

$$|a_{jn}(t,\omega)| \leq e^{nr}\left(e^{(-\mathrm{Im}(\omega)t-r)j} + O\Big(\frac{1}{|\mathrm{Re}(\omega)|}\Big)\right)$$

as $|\mathrm{Re}(\omega)| \to \infty$ in the same way. Therefore, we obtain

$$\begin{aligned} &\sup_{0\leq \mathrm{Im}(\omega)\leq p} \Big| \sum_{n=-\infty}^{\infty} a_{jn}(t,\omega)\hat{Z}_n(0,\omega) \Big| \\ \leq\ & \sup_{0\leq \mathrm{Im}(\omega)\leq p} C \sum_{n=1}^{\infty} e^{|n|(\gamma-r)} \left(e^{(-\mathrm{Im}(\omega)t+r)j} + e^{(-\mathrm{Im}(\omega)t-r)j} + O\Big(\frac{1}{|\mathrm{Re}(\omega)|}\Big)\right). \end{aligned}$$

By taking $r > \gamma$, it turns out that the right hand side of Eq.(7.26) exists and bounded as $\mathrm{Re}(\omega) \to \pm\infty$ for each j and t. This proves that $\hat{Z}_j(t, \omega)$ is bounded as $\mathrm{Re}(\omega) \to \pm\infty$ and $\hat{Z}_j(t, \omega) \in V$ for each j and t.

To verify (F3), put

$$\begin{aligned} A(s, \theta) &= \exp\Bigl[-\frac{K}{2}\int_0^s \Bigl(\eta(\tau)e^{-\sqrt{-1}x(\tau,0;\theta,\omega)} + \overline{\eta(\tau)}e^{\sqrt{-1}x(\tau,0;\theta,\omega)}\Bigr)\hat{\chi}(\tau)d\tau\Bigr] \times \\ &\qquad \left(\frac{K_c + \varepsilon\hat{\chi}(s)}{4\pi}\right)\Bigl(\eta(s)e^{-\sqrt{-1}x(s,0;\theta,\omega)} + \overline{\eta(s)}e^{\sqrt{-1}x(s,0;\theta,\omega)}\Bigr). \end{aligned}$$

Then, $\hat{Z}_j$ is rewritten as

$$\hat{Z}_j(t, \omega) = \sum_{n=-\infty}^{\infty} a_{jn}(t, \omega)\hat{Z}_n(0, \omega) + \int_0^t \int_0^{2\pi} e^{\sqrt{-1}jx(t,0;\theta,\omega)} A(s, \theta) d\theta ds. \tag{7.28}$$

From the above calculation, we obtain

$$\begin{aligned} \sup_{0 \le \mathrm{Im}(\omega) \le p} |\hat{Z}_j(t, \omega)| &\le \sup_{0 \le \mathrm{Im}(\omega) \le p}\ \sup_{0 \le \theta \le 2\pi} C \sum_{n=0}^{\infty} e^{|n|(\gamma - r)} \Bigl(|e^{\sqrt{-1}x(t,0;\theta-\sqrt{-1}r,\omega)}|^j + |e^{\sqrt{-1}x(t,0;\theta+\sqrt{-1}r,\omega)}|^j\Bigr) \\ &\quad + \sup_{0 \le \mathrm{Im}(\omega) \le p}\ \sup_{0 \le \theta \le 2\pi} 2\pi \int_0^t |A(s, \theta)| ds \cdot |e^{\sqrt{-1}x(t,0;\theta,\omega)}|^j, \end{aligned}$$

which proves that $\hat{Z}_j(t, \omega)$ satisfies (7.15) for some C and γ.

Finally, let us verify (F1). Note that Eq.(7.12) provides

$$\frac{\partial x}{\partial \theta}(s, 0; \theta, \omega) = \exp\Bigl[-\frac{K}{2}\int_0^s \Bigl(\eta(\tau)e^{-\sqrt{-1}x(\tau,0;\theta,\omega)} + \overline{\eta(\tau)}e^{\sqrt{-1}x(\tau,0;\theta,\omega)}\Bigr)\hat{\chi}(\tau)d\tau\Bigr].$$

This shows that the second term in the right hand side of Eq.(7.16) is rewritten as

$$\int_0^t \frac{K_c + \varepsilon\hat{\chi}(s)}{4\pi} \int_0^{2\pi} a(x(t, 0; \theta, \omega), \omega)\sqrt{-1}\frac{\partial}{\partial\theta}\Bigl(\eta(s)e^{-\sqrt{-1}x(s,0;\theta,\omega)} - \overline{\eta(s)}e^{\sqrt{-1}x(s,0;\theta,\omega)}\Bigr) d\theta ds.$$

In particular, when $a(\theta, \omega) \equiv 1$, this value vanishes because $x(s, 0; \theta, \omega)$ is periodic in θ. This fact and Eq.(7.16) yield

$$\int_0^{2\pi} \hat{\rho}_t(\theta, \omega) d\theta = \int_0^{2\pi} \hat{h}(\theta, \omega) d\theta. \tag{7.29}$$

Therefore, if an initial condition satisfies (F1), so is $\hat{\rho}_t(\theta, \omega)$ for any $t \geq 0$. Now the proof of Prop.7.2 (i) is completed.

To prove (ii), note that when $\omega \in \mathbf{R}$, $\hat{Z}_j(t, \omega)$ is bounded uniformly in $j \in \mathbf{Z}$ because they are Fourier coefficients of a measure $\hat{\rho}_t$ on S^1. This shows that $\| \,|\overline{Z_j(t, \cdot)}\rangle \|^*_{\beta,n}$ is bounded uniformly in $j \in \mathbf{Z}$. ■



(a) (b)

Fig. 11: The contour for obtaining Eq.(7.27).

7.3 Localization of the semiflow

By using the canonical inclusion, we rewrite Eq.(7.11) as an evolution equation on $\mathbf{R} \times \prod_{k=1}^{\infty} F_p'$ of the form

$$\begin{cases} \dfrac{d}{dt}\varepsilon = 0, \\ \dfrac{d}{dt}|\,\overline{Z}_1\rangle = T_{10}^{\times}|\,\overline{Z}_1\rangle + \dfrac{1}{2}\Big(\varepsilon\overline{\langle P_0\,|\,\overline{Z}_1\rangle}|P_0\rangle - K\langle P_0\,|\,\overline{Z}_1\rangle|\,\overline{Z}_2\rangle\Big)\hat{\chi}(t), \\ \dfrac{d}{dt}|\,\overline{Z}_j\rangle = T_{j}^{\times}|\,\overline{Z}_j\rangle + \dfrac{jK}{2}\Big(\overline{\langle P_0\,|\,\overline{Z}_1\rangle}|\,\overline{Z}_{j-1}\rangle - \langle P_0\,|\,\overline{Z}_1\rangle|\,\overline{Z}_{j+1}\rangle\Big)\hat{\chi}(t), \quad j = 2, 3, \cdots, \end{cases} \tag{7.30}$$

where $|\,\overline{Z}_j\rangle = i(\hat{Z}_j)$. The trivial equation $d\varepsilon/dt = 0$ is added to regard $\varepsilon = K - K_c$ as a dependent variable. The operator T_{10} is defined by (6.3), in which K is replaced by K_c. Note that T_{10} has resonance poles on the imaginary axis.

In what follows, we denote an element of the space $\mathbf{R} \times \prod_{k=1}^{\infty} F_p'$ by $z = (z_0, z_1, z_2, \cdots)$, where $z_0 \in \mathbf{R}$ and $(z_1, z_2, \cdots) \in \prod_{k=1}^{\infty} F_p'$. We also denote it as $z = (z_k)_{k=1}^{\infty}$. A metric on this space is defined as follows: The dual space $F_p(\beta, n)'$ of $F_p(\beta, n)$ is a Banach space with the norm $||\xi||_{\beta,n}^{*} = \sup_{||\phi||_{\beta,n}=1} |\langle \phi\,|\,\xi\rangle|$. The projective limit $F_p(\beta)' = \varprojlim F_p(\beta, n)'$ is a complete metric space with the metric d_β defined by Eq.(5.41). Next, for the projective limit $F_p' = \varprojlim F_p(\beta)'$ we introduce the metric by

$$d(\xi, \zeta) = \sum_{\beta=0}^{\infty} \frac{1}{P(\beta)} \frac{d_\beta(\xi, \zeta)}{1 + d_\beta(\xi, \zeta)}, \tag{7.31}$$

where $\{P(\beta)\}_{\beta=0}^{\infty}$ is a certain sequence of positive numbers such that $\sum_{\beta=0}^{\infty} 1/P(\beta)$ converges. This defines the same projective topology as that induced by the metric (5.42). The constants $P(\beta)$ will be determined in Sec.7.4 so that $d(\xi, 0)$ plays a similar role to a norm. Define d_∞ to be

$$d_\infty(z,z') = \sup_{k\geq 0} d(z_k, z'_k), \quad z = (z_0, z_1, \cdots) \in \mathbf{R}\times\prod_{k=1}^{\infty} F'_p, \tag{7.32}$$

where $d(z_k, z'_k)$ for $k \geq 1$ is the distance on F'_p defined as above, and $d(z_0, z'_0) = |z_0 - z'_0|$ for $z_0, z'_0 \in \mathbf{R}$. Let $\mathcal{G}$ be a subspace of $\mathbf{R}\times\prod_{k=1}^{\infty} F'_p$ consisting of elements $z = (z_0, z_1, \cdots)$ such that $\sup_{k\geq 0} d(z_k, 0)$ is finite. With this metric d_∞, $\mathcal{G}$ is a complete metric vector space. Metric vector spaces and definitions of the metrics used in this section are listed in Table 2.

$F_p(\beta, n)$	Banach space: $\lVert\phi\rVert_{\beta,n} = \sup_{-1/n\leq \mathrm{Im}(z)\leq p+1/n} \lvert\phi(z)\rvert e^{-\beta\lvert z\rvert}$
$F_p(\beta, n)'$	Banach space: $\lVert\xi\rVert^*_{\beta,n} = \sup_{\lVert\phi\rVert_{\beta,n}=1} \lvert\langle\phi\,\vert\,\xi\rangle\rvert$
$F_p(\beta)' = \varprojlim F_p(\beta, n)'$	$d_\beta(\xi,\zeta) = \sum_{n=1}^{\infty} \frac{1}{2^n} \frac{\lVert\xi-\zeta\rVert^*_{\beta,n}}{1+\lVert\xi-\zeta\rVert^*_{\beta,n}}$
$F'_p = \varprojlim F_p(\beta)'$	$d(\xi,\zeta) = \sum_{\beta=0}^{\infty} \frac{1}{P(\beta)} \frac{d_\beta(\xi,\zeta)}{1+d_\beta(\xi,\zeta)}$
$\mathcal{G}\subset\mathbf{R}\times\prod_{k=1}^{\infty} F'_p$	$d_\infty(z,z') = \sup_{k\geq 0} d(z_k, z'_k)$
$\mathcal{F},\ i(\mathcal{F})$	$\mathcal{F}$ is a subspace of $\prod_{k=1}^{\infty} F_p$ satisfying (F1) to (F3); $i(\mathcal{F})$ is its inclusion into $\prod_{k=1}^{\infty} F'_p$. $\mathbf{R}\times i(\mathcal{F})\subset\mathcal{G}$.
$X^{(-m)}\ (= \mathbf{R}\times\overline{i(\mathcal{F})})$	$D_m(z,z') = \kappa^m d_\infty(z,z')$
$X\subset\prod_{m=0}^{\infty} X^{(-m)}$	$D(u,v) = \sup_{m\geq 0} D_m(u^{(-m)}, v^{(-m)})$, $u = (u^{(0)}, u^{(-1)}, \cdots) \in \prod_{m=0}^{\infty} X^{(-m)}$, $u^{(-m)} = (u_k^{(-m)})_{k=0}^{\infty} \in X^{(-m)}$

Table 2: Metric vector spaces used in Section 7. Definitions of the spaces $X^{(-m)}$, X and the constant κ will be given in Section 7.4.

By the definition, it is easy to verify that

$$d_\infty(z, \pm z') = d_\infty(z\mp z', 0) \leq d_\infty(z, 0) + d_\infty(z', 0). \tag{7.33}$$

A sequence $\{z^{(m)} = (z_k^{(m)})_{k=0}^{\infty}\}_{m=1}^{\infty}$ in $\mathcal{G}$ converges to $x = (x_k)_{k=0}^{\infty}$ if and only if $z_0^{(m)} \to x_0$ on $\mathbf{R}$ and $\lVert z_k^{(m)} - x_k\rVert^*_{\beta,n} \to 0$ uniformly in $k \geq 1$ for every $\beta \geq 0$ and $n \geq 1$. On the other hand, since a weakly convergent series in F'_p also converges with respect to the metric d, a sequence $\{z^{(m)} = (z_k^{(m)})_{k=0}^{\infty}\}_{m=1}^{\infty}$ in $\mathcal{G}$ converges to $x = (x_k)_{k=0}^{\infty}$ if and only if $z_0^{(m)} \to x_0$ on $\mathbf{R}$ and $\langle\phi\,\vert\,z_k^{(m)}\rangle \to \langle\phi\,\vert\,x_k\rangle$ uniformly in $k \geq 1$ for every $\phi \in F_p$. It is notable that a mapping $f = (f_0, f_1, \cdots) : \mathcal{G} \to \mathcal{G}$ is continuous with respect to the metric d_∞ if f_k is sequentially continuous with respect to the weak dual topology uniformly in k. Indeed, let $\{z^{(m)}\}_{m=1}^\infty$ be a convergent sequence with respect to d_∞, which satisfies $d_\infty(z^{(m)}, x) \to 0$ as $m \to \infty$. Since the weak dual topology is weaker than that induced by d_∞, $\{z^{(m)}\}_{m=1}^\infty$ weakly converges to x; that is, $z_0^{(m)} \to x_0$ on $\mathbf{R}$ and $\langle \phi \,|\, z_k^{(m)} \rangle \to \langle \phi \,|\, x_k \rangle$ for every $\phi \in F_p$. By the assumption, $f_k(z^{(m)})$ weakly converges to $f_k(x)$. Since F_p is Montel, it also converges with respect to d_∞: $d_\infty(f_k(z^{(m)}), f_k(x)) \to 0$. Since a metric space satisfies the first axiom of countability, this means that f is continuous with respect to the metric d_∞. This fact will be used when proving that the semiflow generated by (7.30) is continuous in initial conditions.

For locally convex topological vector spaces X and Y, a mapping $f : X \to Y$ is called Gâteaux differentiable at $x \in X$ if there exists a map $df(x) : X \to Y$ such that the map G defined by

$$\begin{cases} G(t, x') = \dfrac{f(x + tx') - f(x)}{t} - df(x)x', \quad t \in \mathbf{R},\ x' \in X, \\ G(0, x') = 0, \end{cases} \tag{7.34}$$

is continuous at $(0, 0) \in \mathbf{R} \times X$. Further, if $df(x)$ is linear and if $f : X \to Y$ and $df : X \times X \to Y$ are continuous, f is called C^1 or Fréchet differentiable. Note that f can be Gâteaux differentiable even if it is not continuous (for example, a linear operator is always Gâteaux differentiable). For our situation, it is convenient to use the weak dual topology. As was mentioned in the previous paragraph, we can show that a mapping $f = (f_0, f_1, \cdots) : \mathcal{G} \to \mathcal{G}$ is Gâteaux differentiable with respect to the metric d_∞ if there exist maps $df_k(z)$ such that

$$\lim_{t \to 0} \left| \frac{\langle \phi \,|\, f_k(z + tz') - f_k(z) \rangle}{t} - \langle \phi \,|\, df_k(z)z' \rangle \right| \to 0, \tag{7.35}$$

for any $\phi \in F_p$ uniformly in k. Further, if $df_k(z)$ are linear and if f_k and df_k are sequentially continuous with respect to the weak dual topology uniformly in k, then f is C^1 with respect to the metric d_∞.

Let $i(\mathcal{F})$ be the subspace of $\prod_{k=1}^\infty F_p'$ consisting of elements of the form

$$(i(Z_1), i(Z_2), \cdots), \quad \text{where } (Z_1, Z_2, \cdots) \in \mathcal{F}.$$

Due to Prop.7.2 (ii), $\mathbf{R} \times i(\mathcal{F}) \subset \mathcal{G}$. Thus with the distance d_∞, $\mathbf{R} \times i(\mathcal{F})$ is a metric vector subspace of $\mathcal{G}$, and the closure $\overline{\mathbf{R} \times i(\mathcal{F})}$ is a complete metric vector space.

Now we specify the function $\hat{\chi}(t)$. Let $\mathbf{E}_c$ be the generalized center subspace of T_{10}. Put

$$\hat{\hat{\mathbf{E}}}_c = \mathbf{R} \times \mathbf{E}_c \times \{0\} \times \{0\} \times \cdots \subset \mathbf{R} \times \prod_{k=1}^\infty F_p'. \tag{7.36}$$

Let $P_c : \mathbf{R} \times \prod_{k=1}^\infty F_p' \to \hat{\hat{\mathbf{E}}}_c$ be the projection to $\hat{\hat{\mathbf{E}}}_c$ defined by

$$P_c = (id_{\mathbf{R}},\ \Pi_c,\ 0, 0, \cdots), \tag{7.37}$$

($id_{\mathbf{R}}$ is the identity on $\mathbf{R}$) and $P_s = id - P_c$ the projection to the complement of $\hat{\hat{\mathbf{E}}}_c$. Because of Lemma 5.17 and Prop.5.20, P_c and P_s are continuous on $\mathbf{R}\times i(\mathcal{F})$ and $\hat{\hat{\mathbf{E}}}_c$ is included in the closure $\mathbf{R}\times\overline{i(\mathcal{F})}$. Let $\chi(t)$ be a C^∞ function such that $\chi(t)\equiv 1$ when $0\leq t\leq 1$, $0\leq\chi(t)\leq 1$ when $1\leq t\leq 2$, and $\chi(t)\equiv 0$ when $t\geq 2$. Taking a small positive constant δ_1, we replace $\hat{\chi}(t)$ in (7.30) by

$$\hat{\chi}(t) := \chi\left(\frac{\|P_c z\|_{E_c}}{\delta_1}\right)\cdot\chi(|\eta(t)|), \tag{7.38}$$

where

$$z(t) = (\varepsilon,\, |\overline{Z_1(t,\cdot)}\rangle, |\overline{Z_2(t,\cdot)}\rangle, \cdots) \in \mathbf{R}\times\prod_{k=1}^{\infty} F'_p, \tag{7.39}$$

and $\|\cdot\|_{E_c}$ is a norm on $\hat{\hat{\mathbf{E}}}_c$ defined as follows: An element $y\in\hat{\hat{\mathbf{E}}}_c$ is denoted by $y = (y_0, y_1, 0, \cdots)$ with $y_1 = \sum_{j=0}^{n}\alpha_j|\mu_j\rangle$. Then, $\|y\|_{E_c}$ is defined to be

$$\|y\|_{E_c} = (|y_0|^2 + |\alpha_0|^2 + \cdots + |\alpha_n|^2)^{1/2}. \tag{7.40}$$

Eq.(5.69) shows that $\Pi_c|\overline{Z}_1\rangle$ is given as

$$\Pi_c|\overline{Z}_1\rangle = \sum_{j=0}^{n}\alpha_j|\mu_j\rangle, \quad \alpha_j = \frac{K}{2\overline{D}_j}\langle\overline{Z_1\,|\,\mu_j}\rangle. \tag{7.41}$$

With these α_j, $\|P_c z\|_{E_c}$ is given by

$$\|P_c z\|_{E_c} = (\varepsilon^2 + |\alpha_0|^2 + \cdots + |\alpha_n|^2)^{1/2}, \tag{7.42}$$

for $z = (\varepsilon, |\overline{Z_1}\rangle, \cdots)\in\mathbf{R}\times i(\mathcal{F})$. Since $\hat{\hat{\mathbf{E}}}_c$ is a finite dimensional vector space, the topology on $\hat{\hat{\mathbf{E}}}_c$ induced by $\|\cdot\|_{E_c}$ is equivalent to that induced by the metric d_∞. With this $\hat{\chi}(t)$, we can prove the existence of solutions of Eq.(7.30).

Proposition 7.3. Eq.(7.30) with $\hat{\chi}(t)$ given by (7.38) generates a C^1 semiflow $\tilde{\varphi}_t$ on $\mathbf{R}\times i(\mathcal{F})$. That is, for a given initial condition $z\in\mathbf{R}\times i(\mathcal{F})$, Eq.(7.30) has a unique solution denoted by $\tilde{\varphi}_t(z)$, which is C^1 in z, on $\mathbf{R}\times i(\mathcal{F})$ for any $t\geq 0$.

Proof. At first, we show the existence of solutions of Eq.(7.11) on $\mathcal{F}$. For a given initial condition $(\hat{Z}_1(0,\,\cdot\,),\cdots)\in\mathcal{F}$, there exists a signed measure $\hat{h}(\theta,\omega)$ satisfying (7.14). Such a $\hat{h}$ is uniquely determined because there is a one to one correspondence between a measure on S^1 and its Fourier coefficients (see Shohat and Tamarkin [40]). Thus the existence of a solution of Eq.(7.11) follows from the existence of a solution of Eq.(7.9) with the initial condition $\hat{\rho}_0 = \hat{h}(\theta,\omega)$. Recall that Eq.(7.9) is rewritten as the integro-ODE (7.12) and (7.13). A proof of the existence of solutions of (7.12), (7.13) for any $t\geq 0$ is done by the standard iteration method. Since the proof needs a long calculation, it is left to Appendix B, in which we will also prove that $x(t,s;\theta,\omega)$ is analytic in θ and ω. Once a solution $\hat{\rho}_t$ of (7.9) is obtained, a solution of (7.11) is given through (7.10). Then, Prop.7.2

is applied to show that solutions of Eq.(7.11) are included in $\mathcal{F}$. Note that when $\hat{\chi}$ is given as (7.38), (7.11) becomes an autonomous system. Therefore, solutions define a semiflow on $\mathcal{F}$. This implies that the dual Eq.(7.30) generates a semiflow $\tilde{\varphi}_t$ on $\mathbf{R}\times i(\mathcal{F})$. The proof of smoothness of $\tilde{\varphi}_t$ will be done in Appendix B. It is also proved by the iteration method. ■

The semiflow is also denoted componentwise as

$$\tilde{\varphi}_t(z) = (z_0,\ \tilde{\varphi}^1_t(z), \tilde{\varphi}^2_t(z), \cdots), \quad z = (z_0, z_1, \cdots). \tag{7.43}$$

By virtue of the variation-of-constant formula (see Eq.(5.25)), $\tilde{\varphi}^j_t$ proves to be of the form

$$\tilde{\varphi}^j_t(z_0, z_1, \cdots) = (e^{T_j t})^{\times} z_j + \tilde{g}^j_t(z_0, z_1, \cdots), \quad j = 1, 2, \cdots, \tag{7.44}$$

where $\tilde{g}^j_t$ are nonlinear terms. Now we introduce another localization factor. Let $\delta_2 > 0$ be a sufficiently small positive number. By using a function $\chi(t)$ above, we multiply the function $\chi(d_\infty(P_s z, 0)/\delta_2)$ to the nonlinearity $\tilde{g}^j_t$, and define a perturbed map $\varphi_t = (z_0, \varphi^1_t, \varphi^2_t, \cdots)$ to be

$$\varphi^j_t(z_0, z_1, \cdots) = (e^{T_j t})^{\times} z_j + \tilde{g}^j_t(z_0, z_1, \cdots) \cdot \chi\Bigl(\frac{d_\infty(P_s z, 0)}{\delta_2}\Bigr), \tag{7.45}$$

for $j = 1, 2, \cdots$. Put

$$g^j_t(z_0, z_1, \cdots) = \tilde{g}^j_t(z_0, z_1, \cdots) \cdot \chi\Bigl(\frac{d_\infty(P_s z, 0)}{\delta_2}\Bigr).$$

Fix a positive number $\tau > 0$, and put

$$L = (id_{\mathbf{R}}, (e^{T_1 0\tau})^{\times}, (e^{T_2 \tau})^{\times}, \cdots), \quad g = (0,\ g^1_\tau, g^2_\tau, \cdots).$$

Then, the time τ map φ_τ of φ_t is denoted as

$$\varphi_\tau : \mathbf{R}\times i(\mathcal{F}) \to \mathbf{R}\times i(\mathcal{F}), \quad \varphi_\tau(z) = Lz + g(z). \tag{7.46}$$

This is the desired localization of the semiflow of the original system (6.2). By Prop.7.3, $\tilde{g}^j_t$ is a C^1 mapping on $\mathbf{R}\times i(\mathcal{F})$. Since $d_\infty(\cdot, 0)$ and P_s are continuous on $\mathbf{R}\times i(\mathcal{F})$, $g : \mathbf{R}\times i(\mathcal{F}) \to \mathbf{R}\times i(\mathcal{F})$ is also continuous on $\mathbf{R}\times i(\mathcal{F})$. Hence, the map φ_τ is continuously extended to the map on the closure $\mathbf{R}\times \overline{i(\mathcal{F})}$. Unfortunately, the distance $d_\infty(z, 0)$ is not C^1 in z. However, on the region such that $d_\infty(P_s z, 0) \leq \delta_2$ or $d_\infty(P_s z, 0) \geq 2\delta_2$, g is a C^1 mapping because $\chi(d_\infty(P_s z, 0)/\delta_2)$ becomes a constant. To see that $g(z) \sim O(z^2)$ as $z \to 0$, let us simply denote Eq.(7.30) by $d|\overline{Z}_j\rangle/dt = T^{\times}_j|\overline{Z}_j\rangle + G_j(z)$, where $z = z(t)$ is given as Eq.(7.39). Then, $\tilde{g}^j_t$ in Eq.(7.44) is written as

$$\tilde{g}^j_t(z_0, z_1, \cdots) = \int_0^t (e^{T_j(t-s)})^{\times} G_j(z(s))ds,$$

with the initial condition $z(0) = (z_0, z_1, \cdots)$. Note that G is C^∞ and $G(z) \sim O(z^2)$ as $z \to 0$. Since solutions $z(t)$ is C^1 in initial conditions, we obtain $G(z(s)) \sim O(z(0)^2)$ as $z(0) \to 0$. This shows that $\tilde{g}^j_t$ is of order $O(z^2)$ as $z \to 0$ on $\mathbf{R}\times \overline{i(\mathcal{F})}$ with respect to the metric d_∞.

When $||P_c z||_{E_c} \leq \delta_1$, $|\eta(t)| \leq 1$ and $d_\infty(P_s z, 0) \leq \delta_2$, then $\chi\,(||P_c z||_{E_c}/\delta_1) = 1$, $\chi(|\eta(t)|) = 1$ and $\chi(d_\infty(P_s z, 0)/\delta_2) = 1$. Thus Eq.(7.46) coincides with the time τ map of the semiflow of the original system (6.2). When $||P_c z||_{E_c} \geq 2\delta_1$ or $d_\infty(P_s z, 0) \geq 2\delta_2$, then $\chi\,(||P_c z||_{E_c}/\delta_1) \cdot \chi(d_\infty(P_s z, 0)/\delta_2) = 0$. In this case, $g = 0$ and Eq.(7.46) is reduced to the linear map. Therefore, by taking δ_1 and δ_2 sufficiently small, the Lipschitz constant of g

$$\mathrm{Lip}(g) := \sup_{z, z' \in \mathbf{R} \times \overline{i(\mathcal{F})}} \frac{d_\infty(g(z), g(z'))}{d_\infty(z, z')} \tag{7.47}$$

can be assumed to be sufficiently small.

Remark. We introduced the factors for localization in two steps. The one $\chi\,(||P_c z||_{E_c}/\delta_1)$ is multiplied to the nonlinearity of the equation (7.30), and the other $\chi(d_\infty(P_s z, 0)/\delta_2)$ is multiplied to the nonlinearity of the semiflow (7.44). The reason is that if we multiply both of them to the equation (7.30), then the proof of the existence of solutions for (7.30) (which is completed in Appendix B) becomes too difficult; $\chi\,(||P_c z||_{E_c}/\delta_1)$ is essentially a finite dimensional perturbation, although $\chi(d_\infty(P_s z, 0)/\delta_2)$ includes infinite dimensional terms $P_s z$. On the other hand, if we multiply both of them to the nonlinearity of the semiflow of the original system (6.2), then a center manifold of the resultant perturbed mapping does not coincide with a center manifold of the original system (6.2) because the perturbed mapping is not a semiflow for any differential equations in general (i.e. the property $\varphi_{t+s} = \varphi_t \circ \varphi_s$ is violated because of the perturbation for the semiflow), see Krisztin [24] for details. However, if we introduce these factors in two steps as above, a local center manifold of the original system is correctly obtained as follows: In Sec.7.4, we will prove the existence of a center manifold for the map (7.46). We will show that if δ_1 is sufficiently small, the center manifold is included in the "strip" $d_\infty(P_s z, 0) < \delta_2$. Since the map (7.46) is the same as (7.44), which is a semiflow of the system (7.30), when $d_\infty(P_s z, 0) < \delta_2$, the obtained center manifold is a center manifold of the system (7.30). When $||P_c z||_{E_c} < \delta_1$, (7.30) is reduced to the original system (6.2). Therefore, a local center manifold of (6.2) is obtained as a restriction of the center manifold of (7.30) to the region $||P_c z||_{E_c} < \delta_1$, see Fig.12.

7.4 Proof of the center manifold theorem

Let us prove that the mapping φ_τ defined in Eq.(7.46) has a center manifold, which gives a local center manifold for the original system (6.2). The strategy of the proof is the same as that in Chen, Hale and Bin [9], in which the existence of center manifolds is proved for mappings on Banach spaces. At first, we need the next lemma to treat the metric d_∞ as a norm.

Lemma 7.4. For $u = (u_0, u_1, \cdots) \in \mathbf{R} \times \overline{i(\mathcal{F})}$, suppose that there exists a positive constant δ_3 such that $||u_j||^*_{0,1} \leq \delta_3$ for $j = 1, 2, \cdots$. If constants $\{P(\beta)\}_{\beta=0}^\infty$ are sufficiently large, there exist positive constants A_c and $A_s = A_s(\delta_3)$ such that the inequalities

$$d_\infty(L^m P_c u\,,\, 0) \leq A_c d_\infty(u, 0), \quad m = 0, \pm 1, \pm 2, \cdots, \tag{7.48}$$

and

$$d_\infty(L^m P_s u\,,\, 0) \leq A_s e^{-am\tau} d_\infty(u, 0), \quad m = 0, 1, 2, \cdots, \tag{7.49}$$


$\hat{\hat{\boldsymbol{E}}}_c^{\perp}$
$2\delta_2$
δ_2
δ_1
$2\delta_1$
$\hat{\hat{\boldsymbol{E}}}_c$

Fig. 12: A center manifold (the black curve) for the map (7.46) coincides with that of the semiflow (7.44). The center manifold of the semiflow (7.44) coincides with that of the original system (6.2) in the region $||P_c z||_{E_c} < \delta_1$.

hold, where $a > 0$ is the constant appeared in Prop.5.19.

Note that since P_c is a projection to the finite dimensional vector space $\hat{\hat{\mathbf{E}}}_c$, the linear operator L restricted to $\hat{\hat{\mathbf{E}}}_c$ is equivalent to an exponential of a matrix. Thus the inverse L^{-1} exists on $\hat{\hat{\mathbf{E}}}_c$ and $L^{-m}P_c$ is well-defined.

Proof. For $u = (u_0, u_1, \cdots) \in \mathbf{R} \times \overline{i(\mathcal{F})}$, $L^m P_c u$ is given by

$$L^m P_c u = (u_0\,,\,(e^{T_{10}m\tau})^{\times}\Pi_c u_1\,,\,0\,,0\,,\cdots).$$

Since $u_1 \in \overline{i(V)} \subset \overline{i(W_p(3,0))}$, Eq.(5.79) is applied to yield

$$d_\beta((e^{T_{10}m\tau})^{\times}\Pi_c u_1\,,\,0) = \sum_{n=1}^{\infty}\frac{1}{2^n}\frac{||(e^{T_{10}m\tau})^{\times}\Pi_c u_1||^*_{\beta,n}}{1+||(e^{T_{10}m\tau})^{\times}\Pi_c u_1||^*_{\beta,n}} \leq \sum_{n=1}^{\infty}\frac{1}{2^n}\frac{L_\beta||u_1||^*_{\beta,n}}{1+L_\beta||u_1||^*_{\beta,n}},$$

where $L_\beta = L_{3,0,\beta}$. We can assume without loss of generality that $L_\beta \geq 1$. Then,

$$d_\beta((e^{T_{10}m\tau})^{\times}\Pi_c u_1\,,\,0) \leq L_\beta \sum_{n=1}^{\infty}\frac{1}{2^n}\frac{||u_1||^*_{\beta,n}}{1+||u_1||^*_{\beta,n}} = L_\beta d_\beta(u_1, 0).$$

Eq.(5.74) provides

$$d_0(u_1, 0) = \sum_{n=1}^{\infty}\frac{1}{2^n}\frac{||u_1||^*_{0,n}}{1+||u_1||^*_{0,n}} \leq \sum_{n=1}^{\infty}\frac{1}{2^n}\frac{||u_1||^*_{\beta,n}}{1+||u_1||^*_{\beta,n}} = d_\beta(u_1, 0),$$

and

$$d_\beta(u_1, 0) = \sum_{n=1}^{\infty}\frac{1}{2^n}\frac{||u_1||^*_{\beta,n}}{1+||u_1||^*_{\beta,n}} \leq \sum_{n=1}^{\infty}\frac{1}{2^n}\frac{Q(\beta)||u_1||^*_{0,n}}{1+||u_1||^*_{0,n}} = Q(\beta)d_0(u_1, 0).$$

By using them, $d((e^{T_{10}m\tau})^\times \Pi_c u_1\,,\,0)$ is calculated as

$$\begin{aligned} d((e^{T_{10}m\tau})^\times \Pi_c u_1\,,\,0) &= \sum_{\beta=0}^{\infty} \frac{1}{P(\beta)} \frac{d_\beta((e^{T_{10}m\tau})^\times \Pi_c u_1\,,\,0)}{1+d_\beta((e^{T_{10}m\tau})^\times \Pi_c u_1\,,\,0)} \\ &\leq \sum_{\beta=0}^{\infty} \frac{1}{P(\beta)} \frac{L_\beta d_\beta(u_1,0)}{1+L_\beta d_\beta(u_1,0)} \\ &\leq \sum_{\beta=0}^{\infty} \frac{L_\beta Q(\beta)}{P(\beta)} \frac{d_0(u_1,0)}{1+d_0(u_1,0)}, \end{aligned}$$

where we choose a sequence $\{P(\beta)\}_{\beta=0}^{\infty}$ so that $\sum_{\beta=0}^{\infty} L_\beta Q(\beta)/P(\beta)$ converges. Then,

$$\begin{aligned} d((e^{T_{10}m\tau})^\times \Pi_c u_1\,,\,0) &\leq \sum_{\beta=0}^{\infty} \frac{L_\beta Q(\beta)}{P(\beta)} \cdot \Big(\sum_{\beta=0}^{\infty} \frac{1}{P(\beta)}\Big)^{-1} \cdot \sum_{\beta=0}^{\infty} \frac{1}{P(\beta)} \frac{d_0(u_1,0)}{1+d_0(u_1,0)} \\ &\leq \sum_{\beta=0}^{\infty} \frac{L_\beta Q(\beta)}{P(\beta)} \cdot \Big(\sum_{\beta=0}^{\infty} \frac{1}{P(\beta)}\Big)^{-1} \cdot \sum_{\beta=0}^{\infty} \frac{1}{P(\beta)} \frac{d_\beta(u_1,0)}{1+d_\beta(u_1,0)} \\ &= A_c d(u_1,0), \quad A_c := \sum_{\beta=0}^{\infty} \frac{L_\beta Q(\beta)}{P(\beta)} \cdot \Big(\sum_{\beta=0}^{\infty} \frac{1}{P(\beta)}\Big)^{-1}. \end{aligned} \tag{7.50}$$

Thus we obtain

$$d_\infty(L^m P_c u, 0) \leq \sup\{u_0, A_c d(u_1,0)\} \leq \sup\{A_c u_0, A_c d(u_1,0)\} = A_c d_\infty(u,0),$$

where we suppose that $A_c \geq 1$. Note that we did not use the condition $||u_j||^*_{0,1} \leq \delta_3$ for Eq.(7.48).

Next, $L^m P_s u$ is given by

$$L^m P_s u = (0,\ (e^{T_{10}m\tau})^\times \Pi_s u_1\,,\ (e^{T_2 m\tau})^\times u_2\,,\ \cdots).$$

Eq.(5.80) is used to yield

$$\begin{aligned} d_\beta((e^{T_{10}m\tau})^\times \Pi_s u_1, 0) &= \sum_{n=1}^{\infty} \frac{1}{2^n} \frac{||(e^{T_{10}m\tau})^\times \Pi_s u_1||^*_{\beta,n}}{1+||(e^{T_{10}m\tau})^\times \Pi_s u_1||^*_{\beta,n}} \\ &\leq \sum_{n=1}^{\infty} \frac{1}{2^n} \frac{M_\beta e^{-am\tau}||u_1||^*_{\beta,n}}{1+M_\beta e^{-am\tau}||u_1||^*_{\beta,n}} \leq M_\beta e^{-am\tau} \sum_{n=1}^{\infty} \frac{1}{2^n} \frac{||u_1||^*_{\beta,n}}{1+e^{-am\tau}||u_1||^*_{\beta,n}}, \end{aligned}$$

where $M_\beta := M_{3,0,\beta}$ is assumed to be larger than 1. Note that the condition $||u_j||^*_{0,1} \leq \delta_3$ yields $||u_j||^*_{\beta,n} \leq Q(\beta)\delta_3$. When $||u_j||^*_{\beta,n} \leq Q(\beta)\delta_3$, putting $A'_\beta(\delta_3) = (1+Q(\beta)\delta_3)$ provides

$$\frac{1}{1+e^{-am\tau}||u_1||^*_{\beta,n}} \leq A'_\beta(\delta_3) \frac{1}{1+||u_1||^*_{\beta,n}},$$

uniformly in $m = 0, 1, \cdots$. Therefore, we obtain

$$d_\beta((e^{T_{10}m\tau})^\times \Pi_s u_1, 0) \le A'_\beta(\delta_3) M_\beta e^{-am\tau} d_\beta(u_1, 0).$$

By the same way as in Eq.(7.50), we can verify that there exists a constant $A_s(\delta_3) > 0$ such that

$$d((e^{T_{10}m\tau})^\times \Pi_s u_1, 0) \le A_s(\delta_3) e^{-am\tau} d(u_1, 0).$$

In this calculation, constants $P(\beta)$ are chosen sufficiently large as before. Similarly, Eq.(5.81) shows that

$$d((e^{T_j m\tau})^\times u_j, 0) \le A_s(\delta_3) e^{-am\tau} d(u_j, 0),$$

holds for every $j = 2, 3, \cdots$. Note that the constant A_s can be taken so that it is independent of j because the constant $N_{C,\alpha,\beta}$ in Eq.(5.81) is independent of j. Thus $d_\infty(L^m P_s u, 0)$ satisfies Eq.(7.49). ■

If $||P_c u||_{E_c} \ge 2\delta_1$ or $d_\infty(P_s u, 0) \ge 2\delta_2$, then $g(u) = 0$. Thus there exists $D = D(\delta_1, \delta_2)$ such that the j-th component of g satisfies $d(g(u)_j, 0) \le D$ for every $j = 1, 2, \cdots$. This shows that there exists $\delta_3 = \delta_3(\delta_1, \delta_2)$ such that $||g(u)_j||^*_{0,1} \le \delta_3$ for all j. In what follows, we fix δ_3 in Lemma 7.4 so that g satisfies $||g(u)_j||^*_{0,1} \le \delta_3$ for all j. Then, Eq.(7.49) is applicable to $g(u)$. Note that Eq.(7.48) holds for any $\mathbf{R} \times \overline{i(\mathcal{F})}$ without the assumption.

Lemma 7.5. For a sequence $\{u^{(-m)} = (u_0^{(-m)}, u_1^{(-m)}, \cdots)\}_{m=0}^\infty \subset \mathbf{R} \times \overline{i(\mathcal{F})}$, suppose that

$$L^q P_s u^{(-q)} \to 0 \tag{7.51}$$

as $q \to \infty$, and that there exist constants $C > 0$ and a sufficiently small $b > 0$ such that

$$d_\infty(u^{(-m)}, 0) \le C e^{bm\tau} \tag{7.52}$$

holds for every $m = 0, 1, \cdots$. Then, $\{u^{(-m)}\}_{m=0}^\infty$ satisfies

$$u^{(-m)} = \varphi_\tau(u^{(-m-1)}) = L u^{(-m-1)} + g(u^{(-m-1)}), \quad m = 0, 1, 2, \cdots, \tag{7.53}$$

if and only if it satisfies the equation

$$u^{(-m)} = L^{-m} P_c u_0 - \sum_{k=1}^{m} L^{k-m-1} P_c g(u^{(-k)}) + \sum_{k=m+1}^{\infty} L^{k-m-1} P_s g(u^{(-k)}), \quad m = 0, 1, 2, \cdots. \tag{7.54}$$

Eq.(7.53) means that $\{u^{(-m)}\}_{m=0}^\infty = (u^{(0)}, u^{(-1)}, \cdots)$ is a negative semi-orbit of the mapping φ_τ. Eq.(7.54) is called the *Lyapunov-Perron equation* [9].

Proof. Suppose that $\{u^{(-m)}\}$ satisfies Eq.(7.53). By iterating Eq.(7.53), we obtain

$$\begin{aligned} u^{(0)} &= P_c u^{(0)} + P_s u^{(0)} \\ &= P_c u^{(0)} + L P_s u^{(-1)} + P_s g(u^{(-1)}) \\ &= P_c u^{(0)} + L^2 P_s u^{(-2)} + L P_s g(u^{(-2)}) + P_s g(u^{(-1)}) \\ &\vdots \\ &= P_c u^{(0)} + L^q P_s u^{(-q)} + \sum_{k=1}^{q} L^{k-1} P_s g(u^{(-k)}), \quad q = 0, 1, 2, \cdots. \end{aligned}$$

In a similar manner, we obtain

$$u^{(-m)} = L^{-m}P_c u_0 - \sum_{k=1}^{m} L^{k-m-1}P_c g(u^{(-k)}) + L^{q-m}P_s u^{(-q)} + \sum_{k=m+1}^{q} L^{k-m-1}P_s g(u^{(-k)}), \quad (7.55)$$

for $q = m, m+1, \cdots$ and $m = 0, 1, 2, \cdots$. By the assumption, $L^{q-m}P_s u^{(-q)} \to 0$ as $q \to 0$. Next thing to do is to show that $\sum_{k=m+1}^{q} L^{k-m-1}P_s g(u^{(-k)})$ converges as $q \to \infty$. Eq.(7.49) is applicable to $g(u^{(-m)})$ to yield

$$\begin{aligned} d_\infty(L^{k-m-1}P_s g(u^{(-k)}), 0) &\le A_s e^{-a(k-m-1)\tau} d_\infty(g(u^{(-k)}), 0) \\ &\le \mathrm{Lip}(g) A_s e^{-a(k-m-1)\tau} d_\infty(u^{(-k)}, 0) \\ &\le \mathrm{Lip}(g) A_s C e^{-(a-b)k\tau} e^{a(m+1)\tau}, \end{aligned} \quad (7.56)$$

which shows that $L^{k-m-1}P_s g(u^{(-k)})$ decays exponentially as $k \to \infty$ when $a > b$. Thus taking the limit $q \to \infty$ in Eq.(7.55) yields Eq.(7.54).

Conversely, suppose that $\{u^{(-m)}\}$ satisfies Eq.(7.54). Because of the assumption Eq.(7.52), the series $\sum_{k=m+1}^{\infty} L^{k-m-1}P_s g(u^{(-k)})$ exists. Replacing m by $m+1$ and using L for Eq.(7.54), we obtain

$$Lu^{(-m-1)} = L^{-m}P_c u^{(0)} - \sum_{k=1}^{m+1} L^{k-m-1}P_c g(u^{(-k)}) + \sum_{k=m+2}^{\infty} L^{k-m-1}P_s g(u^{(-k)}). \quad (7.57)$$

Eq.(7.57) is put together with Eq.(7.54) to yield Eq.(7.53). ■

Let $X^{(-m)}$ $(m = 0, 1, \cdots)$ be copies of the space $\mathbf{R} \times \overline{i(\mathcal{F})}$. Define a metric D_m on $X^{(-m)}$ to be

$$D_m(z, z') = e^{-bm\tau} d_\infty(z, z'), \quad z, z' \in \mathbf{R} \times \overline{i(\mathcal{F})}, \quad (7.58)$$

with a small positive constant b. Let X be a subspace of the product $\prod_{m=0}^{\infty} X^{(-m)}$ consisting of elements $u = (u^{(0)}, u^{(-1)}, u^{(-2)}, \cdots)$ such that $\sup_m D_m(u^{(-m)}, 0)$ is finite. With the metric defined by

$$D(u, v) = \sup_{m \ge 0} D_m(u^{(-m)}, v^{(-m)}), \quad (7.59)$$

X is a complete metric vector space (see Table 2). It is easy to verify the inequality

$$D(u, \pm u') = D(u \mp u', 0) \le D(u, 0) + D(u', 0). \quad (7.60)$$

Motivated by Eq.(7.54), let us define the map $J : X \times \hat{\hat{\mathbf{E}}}_c \to X$ to be

$$\begin{cases} J(u, y) = (J^{(0)}(u, y), J^{(-1)}(u, y), J^{(-2)}(u, y), \cdots), \\ J^{(-m)}(u, y) = L^{-m}y - \sum_{k=1}^{m} L^{k-m-1}P_c g(u^{(-k)}) + \sum_{k=m+1}^{\infty} L^{k-m-1}P_s g(u^{(-k)}). \end{cases} \quad (7.61)$$

If the map $J(\cdot\,, y)$ has a fixed point $u = q(y) = (q^{(0)}(y), q^{(-1)}(y), \cdots)$, $q(y)$ is a solution of the Lyapunov-Perron equation (7.54) with $P_c u_0 = y$. If $q(y)$ satisfies conditions (7.51) and (7.52), Lemma 7.5 shows that $q(y)$ is a negative semi-orbit (that is, it satisfies Eq.(7.53)) for each $y \in \hat{\hat{\mathbf{E}}}_c$. We will see that this $q(y)$ gives a desired center manifold. At first, let us show that J is well-defined.

Lemma 7.6. J is a map from $X \times \hat{\hat{\mathbf{E}}}_c$ into X.

Proof. Let us show that $D_m(J^{(-m)}(u, y), 0)$ is bounded uniformly in $m = 0, 1, \cdots$. It satisfies

$$
\begin{aligned}
&D_m(J^{(-m)}(u, y), 0) = e^{-bm\tau} d_\infty(J^{(-m)}(u, y), 0) \\
&\le e^{-bm\tau} d_\infty(L^{-m} y, 0) + e^{-bm\tau} \sum_{k=1}^{m} d_\infty(L^{k-m-1} P_c g(u^{(-k)}), 0) + e^{-bm\tau} \sum_{k=m+1}^{\infty} d_\infty(L^{k-m-1} P_s g(u^{(-k)}), 0)
\end{aligned}
\tag{7.62}
$$

Eq.(7.48) shows that the first term $e^{-b\tau m} d_\infty(L^m y, 0)$ is bounded uniformly in $m = 0, 1, \cdots$. Similarly, we obtain

$$
d_\infty(L^{k-m-1} P_c g(u^{(-k)}), 0) \le A_c d_\infty(g(u^{(-k)}), 0) \le \mathrm{Lip}(g) A_c d_\infty(u^{(-k)}, 0).
$$

Since $u \in X$, there is a constant $C > 0$ such that $d_\infty(u^{(-k)}, 0) \le C e^{bk\tau}$. Therefore,

$$
e^{-bm\tau} \sum_{k=1}^{m} d_\infty(L^{k-m-1} P_c g(u^{(-k)}), 0) \le \mathrm{Lip}(g) A_c C e^{-bm\tau} \sum_{k=1}^{m} e^{bk\tau} \le \mathrm{Lip}(g) A_c C \frac{e^{b\tau} - e^{b\tau(1-m)}}{e^{b\tau} - 1}
$$

is bounded uniformly in $m = 0, 1, \cdots$. The last term in Eq.(7.62) obviously tends to zero as $m \to \infty$. This proves that $J(u, y) \in X$. ■

Proposition 7.7. If the constants δ_1 and δ_2 are sufficiently small, J is a contraction map on X for each $y \in \hat{\hat{\mathbf{E}}}_c$.

Proof. For $u, u' \in X$, we have

$$
\begin{aligned}
&D_m(J^{(-m)}(u, y), J^{(-m)}(u', y)) = e^{-bm\tau} d_\infty(J^{(-m)}(u, y) - J^{(-m)}(u', y), 0) \\
&\le e^{-bm\tau} \sum_{k=1}^{m} d_\infty(L^{k-m-1} P_c(g(u^{(-k)}) - g(u'^{(-k)})), 0) + e^{-bm\tau} \sum_{k=m+1}^{\infty} d_\infty(L^{k-m-1} P_s(g(u^{(-k)}) - g(u'^{(-k)})), 0).
\end{aligned}
\tag{7.63}
$$

Eqs.(7.48) and (7.47) are used to yield

$$
d_\infty(L^{k-m-1} P_c(g(u^{(-k)}) - g(u'^{(-k)})), 0) \le \mathrm{Lip}(g) A_c d_\infty(u^{(-k)}, u'^{(-k)}).
$$

Similarly, we obtain

$$
d_\infty(L^{k-m-1} P_s(g(u^{(-k)}) - g(u'^{(-k)})), 0) \le \mathrm{Lip}(g) A_s e^{-a(k-m-1)\tau} d_\infty(u^{(-k)}, u'^{(-k)}).
$$

Therefore, we obtain

$$\begin{aligned}
& D_m(J^{(-m)}(u,y),\, J^{(-m)}(u',y)) \\
\le\ & e^{-bm\tau}\sum_{k=1}^{m}\mathrm{Lip}(g)A_c d_\infty(u^{(-k)}, u'^{(-k)}) + e^{-bm\tau}\sum_{k=m+1}^{\infty}\mathrm{Lip}(g)A_s e^{-a(k-m-1)\tau} d_\infty(u^{(-k)}, u'^{(-k)}) \\
\le\ & \mathrm{Lip}(g)A_c\sum_{k=1}^{m} e^{b(k-m)\tau} D_k(u^{(-k)}, u'^{(-k)}) + \mathrm{Lip}(g)A_s\sum_{k=m+1}^{\infty} e^{-a(k-m-1)\tau}e^{b(k-m)\tau} D_k(u^{(-k)}, u'^{(-k)}) \\
\le\ & \mathrm{Lip}(g)\left(A_c\sum_{k=1}^{m} e^{b(k-m)\tau} + A_s\sum_{k=m+1}^{\infty} e^{-a(k-m-1)\tau}e^{b(k-m)\tau}\right)\cdot D(u,u').
\end{aligned}$$

This yields

$$\begin{aligned}
D(J(u,y),\, J(u',y)) &= \sup_{m\ge 0} D_m(J^{(-m)}(u,y),\, J^{(-m)}(u',y)) \\
&\le \mathrm{Lip}(g)\left(A_c\frac{e^{b\tau}}{e^{b\tau}-1} + A_s\frac{e^{b\tau}}{1-e^{(b-a)\tau}}\right)\cdot D(u,u').
\end{aligned}$$

We can take δ_1 and δ_2 sufficiently small so that $\mathrm{Lip}(g)$ becomes sufficiently small and

$$\mathrm{Lip}(g)\left(A_c\frac{e^{b\tau}}{e^{b\tau}-1} + A_s\frac{e^{b\tau}}{1-e^{(b-a)\tau}}\right) < 1 \tag{7.64}$$

holds. This implies that $J(\,\cdot\,, y)$ is a contraction map on X for each $y \in \hat{\hat{\mathbf{E}}}_c$. ■

Remark. The numbers a and b are the same as those in Thm.7.1. The reason we introduced metrics D_m and D is that the center manifold is characterized by the "slow" dynamics whose Lyapunov exponent is smaller than b, see Eq.(7.3). The above condition for $\mathrm{Lip}(g)$ shows that if we take $b > 0$ sufficiently small, $\mathrm{Lip}(g)$ (and thus δ_1 and δ_2) should be small accordingly. Since the open set U in Thm.7.1, in which we can prove the existence of the local center manifold, is determined by δ_1 and δ_2, U also becomes small as a result.

By the contraction principle, $J(\,\cdot\,, y)$ has a unique fixed point $u = q(y)$ on X:

$$\left\{\begin{array}{l}
q(y) = (q^{(0)}(y),\, q^{(-1)}(y),\, q^{(-2)}(y), \cdots), \quad q^{(-m)} : \hat{\hat{\mathbf{E}}}_c \to X^{(-m)}, \\
q^{(-m)}(y) = L^{-m}y - \displaystyle\sum_{k=1}^{m} L^{k-m-1}P_c g(q^{(-k)}(y)) + \sum_{k=m+1}^{\infty} L^{k-m-1}P_s g(q^{(-k)}(y)).
\end{array}\right. \tag{7.65}$$

In particular, $q^{(0)}$ defines a map from $\hat{\hat{\mathbf{E}}}_c$ into $X^{(0)} = \mathbf{R}\times\overline{i(\mathcal{F})}$ given by

$$q^{(0)}(y) = y + \sum_{k=1}^{\infty} L^{k-1}P_s g(q^{(-k)}(y)). \tag{7.66}$$

Since $q(y) \in X$, there exists $C > 0$ such that $D_m(q^{(-m)}(y), 0) = e^{-bm\tau}d_\infty(q^{(-m)}(y), 0) \le C$, which verifies the condition (7.52). Further, Eq.(7.65) shows that

$$L^m P_s q^{(-m)}(y) = \sum_{k=m+1}^{\infty} L^{k-1}P_s g(q^{(-k)}(y)).$$

Since this is a convergent series, $L^m P_s q^{(-m)}(y) \to 0$ as $m \to \infty$, which verifies (7.51). Therefore, Lemma 7.5 is applicable to conclude that $\{q^{(-m)}(y)\}_{m=0}^{\infty}$ is a negative semi-orbit for each y.

Proposition 7.8. For any $m = 0, 1, \cdots$,
(i) $q^{(-m)}(0) = 0$.
(ii) $q^{(-m)} : \hat{\hat{\mathbf{E}}}_c \to X^{(-m)}$ is Lipschitz continuous.
(iii) if $||y||_{E_c} \geq 2\delta_1$, then $q^{(-m)}(y) = L^{-m} y \in \hat{\hat{\mathbf{E}}}_c$.
(iv) $q^{(-m)} : \hat{\hat{\mathbf{E}}}_c \to X^{(-m)}$ is a C^1 mapping. In particular, $q^{(0)} : \hat{\hat{\mathbf{E}}}_c \to \mathbf{R} \times \overline{i(\mathcal{F})}$ is C^1.

Proof. (i) Since $g(0) = 0$, $q^{(-m)}(0) = 0$ satisfies Eq.(7.65).
(ii) For $y, y' \in \hat{\hat{\mathbf{E}}}_c$, we estimate $D_m(q^{(-m)}(y), q^{(-m)}(y'))$. By the same calculation as the proof of Prop.7.7, we obtain

$$D(q(y), q(y')) \leq \frac{A_c}{1 - \mathrm{Lip}(g)\left(A_c \dfrac{e^{b\tau}}{e^{b\tau} - 1} + A_s \dfrac{e^{b\tau}}{1 - e^{(b-a)\tau}}\right)} d_\infty(y, y'). \tag{7.67}$$

This means that $q : \hat{\hat{\mathbf{E}}}_c \to X$ is Lipschitz continuous. In particular, we obtain

$$d_\infty(q^{(-m)}(y), q^{(-m)}(y')) \leq \frac{A_c e^{bm\tau}}{1 - \mathrm{Lip}(g)\left(A_c \dfrac{e^{b\tau}}{e^{b\tau} - 1} + A_s \dfrac{e^{b\tau}}{1 - e^{(b-a)\tau}}\right)} d_\infty(y, y'), \tag{7.68}$$

which proves the proposition.
(iii) Put $y = (y_0, y_1, 0, \cdots)$, and $y_1 = \sum_{j=0}^{n} \alpha_j |\mu_j\rangle \in \mathbf{E}_c$. Then, the assumption implies

$$||y||_{E_c} = |y_0|^2 + |\alpha_0|^2 + |\alpha_1|^2 + \cdots + |\alpha_n|^2 \geq 2\delta_1.$$

On the other hand, $L^{-m} y$ is given by $L^{-m} y = (y_0, (e^{-T_{10} m\tau})^{\times} y_1, 0, \cdots)$, and

$$(e^{-T_{10} m\tau})^{\times} y_1 = \sum_{j=0}^{m} \alpha_j (e^{-T_{10} m\tau})^{\times} |\mu_j\rangle = \sum_{j=0}^{m} \alpha_j e^{-\overline{\lambda}_j m\tau} |\mu_j\rangle.$$

Since $\lambda_j \in \sqrt{-1}\mathbf{R}$, it turns out that $||L^{-m} y||_{E_c} = ||y||_{E_c} \geq 2\delta_1$. By the construction of the nonlinearity g, $g(L^{-m} y) = 0$ if $||L^{-m} y||_{E_c} \geq 2\delta_1$. Therefore, $q^{(-m)}(y) = L^{-m} y$ satisfies Eq.(7.65).
(iv) For $y, y^* \in \hat{\hat{\mathbf{E}}}_c$ and $\kappa \in \mathbf{R}$, put

$$\tilde{q}^{(-m)}(y, y^*, \kappa) = \frac{1}{\kappa}(q^{(-m)}(y + \kappa y^*) - q^{(-m)}(y)). \tag{7.69}$$

Then, it satisfies the equation

$$\begin{aligned} \tilde{q}^{(-m)}(y, y^*, \kappa) \quad = \quad & L^{-m} y^* - \frac{1}{\kappa} \sum_{k=1}^{m} L^{k-m-1} P_c(g(q^{(-k)}(y) + \kappa \tilde{q}^{(-k)}(y, y^*, \kappa)) - g(q^{(-k)}(y))) \\ & + \frac{1}{\kappa} \sum_{k=m+1}^{\infty} L^{k-m-1} P_s(g(q^{(-k)}(y) + \kappa \tilde{q}^{(-k)}(y, y^*, \kappa)) - g(q^{(-k)}(y))), \end{aligned} \tag{7.70}$$

for $\kappa \neq 0$. If $\tilde{q}^{(-m)}(y, y^*, 0)$ exists, it should satisfy

$$\begin{aligned}\tilde{q}^{(-m)}(y, y^*, 0) \;=\; & L^{-m}y^* - \sum_{k=1}^{m} L^{k-m-1} P_c \frac{dg}{dx}(q^{(-k)}(y))\tilde{q}^{(-k)}(y, y^*, 0) \\ & + \sum_{k=m+1}^{\infty} L^{k-m-1} P_s \frac{dg}{dx}(q^{(-k)}(y))\tilde{q}^{(-k)}(y, y^*, 0).\end{aligned} \tag{7.71}$$

Motivated by these equations, we define a map $J' : X \times \hat{\hat{\mathbf{E}}}_c \times \hat{\hat{\mathbf{E}}}_c \times \mathbf{R} \to X$ to be $J' = (J'_0, J'_{-1}, J'_{-2}, \cdots)$ and

$$J'_{-m}(u, y, y^*, \kappa) = \begin{cases} L^{-m}y^* - \dfrac{1}{\kappa}\displaystyle\sum_{k=1}^{m} L^{k-m-1} P_c(g(q^{(-k)}(y) + \kappa u^{(-k)}) - g(q^{(-k)}(y))) & \\ \qquad + \dfrac{1}{\kappa}\displaystyle\sum_{k=m+1}^{\infty} L^{k-m-1} P_s(g(q^{(-k)}(y) + \kappa u^{(-k)}) - g(q^{(-k)}(y))), & \kappa \neq 0, \\ L^{-m}y^* - \displaystyle\sum_{k=1}^{m} L^{k-m-1} P_c \frac{dg}{dx}(q^{(-k)}(y))u^{(-k)} & \\ \qquad + \displaystyle\sum_{k=m+1}^{\infty} L^{k-m-1} P_s \frac{dg}{dx}(q^{(-k)}(y))u^{(-k)}, & \kappa = 0. \end{cases} \tag{7.72}$$

We can prove that J' is a contraction map on X for each y, y^* and κ by the completely same way as the proofs of Lemma 7.6 and Proposition 7.7. Hence, there uniquely exists $u^{(-m)} = \tilde{q}^{(-m)}(y, y^*, \kappa)$ satisfying Eq.(7.70) and Eq.(7.71). Taking the limit $\kappa \to 0$ in Eq.(7.70) yields

$$\begin{aligned}\lim_{\kappa\to 0} \tilde{q}^{(-m)}(y, y^*, \kappa) \;=\; & L^{-m}y^* - \sum_{k=1}^{m} L^{k-m-1} P_c \frac{dg}{dx}(q^{(-k)}(y)) \lim_{\kappa\to 0} \tilde{q}^{(-k)}(y, y^*, \kappa) \\ & + \sum_{k=m+1}^{\infty} L^{k-m-1} P_s \frac{dg}{dx}(q^{(-k)}(y)) \lim_{\kappa\to 0} \tilde{q}^{(-k)}(y, y^*, \kappa).\end{aligned} \tag{7.73}$$

This implies that $\lim_{\kappa\to 0} \tilde{q}^{(-m)}(y, y^*, \kappa)$ is a solution of Eq.(7.71). By the uniqueness of a solution, we obtain

$$\lim_{\kappa\to 0} \tilde{q}^{(-m)}(y, y^*, \kappa) = \lim_{\kappa\to 0} \frac{1}{\kappa}\left(q^{(-m)}(y + \kappa y^*) - q^{(-m)}(y)\right) = \tilde{q}^{(-m)}(y, y^*, 0). \tag{7.74}$$

From Eq.(7.71), it turns out that $\tilde{q}^{(-m)}(y, y^*, 0)$ is linear in y^*. Thus we denote it as

$$\tilde{q}^{(-m)}(y, y^*, 0) = dq^{(-m)}(y)y^*. \tag{7.75}$$

Then, $dq^{(-m)}(y) : \hat{\hat{\mathbf{E}}}_c \to X^{(-m)}$ defines a linear operator for each $y \in \hat{\hat{\mathbf{E}}}_c$. The remaining task is to show that $dq^{(-m)} : \hat{\hat{\mathbf{E}}}_c \times \hat{\hat{\mathbf{E}}}_c \to X^{(-m)}$ is continuous. This is done in the same way as the proof of part (ii) of the proposition. For $y', y'^* \in \hat{\hat{\mathbf{E}}}_c$, we estimate

$d_\infty(\tilde{q}^{(-m)}(y, y^*, 0), \tilde{q}^{(-m)}(y', y'^*, 0))$. Then, we can show that $\tilde{q}^{(-m)}$ is Lipschitz continuous in y and y^*. The details are left to the reader. This means that $dq^{(-m)}(y)$ gives the derivative of $q^{(-m)}$ at $y \in \hat{\hat{\mathbf{E}}}_c$. ■

Now we define the center manifold W^c of the map φ_τ by

$$W^c = \{q^{(0)}(y) = y + \hat{q}(y) \,|\, y \in \hat{\hat{\mathbf{E}}}_c\}, \tag{7.76}$$

where

$$\hat{q}(y) = \sum_{k=1}^{\infty} L^{k-1} P_s g(q^{(-k)}(y)) \in \hat{\hat{\mathbf{E}}}_c^{\perp}. \tag{7.77}$$

Proposition 7.9. (i) W^c is a dim-$\hat{\hat{\mathbf{E}}}_c$ dimensional C^1 manifold, which is tangent to the space $\hat{\hat{\mathbf{E}}}_c$. In particular, $q^{(0)}(y)$ is expanded as $q^{(0)}(y) = y + O(y^2)$ as $y \to 0$.
(ii) W^c is φ_τ invariant; that is, $\varphi_\tau(W^c) \subset W^c$.
(iii) For any $\xi_0 \in W^c$, there exists a negative semi-orbit $\{u^{(-m)}\}_{m=0}^\infty \subset W^c$ satisfying $u_0 = \xi_0$ and

$$d_\infty(u^{(-m)}, 0) \leq Ce^{bm\tau},$$

where $b > 0$ as above and C is a positive constant.
(iv) if $\delta_1 > 0$ is sufficiently small, the center manifold W^c is included in the strip region $\{z \in \mathbf{R} \times \overline{i(\mathcal{F})} \,|\, d_\infty(P_s z, 0) \leq \delta_2\}$ (see Fig.12).

Proof. (i) Since $q^{(-k)}(0) = 0$ and $q^{(-k)}(y)$ is C^1, $q^{(-k)}(y)$ is expanded as $q^{(-k)}(y) \sim O(y)$. This shows that $\hat{q}(y) \sim O(y^2)$ because $g(z) \sim O(z^2)$ as $z \to 0$ (see Sec.7.4).
(ii) Recall that $\{q^{(-m)}(y)\}_{m=0}^\infty$ is a negative semi-orbit satisfying Eqs.(7.51) and (7.52). Define $q^{(1)}(y) := \varphi_\tau(q^{(0)}(y))$. Obviously $\{q^{(-m+1)}(y)\}_{m=0}^\infty$ is also a negative semi-orbit satisfying (7.51) and (7.52) with some $C > 0$. Then, Lemma 7.5 implies that $\{q^{(-m+1)}(y)\}_{m=0}^\infty$ is a solution of the Lyapunov-Perron equation (7.54). By the uniqueness of a solution, there exists $y' \in \hat{\hat{\mathbf{E}}}_c$ such that $q^{(-m+1)}(y) = q^{(-m)}(y')$ for $m = 0, 1, \cdots$. In particular, we obtain $\varphi_\tau(q^{(0)}(y)) = \varphi_\tau(q^{(-1)}(y')) = q^{(0)}(y')$, which proves $\varphi_\tau(q^{(0)}(y)) \in W^c$.
(iii) This is obvious from the definition: if $\xi_0 = q^{(0)}(y)$, $\{q^{(-m)}(y)\}_{m=0}^\infty$ is a negative semi-orbit included in W^c.
(iv) Prop.7.8 (iii) implies that $P_s q^{(0)}(y) = 0$ if $\|y\|_{E_c} \geq 2\delta_1$. Thus $\sup_{y \in \hat{\hat{\mathbf{E}}}_c} d_\infty(P_s q^{(0)}(y), 0)$ becomes sufficiently small if δ_1 is sufficiently small. ■

If restricted to a small neighborhood of the origin, W^c gives the desired local center manifold for Eq.(6.2).

Proof of Theorem 7.1. If $\delta_1 > 0$ is sufficiently small, W^c is included in the region $\{z \in \mathbf{R} \times \overline{i(\mathcal{F})} \,|\, d_\infty(P_s z, 0) \leq \delta_2\}$, on which $\chi(d_\infty(P_s z, 0)/\delta_2) = 1$. Thus φ_τ-invariant manifold W^c is also invariant under the map $\tilde{\varphi}_\tau$ given by Eq.(7.44), which is a time τ map of the semiflow of the system (7.30). Take $u_0 \in W^c$. By Prop.7.9 (iii), there is a negative semi-orbit $\{u^{(-m)}\}_{m=0}^\infty \subset W^c$ of $\tilde{\varphi}_\tau$ satisfying Eqs.(7.51) and (7.52). Since $\tilde{\varphi}_\tau$ is a semiflow, we have

$$\tilde{\varphi}_\tau \circ \tilde{\varphi}_t(u^{(-m)}) = \tilde{\varphi}_t(\tilde{\varphi}_\tau(u^{(-m)})) = \tilde{\varphi}_t(u^{(-m+1)}), \tag{7.78}$$

for each $t > 0$. This means that $\{\tilde{\varphi}_t(u^{(-m)})\}_{m=0}^{\infty}$ is a negative semi-orbit of $\tilde{\varphi}_\tau$. Since $\tilde{\varphi}_t$ is C^1 with respect to the metric d_∞, there is a positive number $\tilde{C}$ such that

$$d_\infty(\tilde{\varphi}_t(u^{(-m)}), 0) \leq \tilde{C} d_\infty(u^{(-m)}, 0) \leq \tilde{C} C e^{bm\tau}. \tag{7.79}$$

Further, $L^m P_s \tilde{\varphi}_t(u^{(-m)})$ is estimated as

$$L^m P_s \tilde{\varphi}_t(u^{(-m)}) = L^{m+1} P_s u^{(-m)} + L^m P_s \tilde{g}_t(u^{(-m)}).$$

Since $u^{(-m)}$ satisfies (7.51), $L^{m+1} P_s u^{(-m)}$ tends to zero as $m \to \infty$. By the same calculation as Eq.(7.56), we see that the second term $L^m P_s \tilde{g}_t(u^{(-m)})$ also tends to zero as $m \to \infty$. This shows that $\{\tilde{\varphi}_t(u^{(-m)})\}_{m=0}^{\infty}$ satisfies the conditions (7.51) and (7.52). Therefore, it is a solution of the Lyapunov-Perron equation (7.54). By the uniqueness of a solution, there is $y \in \hat{\hat{\mathbf{E}}}_c$ such that $\tilde{\varphi}_t(u^{(0)}) = q^{(0)}(y) \in W^c$, which proves that W^c is $\tilde{\varphi}_t$-invariant.

In Eq.(7.30), since ε is a constant which is independent of t, $W^c(\varepsilon) := W^c \cap (\{\varepsilon\} \times \overline{i(\mathcal{F})})$ is also $\tilde{\varphi}_t$-invariant for each ε.

On the region $\hat{U} = \{z \,|\, ||P_c z||_{E_c} \leq \delta_1,\ |\eta(t)| \leq 1\}$, $\chi(||P_c z||_{E_c}/\delta_1) \cdot \chi(|\eta(t)|) = 1$ and Eq.(7.30) is reduced to the original system (6.2). Thus $W^c(\varepsilon) \cap \hat{U}$ is invariant under the semiflow generated by (6.2), which gives a local center manifold stated in Thm.7.1 with $U = \hat{U} \cap (\{\varepsilon\} \times \overline{i(\mathcal{F})})$ and $W^c_{loc} = W^c(\varepsilon) \cap U$. Parts (I) and (II) in Thm.7.1 immediately follows from Prop.7.9. It remains to show the part (III) of Thm.7.1. This is proved in the same way as Chen, Hale and Tan [9]. In [9], the existence of invariant foliations is proved for dynamical systems on Banach spaces. Though our phase space $\overline{i(\mathcal{F})}$ is not a Banach space, the distance from the origin $d_\infty(z, 0)$ plays the same role as a norm. Thus with the aid of the estimates (7.48) and (7.49), we can prove the existence of invariant foliations by the same way as [9]. The details are left to the reader. ■

7.5 Reduction to the center manifold

Let us derive the dynamics on the center manifold and prove the Kuramoto's conjecture. Recall that for the continuous limit (2.1) of the Kuramoto model, Putting $Z_j(t, \omega) = \int_0^{2\pi} e^{j\sqrt{-1}\theta} \rho_t(\theta, \omega) d\theta$ yields the system of equations (3.2) and (3.3). Since solutions are included in $V_{1,0} \subset F_p$ (Thm.5.10 (iv)), the canonical inclusion is applied to rewrite Eq.(3.2) and (3.3) as equations of the form (6.2) defined on F_p'. The order parameter η is defined as $\eta(t) = (Z_1, P_0) = \langle P_0 \,|\, \overline{Z}_1 \rangle$. For this system, we have proved that when $0 < K < K_c$, the trivial solution (de-synchronous state) is asymptotically stable because of the existence of resonance poles on the left half plane. In particular, $\eta(t) \to 0$ as $t \to \infty$. When $K > K_c$, we have proved that the trivial solution is unstable because of the existence of eigenvalues on the right half plane. Thus a bifurcation from the trivial solution may occur at $K = K_c$. In Sec.7.1 to Sec.7.4, we have proved that there exists a smooth local center manifold near the origin in $\prod_{k=1}^{\infty} F_p'$ if K is sufficiently close to K_c. Our purpose is to obtain a differential equation describing the dynamics on the center manifold to reveal a bifurcation structure of the Kuramoto model.

To derive the dynamics on the center manifold, we impose the following hypothesis as is assumed in the Kuramoto's conjecture.

(A6) The probability density function $g(\omega)$ is an even and unimodal function such that $g''(0) \neq 0$.

Note that $g''(0) \leq 0$ when $g(\omega)$ is unimodal. Thus (A6) implies $g''(0) < 0$. Under this assumption, there exists only one resonance pole $\lambda_0 = 0$ on the imaginary axis when $K = K_c$. Indeed, the imaginary parts of resonance poles on the imaginary axis are given as roots of the second equation of (3.17), which is calculated as

$$0 = \lim_{x \to +0} \int_{\mathbf{R}} \frac{\omega - y}{x^2 + (\omega - y)^2} g(\omega) d\omega = \lim_{x \to +0} \int_0^\infty \frac{\omega}{x^2 + \omega^2} \left(g(y + \omega) - g(y - \omega) \right) d\omega.$$

Since g is even, $y = 0$ is a root of this equation. Since g is unimodal, $g(y+\omega) - g(y-\omega) > 0$ when $y < 0, \omega > 0$ and $g(y + \omega) - g(y - \omega) < 0$ when $y > 0, w > 0$. Thus $y = 0$ is a unique root. This proves that $\lambda_0 = 0$ is the unique resonance pole on the imaginary axis when $K = K_c$. Further, the resonance pole is a pole of first order. Indeed,

$$\lim_{x \to +0} \frac{d}{dy}\Big|_{y=0} \int_0^\infty \frac{\omega}{x^2 + \omega^2} \left(g(y + \omega) - g(y - \omega) \right) d\omega = \lim_{x \to +0} \int_0^\infty \frac{2\omega}{x^2 + \omega^2} \frac{dg}{d\omega}(\omega) d\omega.$$

Since g is unimodal, we have $dg/d\omega \leq 0$, and the above value is not zero. This means that $y = 0$ is a single root. Thus the center subspace $\mathbf{E}_c$ is of one dimensional. Let $|\,\mu_0\rangle$ be the generalized eigenfunction associated with $\lambda_0 = 0$. By the definition, $|\,\mu_0\rangle$ is given by

$$\langle \phi\,|\,\mu_0 \rangle = \lim_{x \to +0} \int_{\mathbf{R}} \frac{1}{x - \sqrt{-1}\omega} \phi(\omega) g(\omega) d\omega. \tag{7.80}$$

This is also written as

$$|\,\mu_0\rangle = \lim_{x \to +0} \left| \overline{\frac{1}{x - \sqrt{-1}\omega}} \right\rangle = \lim_{x \to +0} i\left(\frac{1}{x - \sqrt{-1}\omega}\right), \tag{7.81}$$

where the limit is taken with respect to the weak dual topology on F'_p. The main theorem in this section, which confirms the Kuramoto's conjecture, is stated as follows:

Theorem 7.10. For the continuous model (2.1) of the Kuramoto model, suppose that the function $g(\omega)$ satisfies assumptions (A1) to (A3),(A5) and (A6). Then, there exist positive constants ε_0 and δ such that if $K_c < K < K_c + \varepsilon_0$ and if the initial condition $h(\theta)$ satisfies

$$\left| \int_0^{2\pi} e^{\sqrt{-1} j \theta} h(\theta) d\theta \right| < \delta \tag{7.82}$$

for $j = 1, 2, \cdots$, then the order parameter $\eta(t)$ tends to the constant expressed as

$$r(t) = |\eta(t)| = \sqrt{\frac{-16}{\pi K_c^4 g''(0)}} \sqrt{K - K_c} + O(K - K_c), \tag{7.83}$$

as $t \to \infty$. In particular, the bifurcation diagram of the order parameter is given as Fig.2 (a).



Fig. 13: Eq.(7.83) holds even if the shape of $g(\omega)$ is slightly deformed from an even and unimodal function. If $g''(0) > 0$, the unstable synchronous solution appears backward.

Remark. A few remarks are in order. When $g(\omega)$ is a rational function with the assumption (A6), the theorem also holds without any modifications, though the existence of the center manifold is trivial because $i(H_+)$ is of finite dimensional. For both cases, we do not need the assumption (A4) because (A6) implies that the resonance pole on the imaginary axis is a pole of first order. The assumption (A6) is also not essential. The theorem is true as long as only one resonance pole gets across the imaginary axis at the origin. For example, let us slightly deform an even and unimodal $g(\omega)$ so that it sags in the center as is shown in Fig.13. If the sag is sufficiently small, only one resonance pole gets across the imaginary axis along the real axis, and Eq.(7.83) holds. In this case, since $g''(0) > 0$, $|\eta(t)|$ in Eq.(7.83) is positive when $K < K_c$. This solution is unstable as is verified from the dynamics on the center manifold given as (7.99). This means that a subcritical bifurcation occurs and the bifurcation diagram shown in Fig.2 (b) is (partially) obtained. In general, the more complex a shape of $g(\omega)$ becomes, the more complex bifurcations occur. The result (7.83) coincides with formal discussions and numerical simulations obtained by many physicists, see Daido [15] and Strogatz [42], for example.

Proof of Thm.7.10. Suppose that an initial condition $h(\theta)$ satisfies Eq.(7.82). Then, we have

$$\begin{aligned}
||\, |\overline{Z_j(0,\cdot)}\rangle\, ||^*_{\beta,n} &= \sup_{||\phi||_{\beta,n}=1} \Big| \int_{\mathbf{R}} Z_j(0,\omega)\phi(\omega)g(\omega)d\omega \Big| \\
&= \sup_{||\phi||_{\beta,n}=1} \Big| \int_{\mathbf{R}} \int_0^{2\pi} e^{\sqrt{-1}j\theta} h(\theta)\phi(\omega)g(\omega)d\theta d\omega \Big| \\
&= \Big| \int_0^{2\pi} e^{\sqrt{-1}j\theta} h(\theta)d\theta \Big| \cdot \sup_{||\phi||_{\beta,n}=1} \Big| \int_{\mathbf{R}} \phi(\omega)g(\omega)d\omega \Big| \\
&\leq \delta \cdot ||P_0||^*_{\beta,n},
\end{aligned}$$

for every j, β and n. Thus we can take δ sufficiently small so that the initial condition $(|\overline{Z_1(0,\cdot)}\rangle, |\overline{Z_2(0,\cdot)}\rangle, \cdots)$ for Eq.(6.2) is included in the neighborhood U (with respect to the metric d_∞) of the origin given in Thm.7.1. Then, the center manifold theorem is applicable. Let us derive the dynamics on the center manifold.

Since we are interested in a bifurcation at $K = K_c$, put $\varepsilon = K - K_c$ and divide the operator T_1 as

$$T_1\phi(\omega) = T_{10}\phi(\omega) + \frac{\varepsilon}{2}\langle P_0 \,|\, \overline{\phi}\rangle P_0(\omega), \tag{7.84}$$

where

$$T_{10}\phi(\omega) = \sqrt{-1}\omega\phi(\omega) + \frac{K_c}{2}\langle P_0 \,|\, \overline{\phi}\rangle P_0(\omega). \tag{7.85}$$

Then, the operator T_{10} has a resonance pole at the origin and all other resonance poles are on the left half plane. Eq.(6.2) is rewritten as

$$\frac{d}{dt}|\,\overline{Z}_1\rangle = T_{10}^{\times}|\,\overline{Z}_1\rangle + \frac{\varepsilon}{2}\overline{\langle P_0 \,|\, \overline{Z}_1\rangle}|\,P_0\rangle - \frac{K}{2}\langle P_0 \,|\, \overline{Z}_1\rangle|\,\overline{Z}_2\rangle. \tag{7.86}$$

To obtain the dynamics on the center manifold, by using the spectral decomposition (5.65), we put

$$|\,\overline{Z}_1\rangle = \frac{K_c}{2}\alpha(t)|\,\mu_0\rangle + |\,Y_1\rangle, \tag{7.87}$$

where $|\,\mu_0\rangle$ is defined by Eq.(7.81), $|\,Y_1\rangle$ is included in the complement $\mathbf{E}_c^{\perp}$ of $\mathbf{E}_c$, and where

$$\alpha(t) = \frac{1}{\overline{D}_0}\overline{\langle Z_1 \,|\, \mu_0\rangle}. \tag{7.88}$$

We will derive the dynamics of α. Since $\langle P_0 \,|\, \mu_n\rangle = 2/K_c$ for $n = 0, 1, \cdots$ by the definitions of $|\,\mu_n\rangle$ and the resonance poles, we obtain

$$\langle P_0 \,|\, \overline{Z}_1\rangle = \overline{\alpha(t)} + \langle P_0 \,|\, Y_1\rangle, \tag{7.89}$$

and $|\,P_0\rangle$ is decomposed as

$$|\,P_0\rangle = \frac{1}{\overline{D}_0}|\,\mu_0\rangle + |\,Y_0\rangle, \tag{7.90}$$

where $|\,Y_0\rangle \in \mathbf{E}_c^{\perp}$. By Thm.7.1 (I), on the local center manifold, we can suppose that

$$\langle \phi \,|\, Y_1\rangle,\ \langle \phi \,|\, \overline{Z}_j\rangle \sim O(\alpha^2, \alpha\varepsilon, \varepsilon^2), \tag{7.91}$$

for $j = 2, 3, \cdots$ and for every $\phi \in F_p$. Let us calculate the expression of the center manifold. Substituting Eqs.(7.89),(7.87) into Eq.(6.2) for $j = 2$ yields

$$\frac{d}{dt}|\,\overline{Z}_2\rangle = T_2^{\times}|\,\overline{Z}_2\rangle + K\Big((\alpha + \overline{\langle P_0 \,|\, Y_1\rangle})\Big(\frac{K_c}{2}\alpha|\,\mu_0\rangle + |\,Y_1\rangle\Big) - (\overline{\alpha} + \langle P_0 \,|\, Y_1\rangle)|\,\overline{Z}_3\rangle\Big). \tag{7.92}$$

We suppose that $d\alpha/dt \sim O(\alpha^2, \alpha\varepsilon, \varepsilon^2)$, which will be justified later. Then, the above equation yields

$$T_2^{\times}|\,\overline{Z}_2\rangle = -\frac{KK_c}{2}\alpha^2|\,\mu_0\rangle + O(\alpha^3, \alpha^2\varepsilon, \alpha\varepsilon^2, \varepsilon^3). \tag{7.93}$$

Lemma 7.11. Define the operator $(T_2^{\times})^{-1} : i(F_p) \to F_p'$ to be

$$\langle \phi \,|\, (T_2^{\times})^{-1}\overline{\psi}\rangle = -\frac{1}{2}\lim_{x\to+0}\int_{\mathbf{R}}\frac{1}{x - \sqrt{-1}\omega}\phi(\omega)\psi(\omega)g(\omega)d\omega. \tag{7.94}$$

Then,

$$(T_2^\times)(T_2^\times)^{-1}|\,\overline{\psi}\rangle = (T_2^\times)^{-1}(T_2^\times)|\,\overline{\psi}\rangle = |\,\overline{\psi}\rangle \tag{7.95}$$

for any $|\,\overline{\psi}\rangle \in i(F_p)$, and it is continuous on $i(V)$.

Proof. The straightforward calculation shows that

$$\begin{aligned}
\langle \phi\,|\,(T_2^\times)(T_2^\times)^{-1}\overline{\psi}\rangle &= \langle T_2\phi\,|\,(T_2^\times)^{-1}\overline{\psi}\rangle \\
&= -\frac{1}{2}\lim_{x\to+0}\int_{\mathbf{R}}\frac{2\sqrt{-1}\omega}{x-\sqrt{-1}\omega}\phi(\omega)\psi(\omega)g(\omega)d\omega \\
&= \int_{\mathbf{R}}\phi(\omega)\psi(\omega)g(\omega)d\omega - \lim_{x\to+0}\int_{\mathbf{R}}\frac{x}{x-\sqrt{-1}\omega}\phi(\omega)\psi(\omega)g(\omega)d\omega.
\end{aligned}$$

Since the limit

$$\lim_{x\to+0}\int_{\mathbf{R}}\frac{1}{x-\sqrt{-1}\omega}\phi(\omega)\psi(\omega)g(\omega)d\omega = \langle \phi\cdot\psi\,|\,\mu_0\rangle$$

exists, the second term in the right hand side above is zero. Thus we obtain

$$\langle \phi\,|\,(T_2^\times)(T_2^\times)^{-1}\overline{\psi}\rangle = \langle \phi\,|\,\overline{\psi}\rangle.$$

In the same way,

$$\begin{aligned}
\langle \phi\,|\,(T_2^\times)^{-1}(T_2^\times)\overline{\psi}\rangle &= \langle \phi\,|\,(T_2^\times)^{-1}(-2\sqrt{-1}\omega)\overline{\psi}\rangle \\
&= -\frac{1}{2}\lim_{x\to+0}\int_{\mathbf{R}}\frac{2\sqrt{-1}\omega}{x-\sqrt{-1}\omega}\phi(\omega)\psi(\omega)g(\omega)d\omega \\
&= \langle \phi\,|\,\overline{\psi}\rangle.
\end{aligned}$$

Note that the right hand side of (7.94) is also written as $-\langle \phi\cdot\psi\,|\,\mu_0\rangle/2$, which coincides with the coefficient of $\Pi_0|\,\overline{\phi\cdot\psi}\rangle$, up to the constant factor. Thus the proof of continuity of $(T_2^\times)^{-1}$ on $i(V)$ is the same as that of the continuity of Π_0 on $i(V)$. ■

Since $(T_2^\times)^{-1}$ is continuous on $i(V)$, its domain is continuously extended to the closure $\overline{i(V)}$. Since $|\,\mu_0\rangle \in \overline{i(V)}$ (see Prop.5.20) and it is given as Eq.(7.81), $(T_2^\times)^{-1}|\,\mu_0\rangle$ is calculated as

$$\langle \phi\,|\,(T_2^\times)^{-1}\mu_0\rangle = \lim_{x\to+0}\left\langle \phi\,\middle|\,(T_2^\times)^{-1}\overline{\frac{1}{x-\sqrt{-1}\omega}}\right\rangle = -\frac{1}{2}\lim_{x\to+0}\int_{\mathbf{R}}\frac{1}{(x-\sqrt{-1}\omega)^2}\phi(\omega)g(\omega)d\omega. \tag{7.96}$$

Then, Eq.(7.93) provides

$$\langle \phi\,|\,\overline{Z}_2\rangle = \frac{KK_c}{4}\overline{\alpha}^2\lim_{x\to+0}\int_{\mathbf{R}}\frac{1}{(x-\sqrt{-1}\omega)^2}\phi(\omega)g(\omega)d\omega + O(\alpha^3,\alpha^2\varepsilon,\alpha\varepsilon^2,\varepsilon^3), \tag{7.97}$$

which gives the expression of the center manifold to the $|\,\overline{Z}_2\rangle$ direction. The projection of it to the center subspace is given as

$$\begin{aligned}
\Pi_0|\,\overline{Z}_2\rangle &= \frac{K_c}{2\overline{D}_0}\langle\overline{Z_2\,|\,\mu_0\rangle}\cdot|\,\mu_0\rangle \\
&= \frac{K_c}{2\overline{D}_0}\lim_{x\to+0}\overline{\Big\langle Z_2\,\Big|\,\overline{\frac{1}{x-\sqrt{-1}\omega}}\Big\rangle}\cdot|\,\mu_0\rangle = \frac{K_c}{2\overline{D}_0}\lim_{x\to+0}\Big\langle\overline{\frac{1}{x-\sqrt{-1}\omega}\,\Big|\,\overline{Z}_2}\Big\rangle\cdot|\,\mu_0\rangle,
\end{aligned}$$

where

$$\begin{aligned}
\lim_{x\to+0}\Big\langle\frac{1}{x-\sqrt{-1}\omega}\,\Big|\,\overline{Z}_2\Big\rangle &= \frac{KK_c}{4}\overline{\alpha}^2\lim_{x\to+0}\int_{\mathbf{R}}\frac{1}{(x-\sqrt{-1}\omega)^3}g(\omega)d\omega + O(\alpha^3,\alpha^2\varepsilon,\alpha\varepsilon^2,\varepsilon^3) \\
&= -\frac{KK_c}{8}\overline{\alpha}^2\lim_{x\to+0}\int_{\mathbf{R}}\frac{1}{x-\sqrt{-1}\omega}g''(\omega)d\omega + O(\alpha^3,\alpha^2\varepsilon,\alpha\varepsilon^2,\varepsilon^3) \\
&= -\frac{KK_c}{8}\overline{\alpha}^2\cdot\pi g''(0) + O(\alpha^3,\alpha^2\varepsilon,\alpha\varepsilon^2,\varepsilon^3).
\end{aligned}$$

Thus we obtain

$$\Pi_0|\,\overline{Z}_2\rangle = -\frac{KK_c^2}{16\overline{D}_0}\alpha^2\cdot\pi g''(0)\cdot|\,\mu_0\rangle + O(\alpha^3,\alpha^2\varepsilon,\alpha\varepsilon^2,\varepsilon^3). \tag{7.98}$$

Finally, the projection of Eq.(7.86) to the center subspace is given by

$$\frac{d}{dt}\Pi_0|\,\overline{Z}_1\rangle = T_{10}^{\times}\Pi_0|\,\overline{Z}_1\rangle + \frac{\varepsilon}{2}\langle\overline{P_0\,|\,\overline{Z}_1\rangle}\Pi_0|\,P_0\rangle - \frac{K}{2}\langle P_0\,|\,\overline{Z}_1\rangle\Pi_0|\,\overline{Z}_2\rangle.$$

By using Eqs.(7.87),(7.89),(7.90) and (7.98), we obtain

$$\begin{aligned}
\frac{d}{dt}\frac{K_c}{2}\alpha|\,\mu_0\rangle &= \frac{K_c}{2}\alpha T_{10}^{\times}|\,\mu_0\rangle + \frac{\varepsilon}{2}\Big(\alpha + \langle\overline{P_0\,|\,Y_1\rangle}\Big)\frac{1}{\overline{D}_0}|\,\mu_0\rangle \\
&\quad -\frac{K}{2}(\overline{\alpha} + \langle P_0\,|\,Y_1\rangle)\cdot\left(-\frac{\pi g''(0)KK_c^2}{16\overline{D}_0}\alpha^2|\,\mu_0\rangle + O(\alpha^3,\alpha^2\varepsilon,\alpha\varepsilon^2,\varepsilon^3)\right), \\
&= \frac{\varepsilon}{2\overline{D}_0}\alpha|\,\mu_0\rangle + \frac{\pi g''(0)K_c^4}{32\overline{D}_0}\alpha|\alpha|^2|\,\mu_0\rangle + O(\varepsilon\alpha^2,\varepsilon^2\alpha,\varepsilon^3,\alpha^4),
\end{aligned}$$

which yields the dynamics on the center manifold as

$$\frac{d}{dt}\alpha = \frac{\alpha}{\overline{D}_0K_c}\left(\varepsilon + \frac{\pi g''(0)K_c^4}{16}|\alpha|^2\right) + O(\varepsilon\alpha^2,\varepsilon^2\alpha,\varepsilon^3,\alpha^4). \tag{7.99}$$

Since $g''(0) < 0$ because of the assumption (A6), this equation has a fixed point expressed as Eq.(7.83) when $\varepsilon = K - K_c > 0$. Note that the order parameter $\eta(t) = (Z_1, P_0)$ is rewritten as

$$\eta(t) = (Z_1,P_0) = (P_0,\overline{Z}_1) = \langle P_0\,|\,\overline{Z}_1\rangle = \frac{K_c}{2}\overline{\alpha}\langle P_0\,|\,\mu_0\rangle + \langle P_0\,|\,Y_1\rangle = \overline{\alpha} + O(\alpha^2,\alpha\varepsilon,\varepsilon^2). \tag{7.100}$$

Thus the dynamics of the order parameter is also given by Eq.(7.99). To prove that the fixed point (7.83) is asymptotically stable, it is sufficient to show the following.

Lemma 7.12. $D_0 > 0$.

Proof Put

$$f(\lambda) = 1 - \frac{K_c}{2}\int_{\mathbf{R}} \frac{1}{\lambda - \sqrt{-1}\omega} g(\omega) d\omega - \pi K_c g(-\sqrt{-1}\lambda).$$

By the definition of D_0,

$$\begin{aligned} D_0 = f'(0) &= \lim_{\lambda\to 0}\frac{K_c}{2}\int_{\mathbf{R}} \frac{1}{(\lambda - \sqrt{-1}\omega)^2} g(\omega) d\omega + \sqrt{-1}\pi K_c g'(0) \\ &= \frac{\sqrt{-1}K_c}{2}\lim_{\lambda\to 0}\int_{\mathbf{R}} \frac{1}{\lambda - \sqrt{-1}\omega} g'(\omega) d\omega + \sqrt{-1}\pi K_c g'(0). \end{aligned}$$

Since $g(\omega)$ is even,

$$D_0 = -\frac{K_c}{2}\lim_{x\to 0}\int_{\mathbf{R}} \frac{\omega}{x^2+\omega^2} g'(\omega) d\omega = -K_c \lim_{x\to 0}\int_0^\infty \frac{\omega}{x^2+\omega^2} g'(\omega) d\omega.$$

Since $g(\omega)$ is unimodal, $g'(\omega) \leq 0$ when $\omega > 0$, which proves that $D_0 > 0$. ■

Since $D_0 > 0, K_c > 0, g''(0) < 0$, the fixed point $\alpha = 0$ (de-synchronous state) is unstable and the fixed point Eq.(7.83) (synchronous state) is asymptotically stable when $\varepsilon = K - K_c > 0$. This completes the proof. ■

A Resonance poles of the Gaussian distribution

In this appendix, we give a few simple properties of resonance poles for the Gaussian distribution $g(\omega) = e^{-\omega^2/2}/\sqrt{2\pi}$. In particular, we prove that it satisfies the assumption (A4) (the other assumptions (A1) to (A3), (A5) and (A6) are trivially satisfied).

The transition point is given by $K_c = 2\sqrt{2/\pi}$. The equation (4.11) for obtaining the resonance poles is reduced to

$$\int_{\mathbf{R}} \frac{1}{\lambda - \sqrt{-1}\omega} g(\omega) d\omega + 2\pi g(-\sqrt{-1}\lambda) = e^{\lambda^2/2}\left(\sqrt{\frac{\pi}{2}} - \int_0^\lambda e^{-x^2/2}dx\right) = \frac{2}{K}. \qquad \text{(A.1)}$$

Let $\lambda_0, \lambda_1, \cdots$ be roots of this equation with $0 \geq \mathrm{Re}(\lambda_0) \geq \mathrm{Re}(\lambda_1) \geq \cdots$.

Proposition A.1. If λ_n is a resonance pole, so is its complex conjugate $\overline{\lambda}_n$.

Proof. This is always true when $g(\omega)$ is an even function, and immediately follows from Eq.(4.11). ■

Proposition A.2. There exist infinitely many resonance poles. As $n \to \infty$, $\mathrm{Re}(\lambda_n) \to -\infty$ and they approach to the rays $\arg(z) = 3\pi/4, 5\pi/4$.

Proof. If $|\lambda|$ is sufficiently large, Eq.(A.1) is estimated as

$$e^{\lambda^2/2} \sim \frac{2}{\sqrt{2\pi}K} + O(1/\lambda), \tag{A.2}$$

which gives the desired results. ■

Proposition A.3. When $0 < K < K_c$, there exists only one resonance pole on the real axis.

Proof. Put

$$f(\lambda) = e^{\lambda^2/2}\left(\sqrt{\frac{\pi}{2}} - \int_0^\lambda e^{-x^2/2}dx\right) - \frac{2}{K}. \tag{A.3}$$

If λ is negative,

$$f'(\lambda) = \lambda e^{\lambda^2/2}\left(\sqrt{\frac{\pi}{2}} - \int_0^\lambda e^{-x^2/2}dx\right) - 1 < 0. \tag{A.4}$$

Thus $f(\lambda)$ is monotonically decreasing on the negative real axis. If $0 < K < K_c$, then $f(0) = \sqrt{\pi/2} - 2/K < 0$, and if $\lambda < 0$ is sufficiently small, $f(\lambda) > 0$. Thus the intermediate-value theorem proves the proposition. ■

Proposition A.4. All roots of Eq.(A.1) are simple roots. Thus the Gaussian distribution satisfies the assumption (A4).

Proof. Put

$$f(\lambda) = \int_{\mathbf{R}} \frac{1}{\lambda - \sqrt{-1}\omega} g(\omega) + 2\pi g(-\sqrt{-1}\lambda) - \frac{2}{K} \tag{A.5}$$

(this is the same function as Eq.(A.3)). The derivative is calculated as

$$\begin{aligned} f'(\lambda) &= -\int_{\mathbf{R}} \frac{1}{(\lambda - \sqrt{-1}\omega)^2} g(\omega)d\omega - 2\pi\sqrt{-1}g'(-\sqrt{-1}\lambda) \\ &= -\int_{\mathbf{R}} \frac{\sqrt{-1}}{\lambda - \sqrt{-1}\omega} g'(\omega)d\omega - 2\pi\sqrt{-1}g'(-\sqrt{-1}\lambda). \end{aligned}$$

Since the Gaussian distribution satisfies $g'(\omega) = -\omega g(\omega)$, we obtain

$$\begin{aligned} f'(\lambda) &= \int_{\mathbf{R}} \frac{\sqrt{-1}\omega}{\lambda - \sqrt{-1}\omega} g(\omega)d\omega + 2\pi\lambda g(-\sqrt{-1}\lambda) \\ &= \lambda \int_{\mathbf{R}} \frac{1}{\lambda - \sqrt{-1}\omega} g(\omega)d\omega + 2\pi\lambda g(-\sqrt{-1}\lambda) - 1. \end{aligned}$$

If λ_n is a resonance pole, it satisfies $f(\lambda_n) = 0$. By using this, the above is rewritten as

$$f'(\lambda_n) = \lambda_n \frac{2}{K} - 1. \tag{A.6}$$

Since $\mathrm{Re}(\lambda_n) \leq 0$, $f'(\lambda_n) \neq 0$. This proves the proposition. ■

B Proof of the existence of solutions of Eq.(7.9)

In this Appendix B, for Eq.(7.9) with $\hat{\chi}$ given by (7.38), we prove the existence of solutions and smoothness of them with respect to initial conditions to complete the proof of Prop.7.3. Recall that Eq.(7.9) is equivalent to the integro-ODE (7.12) and (7.13). Put

$$\left\{\begin{array}{l} f_1(x,\eta,\alpha) = \dfrac{K}{2\sqrt{-1}}(\eta e^{-\sqrt{-1}x} - \overline{\eta} e^{\sqrt{-1}x}) \cdot \chi\Big(\dfrac{||P_c z||}{\delta_1}\Big) \cdot \chi(|\eta(t)|), \\ f_2(x,\eta,\alpha) = \dfrac{K}{2}(\eta e^{-\sqrt{-1}x} + \overline{\eta} e^{\sqrt{-1}x}) \cdot \chi\Big(\dfrac{||P_c z||}{\delta_1}\Big) \cdot \chi(|\eta(t)|), \\ f_3(x,\eta,\alpha) = \dfrac{1}{2}(\eta e^{-\sqrt{-1}x} + \overline{\eta} e^{\sqrt{-1}x}) \cdot \Big(K_c + \varepsilon\chi\Big(\dfrac{||P_c z||}{\delta_1}\Big) \cdot \chi(|\eta(t)|)\Big), \end{array}\right.$$

where $\alpha = (\alpha_0, \alpha_1, \cdots, \alpha_n)$ is defined through Eq.(7.41) and (7.42). Then, (7.12) and (7.13) are rewritten as a system of integral equations

$$\left\{\begin{array}{l} x(t,0;\theta,\omega) = \omega t + \theta + \displaystyle\int_0^t f_1(x(s,0;\theta,\omega),\eta(s),\alpha(s))ds, \\ \hat{\rho}_t(t,\omega) = \hat{h}(x(0,t;\theta,\omega),\omega)\exp\Big[\displaystyle\int_0^t f_2(x(s,t;\theta,\omega),\eta(s),\alpha(s))ds\Big] \\ \quad +\dfrac{1}{2\pi}\displaystyle\int_0^t \exp\Big[\int_s^t f_2(x(\tau,t;\theta,\omega),\eta(\tau),\alpha(\tau))d\tau\Big] f_3(x(s,t;\theta,\omega),\eta(s),\alpha(s))ds, \\ \eta(t) = (\hat{Z}_1, P_0) = \displaystyle\int_{\mathbf{R}} \hat{Z}_1(t,\omega)g(\omega)d\omega, \\ \alpha_j(t) = \dfrac{K}{2\overline{D}_j}\overline{\langle \hat{Z}_1 \,|\, \mu_j\rangle}. \end{array}\right. \tag{B.1}$$

In particular, Eq.(7.17) is rewritten as

$$\hat{Z}_k(t,\omega) = \int_0^{2\pi} e^{\sqrt{-1}k\theta}\hat{\rho}_t(\theta,\omega)d\theta = \int_0^{2\pi} e^{\sqrt{-1}kx(t,0;\theta,\omega)}\hat{h}(\theta,\omega)d\theta \tag{B.2}$$

$$+\frac{1}{2\pi}\int_0^{2\pi} e^{\sqrt{-1}kx(t,0;\theta,\omega)}\int_0^t \exp\Big[\int_s^0 f_2(x(\tau,0;\theta,\omega),\eta(\tau),\alpha(\tau))d\tau\Big] f_3(x(s,0;\theta,\omega),\eta(s),\alpha(s))dsd\theta.$$

Since there is a one to one correspondence between a measure on S^1 and its Fourier coefficients (see Shohat and Tamarkin [40]), $\hat{\rho}_t$ is uniquely determined by $\{\hat{Z}_k\}_{k=0}^\infty$.

Proposition B.1. For any signed measure $\hat{h}(\theta,\omega)$ on S^1 parameterized by $\omega \in \mathbf{R}$ satisfying Eq.(7.14), the system of integral equations (B.1) has a unique solution for any $t \geq 0$.

Proof. At first, we construct a local solution by the Picard's iteration method. Once we obtain x, η and α, then $\hat{Z}_1, \hat{Z}_2, \cdots$ are obtained by substituting them into Eq.(B.2). Thus we consider to construct x, η and α by the iteration. Define sequences $\{x_m(t,0;\theta,\omega)\}_{m=0}^\infty$, $\{Z_{m,1}(t,\omega)\}_{m=0}^\infty$, $\{\eta_m(t)\}_{m=0}^\infty$ and $\{\alpha_m(t)\}_{m=0}^\infty$ to be

$$x_0(t,0;\theta,\omega) = \omega t + \theta, \quad Z_{0,1}(t,\omega) = \int_0^{2\pi} e^{\sqrt{-1}\theta}\hat{h}(\theta,\omega)d\theta, \tag{B.3}$$

and

$$\left\{\begin{aligned} & x_{m+1}(t,0;\theta,\omega) = x_0 + \int_0^t f_1(x_m(s,0;\theta,\omega),\eta_m(s),\alpha_m(s))ds, \\ & Z_{m+1,1}(t,\omega) = \int_0^{2\pi} e^{\sqrt{-1}x_{m+1}(t,0;\theta,\omega)}\hat{h}(\theta,\omega)d\theta + \frac{1}{2\pi}\int_0^{2\pi} e^{\sqrt{-1}x_{m+1}(t,0;\theta,\omega)}\times \\ & \qquad \int_0^t \exp\left[\int_s^0 f_2(x_m(\tau,0;\theta,\omega),\eta_m(\tau),\alpha_m(\tau))d\tau\right] f_3(x_m(s,0;\theta,\omega),\eta_m(s),\alpha_m(s))dsd\theta, \\ & \eta_m(t) = \int_{\mathbf{R}} Z_{m,1}(t,\omega)g(\omega)d\omega, \\ & \alpha_{m,j}(t) = \frac{K}{2\overline{D}_j}\langle\overline{Z_{m,1}\,|\,\mu_j}\rangle, \quad \alpha_m(t) = (\alpha_{m,0}(t),\cdots,\alpha_{m,n}(t)). \end{aligned}\right. \tag{B.4}$$

In what follows, we suppose that $\theta,\omega\in\mathbf{R}$. In this case, we can verify by induction that $x_m(t,0;\theta,\omega)\in\mathbf{R}$ for every m. We will show later that $x(t,0;\theta,\omega)$ has an analytic continuation in θ and ω.

Take a positive number M_1 and put

$$R = \{(x,\eta,\alpha)\,|\,|\eta-\eta_0|\le M_1\}\subset\mathbf{R}\times\mathbf{C}\times\mathbf{C}^{n+1}. \tag{B.5}$$

When $(x,\eta,\alpha)\in R$, there exists a positive constant M_2 such that

$$|f_i(x,\eta,\alpha)|\le K|\eta|\le M_2, \tag{B.6}$$

for $i=1,2,3$. If $(x_{m-1},\eta_{m-1},\alpha_{m-1})\in R$, $|\eta_m-\eta_0|$ is estimated as

$$\begin{aligned} |\eta_m-\eta_0| \;\le\; & \int_{\mathbf{R}}\int_0^{2\pi} |e^{\sqrt{-1}x_m(t,0;\theta,\omega)} - e^{\sqrt{-1}\theta}|\cdot|\hat{h}(\theta,\omega)|g(\omega)d\omega d\theta \\ & + \frac{1}{2\pi}\int_{\mathbf{R}}\int_0^{2\pi}\int_0^t \exp\left[\int_0^s |f_2(x_{m-1},\eta_{m-1},\alpha_{m-1})|d\tau\right]|f_3(x_{m-1},\eta_{m-1},\alpha_{m-1})|g(\omega)dsd\theta d\omega \\ \le\; & \int_{\mathbf{R}}\int_0^{2\pi} |x_m(t,0;\theta,\omega)-\theta|\cdot|\hat{h}(\theta,\omega)|g(\omega)d\omega d\theta + \frac{1}{2\pi}\int_{\mathbf{R}}\int_0^{2\pi}\int_0^t e^{M_2 s}M_2 g(\omega)dsd\theta d\omega, \end{aligned}$$

and $|x_m-\theta|$ is estimated as

$$|x_m-\theta|\le|\omega|t+\int_0^t|f_1(x_{m-1},\eta_{m-1},\alpha_{m-1})|ds\le(|\omega|+M_2)t.$$

Thus we obtain

$$|\eta_m-\eta_0|\le\int_{\mathbf{R}}\int_0^{2\pi}(|\omega|+M_2)\cdot|\hat{h}(\theta,\omega)|g(\omega)d\omega d\theta\cdot t+(e^{M_2 t}-1).$$

By the assumption (A2) for $g(\omega)$ and the assumption (F2) for $\hat{h}$, the integral above exists. Hence, there exists $T_1>0$ such that when $0\le t\le T_1$, $|\eta_m-\eta_0|\le M_1$ holds.

Now the induction shows that $(x_m, \eta_m, \alpha_m) \in R$ and thus Eq.(B.6) is valid for every (x_m, η_m, α_m), $m = 0, 1, \cdots$ on the interval $0 \leq t \leq T_1$.

When $0 \leq t \leq T_1$, we obtain

$$\begin{aligned}
|x_1 - x_0| &\leq \int_0^t |f_1(x_0, \eta_0, \alpha_0)| ds \leq M_2 t, \\
|Z_{1,1} - Z_{0,1}| &\leq \int_0^{2\pi} |e^{\sqrt{-1}x_1} - e^{\sqrt{-1}\theta}| \cdot |\hat{h}(\theta, \omega)| d\theta \\
&\quad + \frac{1}{2\pi} \int_0^{2\pi} \int_0^t \exp\left[\int_0^s |f_2(x_0, \eta_0, \alpha_0)| d\tau\right] \cdot |f_3(x_0, \eta_0, \alpha_0)| ds d\theta \\
&\leq (|\omega| + M_2)t \int_0^{2\pi} |\hat{h}(\theta, \omega)| d\theta + \frac{1}{2\pi} \int_0^{2\pi} \int_0^t e^{M_2 s} M_2 ds d\theta \\
&= C_1(|\omega| + M_2)t + (e^{M_2 t} - 1),
\end{aligned}$$

where we put $C_1 = \int_0^{2\pi} |\hat{h}(\theta, \omega)| d\theta$. There exists a constant $M_3 > 0$ such that $e^{M_2 t} - 1 \leq M_3 t$ on $0 \leq t \leq T_1$. Then,

$$\begin{aligned}
|Z_{1,1} - Z_{0,1}| &\leq (C_1|\omega| + C_1 M_2 + M_3)t, \\
|\eta_1 - \eta_0| &\leq \int_{\mathbf{R}} (C_1|\omega| + C_1 M_2 + M_3)t\, g(\omega) d\omega := M_4 t.
\end{aligned}$$

Let $\lambda_j = \sqrt{-1} y_j$ $(j = 0, \cdots, n)$ be resonance poles on the imaginary axis. By the definition, $|\alpha_{1,j} - \alpha_{0,j}|$ is calculated as

$$\begin{aligned}
|\alpha_{1,j} - \alpha_{0,j}| &\leq \frac{K}{2|D_j|} |\langle Z_{1,1} - Z_{0,1} \,|\, \mu_j \rangle| \\
&= \frac{K}{2|D_j|} \left| \lim_{x \to +0} \int_{\mathbf{R}} \frac{1}{x + \sqrt{-1}(y_j - \omega)} (Z_{1,1}(t, \omega) - Z_{0,1}(t, \omega)) g(\omega) d\omega \right| \\
&\leq \frac{K}{2|D_j|} \lim_{x \to +0} \int_{\mathbf{R}} \frac{C_1|\omega| + C_1 M_2 + M_3}{|x + \sqrt{-1}(y_j - \omega)|} g(\omega) d\omega \cdot t := M_{5,j}\, t.
\end{aligned}$$

Putting $M = \max\{M_2, M_4, M_{5,j}\}$ yields

$$|x_1 - x_0| \leq Mt, \ |\eta_1 - \eta_0| \leq Mt, \ |\alpha_{1,j} - \alpha_{0,j}| \leq Mt \tag{B.7}$$

for $0 \leq t \leq T_1$. Since f_1, f_2, f_3 are Lipschitz continuous, there exists a positive constant L such that

$$|f_i(x, \eta, \alpha) - f_i(x', \eta', \alpha')| \leq L\left(|x - x'| + |\eta - \eta'| + \sum_{j=0}^{n} |\alpha_j - \alpha'_j|\right), \quad i = 1, 2, 3, \tag{B.8}$$

for $(x, \eta, \alpha) \in R$ and $(x', \eta', \alpha') \in R$. Let us prove by induction that there exists a positive constant N such that inequalities

$$|x_{m+1} - x_m|, \ |\eta_{m+1} - \eta_m|, \ |\alpha_{m+1,j} - \alpha_{m,j}| \leq M \frac{N^m L^m (n+3)^m}{(m+1)!} t^{m+1} \tag{B.9}$$

hold on $0 \leq t \leq T_1$ for $m = 1, 2, \cdots$. Indeed,

$$\begin{aligned}
|x_{m+1} - x_m| &\leq \int_0^t |f_1(x_m, \eta_m, \alpha_m) - f_1(x_{m-1}, \eta_{m-1}, \alpha_{m-1})| ds \\
&\leq \int_0^t L\left(|x_m - x_{m-1}| + |\eta_m - \eta_{m-1}| + \sum_{j=0}^{n} |\alpha_{m,j} - \alpha_{m-1,j}|\right) ds \\
&\leq \int_0^t L(n+3)M \frac{N^{m-1}L^{m-1}(n+3)^{m-1}}{m!} s^m ds = M \frac{N^{m-1}L^m(n+3)^m}{(m+1)!} t^{m+1}.
\end{aligned}$$

Next,

$$\begin{aligned}
& |Z_{m+1,1} - Z_{m,1}| \\
\leq\ & \int_0^{2\pi} |e^{\sqrt{-1}x_{m+1}} - e^{\sqrt{-1}x_m}| \cdot |\hat{h}(\theta, \omega)| d\theta \\
+\ & \frac{1}{2\pi} \int_0^{2\pi} |e^{\sqrt{-1}x_{m+1}} - e^{\sqrt{-1}x_m}| \int_0^t \exp\Big[\int_0^s |f_2(x_m, \eta_m, \alpha_m)| d\tau\Big] \cdot |f_3(x_m, \eta_m, \alpha_m)| ds d\theta \\
+\ & \frac{1}{2\pi} \int_0^{2\pi} \int_0^t \left| \exp\Big[\int_s^0 f_2(x_m, \eta_m, \alpha_m) d\tau\Big] - \exp\Big[\int_s^0 f_2(x_{m-1}, \eta_{m-1}, \alpha_{m-1}) d\tau\Big] \right| \cdot |f_3(x_m, \eta_m, \alpha_m)| ds d\theta \\
+\ & \frac{1}{2\pi} \int_0^{2\pi} \int_0^t \exp\Big[\int_0^s |f_2(x_{m-1}, \eta_{m-1}, \alpha_{m-1})| d\tau\Big] \cdot |f_3(x_m, \eta_m, \alpha_m) - f_3(x_{m-1}, \eta_{m-1}, \alpha_{m-1})| ds d\theta.
\end{aligned}$$

There exists $M_6 \geq 0$ such that

$$\begin{aligned}
& \left| \exp\Big[\int_s^0 f_2(x_m, \eta_m, \alpha_m) d\tau\Big] - \exp\Big[\int_s^0 f_2(x_{m-1}, \eta_{m-1}, \alpha_{m-1}) d\tau\Big] \right| \\
\leq\ & M_6 \int_0^s |f_2(x_m, \eta_m, \alpha_m) - f_2(x_{m-1}, \eta_{m-1}, \alpha_{m-1})| d\tau
\end{aligned}$$

holds when $0 \leq s \leq T_1$. This provides

$$\begin{aligned}
& |Z_{m+1,1} - Z_{m,1}| \\
\leq\ & \int_0^{2\pi} |x_{m+1} - x_m| \cdot |\hat{h}(\theta,\omega)| d\theta + \frac{1}{2\pi}\int_0^{2\pi} |x_{m+1} - x_m| \int_0^t e^{M_2 s} M_2 ds d\theta \\
& + \frac{1}{2\pi}\int_0^{2\pi}\int_0^t\int_0^s M_2 M_6 L \left(|x_m - x_{m-1}| + |\eta_m - \eta_{m-1}| + \sum_{j=0}^n |\alpha_{m,j} - \alpha_{m-1,j}| \right) d\tau ds d\theta \\
& + \frac{1}{2\pi}\int_0^{2\pi}\int_0^t e^{M_2 s} L \left(|x_m - x_{m-1}| + |\eta_m - \eta_{m-1}| + \sum_{j=0}^n |\alpha_{m,j} - \alpha_{m-1,j}| \right) ds d\theta \\
\leq\ & M \frac{N^{m-1} L^m (n+3)^m}{(m+1)!} t^{m+1} \int_0^{2\pi} |\hat{h}(\theta,\omega)| d\theta + M \frac{N^{m-1} L^m (n+3)^m}{(m+1)!} t^{m+1} (e^{M_2 t} - 1) \\
& + \frac{1}{2\pi}\int_0^{2\pi}\int_0^t\int_0^s M_2 M_6 L (n+3) M \frac{N^{m-1} L^{m-1} (n+3)^{m-1}}{m!} \tau^m d\tau ds d\theta \\
& + \frac{1}{2\pi}\int_0^{2\pi}\int_0^t L (n+3) M \frac{N^{m-1} L^{m-1} (n+3)^{m-1}}{m!} s^m e^{M_2 s} ds d\theta \\
\leq\ & M \frac{N^{m-1} L^m (n+3)^m}{(m+1)!} t^{m+1} \left(\int_0^{2\pi} |\hat{h}(\theta,\omega)| d\theta + (e^{M_2 t} - 1) + \frac{M_2 M_6}{m+2} t + e^{M_2 t} \right).
\end{aligned}$$

Take a number $N_1(\omega)$, which is independent of m, so that

$$\int_0^{2\pi} |\hat{h}(\theta,\omega)| d\theta + (e^{M_2 t} - 1) + \frac{M_2 M_6}{m+2} t + e^{M_2 t} \leq N_1(\omega)$$

on the interval $0 \leq t \leq T_1$. By the assumption (F3) for $\hat{h}$, $N_1(\omega)$ is bounded as $\omega \to \pm\infty$. Thus we obtain

$$|\eta_{m+1} - \eta_m| \leq \int_{\mathbf{R}} |Z_{m+1,1} - Z_{m,1}| g(\omega) d\omega \leq M \frac{N^{m-1} L^m (n+3)^m}{(m+1)!} t^{m+1} \int_{\mathbf{R}} N_1(\omega) g(\omega) d\omega.$$

By the assumption (A2) for $g(\omega)$, $N_2 := \int_{\mathbf{R}} N_1(\omega) g(\omega) d\omega$ exists:

$$|\eta_{m+1} - \eta_m| \leq M \frac{N^{m-1} L^m (n+3)^m}{(m+1)!} t^{m+1} \cdot N_2.$$

Similarly, $|\alpha_{m+1,j} - \alpha_{m,j}|$ is estimated as

$$\begin{aligned}
|\alpha_{m+1,j} - \alpha_{m,j}| &\leq \frac{K}{2|D_j|} \lim_{x \to +0} \int_{\mathbf{R}} \frac{1}{|x + \sqrt{-1}(y_j - \omega)|} N_1(\omega) g(\omega) d\omega \cdot M \frac{N^{m-1} L^m (n+3)^m}{(m+1)!} t^{m+1} \\
&:= M \frac{N^{m-1} L^m (n+3)^m}{(m+1)!} t^{m+1} \cdot N_3,
\end{aligned}$$

with some constant $N_3 > 0$. Now Eq.(B.9) is obtained by putting $N = \max\{1, N_2, N_3\}$.

Eq.(B.9) shows that sequences $\{x_m\}, \{\eta_m\}$ and $\{\alpha_m\}$ converge to solutions $x(t, 0; \theta, \omega), \eta(t)$ and $\alpha(t)$ uniformly on $0 \leq t \leq T_1$, respectively. Substituting them into (B.2) provides $\hat{Z}_1, \hat{Z}_2, \cdots$. They give a local solution of Eq.(B.1) on the interval $0 \leq t \leq T_1$.

Next thing to do is to show that the solution is extended to any $t \geq 0$. For this purpose, let us show that $x(t, 0; \theta, \omega), \eta(t)$ and $\alpha(t)$ are bounded for any bounded $t \geq 0$ if they exist.

Suppose that there exists $t_0 > 0$ such that $|\eta(t)| \geq 2$ for $t \geq t_0$. Then, $\chi(|\eta(t)|) = 0$, and thus Eq.(7.11) is reduced to the linear equation $d\hat{Z}_1/dt = T_{10}\hat{Z}_1$. This shows that when $t \geq t_0$, $\eta(t)$ is given by $\eta(t) = (e^{T_{10}(t-t_0)}\hat{Z}_1(t_0, \omega), P_0)$, which is bounded for every $t \geq t_0$ due to Prop.5.13. Thus $\eta(t)$ is bounded for every $t \geq 0$. In a similar manner, we can verify that $\alpha_0(t), \cdots, \alpha_n(t)$ are bounded for every $t \geq 0$ because if $|\alpha_j| \geq 2\delta_1$, then $\chi(||P_c z||_{E_c}/\delta_1) = 0$ and Eq.(7.11) is reduced to the linear equation. If $\eta(t)$ and $\alpha(t)$ exist and are bounded, the right hand side of Eq.(7.12) defines a time-dependent vector field on the compact space S^1. Thus a solution $x(t, 0; \theta, \omega)$ exists. Then, substitution of them into Eq.(B.2) shows that $\hat{Z}_1, \hat{Z}_2, \cdots$ are bounded for any bounded $t \geq 0$.

We have proved that if solutions exist, they should be bounded. Now we can extend a local solution defined on $0 \leq t \leq T_1$ in usual way: Repeating the iteration method by using an initial condition given at $t = T_1$, we can extend a local solution to a larger interval $0 \leq t \leq T_2$. Repeating this procedure, we obtain a sequence $T_2 < T_3 < \cdots$ such that a solution exists on $0 \leq t \leq T_n$. Suppose that $T_\infty := \lim_{n\to\infty} T_n$ is finite. Then, since the solution is bounded at $t = T_\infty$, we again use the iteration method with the initial condition given at $t = T_\infty$ to extend the solution beyond T_∞. Therefore, the solution can be extended for every $t \geq 0$. This proves that Eq.(B.1) has a solution for any $t \geq 0$. The uniqueness of solutions is proved by a standard argument using the Gronwall inequality and is omitted here. ■

Proposition B.2. A solution $x(t, 0; \theta, \omega)$ of Eq.(7.12) has an analytic continuation with respect to $\theta \in \mathbf{C}$ and $\omega \in \mathbf{C}$.

Proof. Once $\eta(t)$ and $\alpha(t)$ are determined, (7.12) is an ODE of x. Since the right hand side of (7.12) is analytic in the parameter ω and the dependent variable x, the proposition follows from the fundamental theorem of ODE. ■

At this time, Prop.7.2 is applicable: solutions of Eq.(7.11) exist in $\mathcal{F}$, and thus solutions of Eq.(7.30) exist in $\mathbf{R} \times i(\mathcal{F})$ for any $t \geq 0$.

Proposition B.3. Solutions of Eq.(7.30) in $\mathbf{R} \times i(\mathcal{F})$ are continuous in initial conditions $(|\overline{Z_1(0)}\rangle, |\overline{Z_2(0)}\rangle, \cdots)$ with respect to the metric d_∞.

One of the modern ways to prove the continuity of solutions of ODEs is to use the implicit function theorem on a Banach space. Unfortunately, our space $i(\mathcal{F})$ is not a Banach space and the implicit function theorem does not hold in general. Thus we use more "classical" way by using the iteration method.

Proof. Take two elements $Z(0) = (Z_1(0), Z_2(0), \cdots)$ and $Z^*(0) = (Z_1^*(0), Z_2^*(0), \cdots)$ in $\mathcal{F}$. Let $(|\overline{Z_1(t)}\rangle, |\overline{Z_2(t)}\rangle, \cdots)$ and $(|\overline{Z_1'(t)}\rangle, |\overline{Z_2'(t)}\rangle, \cdots)$ be solutions of (7.30) in $i(\mathcal{F})$ with the initial conditions

$$(|\overline{Z_1(0)}\rangle, |\overline{Z_2(0)}\rangle, \cdots), \quad \text{and} \quad (|\overline{Z_1(0)}\rangle - \kappa|\overline{Z_1^*(0)}\rangle, |\overline{Z_2(0)}\rangle - \kappa|\overline{Z_2^*(0)}\rangle, \cdots),$$

respectively, where $\kappa \in \mathbf{R}$ and $|\,\overline{Z_k(0)}\rangle = i(Z_k(0)), |\,\overline{Z_k^*(0)}\rangle = i(Z_k^*(0))$. We want to prove that if $|\,\overline{Z_k^*(0)}\rangle \to 0$ with respect to the weak dual topology uniformly in $k = 1, 2, \cdots$, then $|\,\overline{Z_k'(t)}\rangle \to |\,\overline{Z_k(t)}\rangle$ uniformly in $k = 1, 2, \cdots$. The constant κ is introduced to prove the next Prop.B.4. In this proof, the reader can assume that $\kappa = 1$. Since $Z(0), Z^*(0) \in \mathcal{F}$, there exist signed measures h and h^* such that

$$Z_k(0) = \int_0^{2\pi} e^{\sqrt{-1}k\theta} h(\theta, \omega) d\theta, \quad Z_k^*(0) = \int_0^{2\pi} e^{\sqrt{-1}k\theta} h^*(\theta, \omega) d\theta, \tag{B.10}$$

for $k = 1, 2, \cdots$. Put $h'(\theta, \omega) = h(\theta, \omega) - \kappa h^*(\theta, \omega)$. Then, $Z_k(t)$ and $Z_k'(t)$ are Fourier coefficients of solutions $\hat{\rho}$ and $\hat{\rho}'$ of Eq.(7.9) satisfying the initial conditions h and h', respectively, and $i(Z_k(t)) = |\,\overline{Z_k(t)}\rangle$, $i(Z_k'(t)) = |\,\overline{Z_k'(t)}\rangle$. Let (x, η, α) and (x', η', α') be corresponding variables; x and x' are solutions of the first equation of (B.1), in which $\hat{h}$ is replaced by h and h', respectively; $\eta(t) = \langle P_0\,|\,\overline{Z}_1\rangle$, $\alpha_j(t) = K/(2\overline{D}_j) \cdot \langle \overline{Z_1\,|\,\mu_j}\rangle$, and η', α' are defined in a similar manner. Put

$$y = x - x', \ Y_k = Z_k - Z_k', \ \xi = \eta - \eta', \ \gamma_j = \alpha_j - \alpha_j'. \tag{B.11}$$

Then, they satisfy the integral equations

$$\left\{ \begin{array}{l} y(t, 0; \theta, \omega) = \displaystyle\int_0^t \left(f_1(x, \eta, \alpha) - f_1(x - y, \eta - \xi, \alpha - \gamma) \right) ds, \\ Y_k(t, \omega) = \displaystyle\int_0^{2\pi} e^{\sqrt{-1}kx} h(\theta, \omega) d\theta - \int_0^{2\pi} e^{\sqrt{-1}k(x-y)} h'(\theta, \omega) d\theta \\ \qquad\qquad + \displaystyle\frac{1}{2\pi} \int_0^{2\pi} e^{\sqrt{-1}kx} \int_0^t \exp\Big[\int_s^0 f_2(x, \eta, \alpha) d\tau \Big] f_3(x, \eta, \alpha) ds d\theta \\ \qquad\qquad - \displaystyle\frac{1}{2\pi} \int_0^{2\pi} e^{\sqrt{-1}k(x-y)} \int_0^t \exp\Big[\int_s^0 f_2(x - y, \eta - \xi, \alpha - \gamma) d\tau \Big] f_3(x - y, \eta - \xi, \alpha - \gamma) ds d\theta, \\ \xi(t) = \langle P_0\,|\,\overline{Y}_1\rangle, \\ \gamma_j(t) = \displaystyle\frac{K}{2\overline{D}_j} \langle \overline{Y_1\,|\,\mu_j}\rangle. \end{array} \right. \tag{B.12}$$

Let us show by the iteration method that these equations have solutions which tend to zero as $|\,\overline{Y_k(0, \omega)}\rangle = \kappa |\,\overline{Z_k^*(0)}\rangle \to 0$. Define sequences $\{y_m\}_{m=0}^\infty, \{Y_{m,k}\}_{m=0}^\infty, \{\xi_m\}_{m=0}^\infty$ and $\{\gamma_{m,j}\}_{m=0}^\infty$ by

$$y_0 = 0, \quad Y_{0,k} = Z_k(0) - Z_k'(0) = \kappa Z_k^*(0) = \kappa \int_0^{2\pi} e^{\sqrt{-1}k\theta} h^*(\theta, \omega) d\theta, \tag{B.13}$$

and

$$\begin{cases} y_{m+1} = \displaystyle\int_0^t (f_1(x,\eta,\alpha) - f_1(x-y_m,\eta-\xi_m,\alpha-\gamma_m))\,ds, \\ Y_{m+1,k} = \displaystyle\int_0^{2\pi} e^{\sqrt{-1}kx} h(\theta,\omega)d\theta - \int_0^{2\pi} e^{\sqrt{-1}k(x-y_{m+1})} h'(\theta,\omega)d\theta \\ \quad +\dfrac{1}{2\pi}\displaystyle\int_0^{2\pi} e^{\sqrt{-1}kx}\int_0^t \exp\Big[\int_s^0 f_2(x,\eta,\alpha)d\tau\Big] f_3(x,\eta,\alpha)dsd\theta \\ \quad -\dfrac{1}{2\pi}\displaystyle\int_0^{2\pi} e^{\sqrt{-1}k(x-y_{m+1})}\int_0^t \exp\Big[\int_s^0 f_2(x-y_m,\eta-\xi_m,\alpha-\gamma_m)d\tau\Big] f_3(x-y_m,\eta-\xi_m,\alpha-\gamma_m)dsd\theta, \\ \xi_m(t) = \langle P_0\,|\,\overline{Y}_{m,1}\rangle, \\ \gamma_{m,j}(t) = \dfrac{K}{2\overline{D}_j}\langle \overline{Y_{m,1}\,|\,\mu_j}\rangle. \end{cases} \tag{B.14}$$

Since $\xi_0(t) = \langle P_0\,|\,\overline{Y}_{0,1}\rangle = \kappa\langle P_0\,|\,\overline{Z_1^*(0)}\rangle$, $\xi_0(t)\to 0$ as $|\,\overline{Z_1^*(0)}\rangle \to 0$ with respect to the weak dual topology. Since $\gamma_{0,j}(t) = K/2\overline{D}_j\cdot\kappa\langle\overline{Z_1^*(0)\,|\,\mu_j}\rangle = \kappa\Pi_j|\,\overline{Z_1^*(0)}\rangle$ and since Π_j is continuous on $i(V)$, $\gamma_{0,j}\to 0$ as $|\,\overline{Z_1^*(0)}\rangle\to 0$ in $i(V)$. To estimate Eq.(B.14), we use the same notation as the proof of Prop.B.1. For a small interval $0\le t\le T_1$, there exists a constant $L>0$ such that

$$\begin{aligned} |y_1-y_0| &\le \int_0^t |f_1(x,\eta,\alpha)-f_1(x-y_0,\eta-\xi_0,\alpha-\gamma_0)|ds \\ &\le L\int_0^t(|y_0|+|\xi_0|+|\gamma_0|)ds = Lt(|\xi_0|+|\gamma_0|)\sim O(\kappa t), \end{aligned} \tag{B.15}$$

where $|\gamma_0| = |\gamma_{0,0}|+|\gamma_{0,1}|+\cdots+|\gamma_{0,n}|$. This shows that

$$|y_1-y_0| = |y_1|\to 0, \tag{B.16}$$

as $|\,\overline{Z_1^*(0)}\rangle\to 0$. Next, for any $\phi\in F_p$,

$$\begin{aligned} &|\langle\phi\,|\,\overline{Y}_{1,k}-\overline{Y}_{0,k}\rangle| \\ \le\ & \Big|\kappa\int_{\mathbf{R}}\int_0^{2\pi}(e^{\sqrt{-1}k(x-y_1)}-e^{\sqrt{-1}k\theta})\phi(\omega)h^*(\theta,\omega)g(\omega)d\theta d\omega\Big| \\ &+\int_{\mathbf{R}}\int_0^{2\pi}|e^{\sqrt{-1}kx}-e^{\sqrt{-1}k(x-y_1)}|\cdot|\phi(\omega)h(\theta,\omega)|g(\omega)d\theta d\omega \\ &+\frac{1}{2\pi}\int_{\mathbf{R}}\int_0^{2\pi}|e^{\sqrt{-1}kx}-e^{\sqrt{-1}k(x-y_1)}|\int_0^t\exp\Big[\int_0^s|f_2(x,\eta,\alpha)|d\tau\Big]|f_3(x,\eta,\alpha)|\cdot|\phi(\omega)|g(\omega)dsd\theta d\omega \\ &+\frac{1}{2\pi}\int_{\mathbf{R}}\int_0^{2\pi}\int_0^t\Big|\exp\Big[\int_s^0 f_2(x,\eta,\alpha)d\tau\Big]-\exp\Big[\int_s^0 f_2(x-y_0,\eta-\xi_0,\alpha-\gamma_0)d\tau\Big]\Big|\times \\ &\qquad\qquad |f_3(x,\eta,\alpha)|\cdot|\phi(\omega)|g(\omega)dsd\theta d\omega \\ &+\frac{1}{2\pi}\int_{\mathbf{R}}\int_0^{2\pi}\int_0^t\exp\Big[\int_0^s|f_2(x,\eta,\alpha)|d\tau\Big]|f_3(x,\eta,\alpha)-f_3(x-y_0,\eta-\xi_0,\alpha-\gamma_0)|\,|\phi(\omega)|g(\omega)dsd\theta d\omega. \end{aligned}$$

Let us show that $|\langle \phi \,|\, \overline{Y}_{1,k} - \overline{Y}_{0,k} \rangle| \to 0$ as $|\,\overline{Z_j^*(0)}\rangle \to 0$ uniformly in $k = 1, 2 \cdots$, and it is of order $O(\kappa t)$. Since $y_1, \xi_0, \gamma_0 \to 0$ as $|\,\overline{Z_1^*(0)}\rangle \to 0$ and since $\xi_0, \gamma_0 \sim O(\kappa), y_0 \sim O(\kappa t)$, it is easy to verify that the second term to the fifth term in the right hand side above tend to zero as $|\,\overline{Z_1^*(0)}\rangle \to 0$ uniformly in $k = 1, 2, \cdots$, and they have an order $O(\kappa t)$. Next, we estimate the first term $A := \int_{\mathbf{R}} \int_0^{2\pi} (e^{\sqrt{-1}k(x-y_1)} - e^{\sqrt{-1}k\theta}) \phi(\omega) h^*(\theta, \omega) g(\omega) d\theta d\omega$ as

$$
\begin{aligned}
|A| \leq & \int_{\mathbf{R}} \int_0^{2\pi} k|x - y_1 - \theta| \cdot |\phi(\omega) h^*(\theta, \omega) g(\omega)| d\theta d\omega \\
\leq & \int_{\mathbf{R}} \int_0^{2\pi} k|\omega t + \int_0^t f_1(x, \eta, \alpha) ds - y_1| \cdot |\phi(\omega) h^*(\theta, \omega) g(\omega)| d\theta d\omega \\
\leq & \, kt \int_{\mathbf{R}} \int_0^{2\pi} |\omega + M_2| \cdot |\phi(\omega) h^*(\theta, \omega) g(\omega)| d\theta d\omega + k|y_1| \int_{\mathbf{R}} \int_0^{2\pi} |\phi(\omega) h^*(\theta, \omega) g(\omega)| d\theta d\omega. \quad \text{(B.17)}
\end{aligned}
$$

This and Eq.(B.15) show $A \sim O(\kappa t)$, and thus $|\langle \phi \,|\, \overline{Y}_{1,k} - \overline{Y}_{0,k} \rangle| \sim O(\kappa t)$. On the other hand, with the aid of the Fourier expansion of the factor $e^{\sqrt{-1}k(x-y_1)}$ in A, we can verify that A tends to zero uniformly in $k = 1, 2 \cdots$ as $\langle \phi \,|\, \overline{Z_j^*(0)} \rangle$ tends to zero uniformly for $j = 1, 2 \cdots$;

$$
\langle \phi \,|\, \overline{Z_j^*(0)} \rangle = \int_{\mathbf{R}} \int_0^{2\pi} e^{\sqrt{-1}j\theta} \phi(\omega) h^*(\theta, \omega) g(\omega) d\theta d\omega \to 0, \quad j = 1, 2, \cdots. \tag{B.18}
$$

These facts prove that there exists a number $B = B(k, \phi, Z_k^*(0))$ such that

$$
|\langle \phi \,|\, \overline{Y}_{1,k} - \overline{Y}_{0,k} \rangle| \leq B \cdot \kappa t,
$$

and $B \to 0$ as $|\,\overline{Z_k^*(0)}\rangle \to 0$ uniformly in k. Therefore, $\xi_1 - \xi_0$ and $\gamma_{1,j} - \gamma_{0,j}$ are also of order $O(\kappa t)$ and they tend to zero as $|\,\overline{Z_k^*(0)}\rangle \to 0$. With these estimates, we can show by induction that $|y_{m+1} - y_m|, |\langle \phi \,|\, \overline{Y}_{m+1,k} - \overline{Y}_{m,k} \rangle|, |\xi_{m+1} - \xi_m|$ and $|\gamma_{m+1,j} - \gamma_{m,j}|$ have order $O(\kappa t^n / n!)$, and they tend to zero as $|\,\overline{Z_k^*(0)}\rangle \to 0$. Thus the sequences converge to solutions $y, \langle \phi \,|\, \overline{Y}_k \rangle, \xi$ and γ_j of Eq.(B.12), and it turns out that the solution is of order $O(\kappa)$ and it tends to zero as $|\,\overline{Z_k^*(0)}\rangle \to 0$ uniformly in k. The detailed calculation is the same as the proof of Prop.B.1. In particular, since $\langle \phi \,|\, \overline{Z_k(t)} - \overline{Z_k'(t)} \rangle \to 0$ as $|\,\overline{Z_k^*(0)}\rangle \to 0$ uniformly in k, this result means that solutions of Eq.(7.30) are continuous in initial conditions with respect to the weak dual topology. As was remarked in Sec.7.4, this also proves that solutions are continuous with respect to the metric d_∞. ■

The remaining task to complete the proof of Prop.7.3 is to show that solutions are differentiable with respect to initial conditions.

Proposition B.4. Solutions of Eq.(7.30) in $\mathbf{R} \times i(\mathcal{F})$ are Fréchet differentiable in initial conditions with respect to the metric d_∞.

Proof. We use the same notation as the proof of Prop.B.3. Define two solutions $(|\,\overline{Z_1(t)}\rangle, |\,\overline{Z_2(t)}\rangle, \cdots)$ and $(|\,\overline{Z_1'(t)}\rangle, |\,\overline{Z_2'(t)}\rangle, \cdots)$ as the proof of Prop.B.3. We consider the limit $\kappa \to 0$ for fixed

$(|\,\overline{Z_1^*(0)}\rangle, |\,\overline{Z_2^*(0)}\rangle, \cdots)$. In the proof, we have proved that the integral equation (B.12) has a solution, which is of $O(\kappa)$ as $\kappa \to 0$. Therefore, the following quantities exist:

$$\begin{cases} \tilde{y}(t, Z(0), Z^*(0)) := \lim\limits_{\kappa\to 0} \dfrac{y}{\kappa} = \lim\limits_{\kappa\to 0} \dfrac{x - x'}{\kappa}, & \tilde{y}(0, Z(0), Z^*(0)) = 0, \\ \tilde{Y}_k(t, Z(0), Z^*(0)) := \lim\limits_{\kappa\to 0} \dfrac{Y_k}{\kappa} = \lim\limits_{\kappa\to 0} \dfrac{Z_k - Z'_k}{\kappa}, & \tilde{Y}_k(0, Z(0), Z^*(0)) = Z_k^*(0), \\ \tilde{\xi}(t, Z(0), Z^*(0)) := \lim\limits_{\kappa\to 0} \dfrac{\xi}{\kappa} = \lim\limits_{\kappa\to 0} \dfrac{\eta - \eta'}{\kappa}, & \tilde{\xi}(0, Z(0), Z^*(0)) = \langle P_0\,|\,\overline{Z_1^*(0)}\rangle, \\ \tilde{\gamma}_j(t, Z(0), Z^*(0)) := \lim\limits_{\kappa\to 0} \dfrac{\gamma_j}{\kappa} = \lim\limits_{\kappa\to 0} \dfrac{\alpha_j - \alpha'_j}{\kappa}, & \tilde{\gamma}_j(0, Z(0), Z^*(0)) = \dfrac{K}{2\overline{D}_j}\overline{\langle Z_1^*(0)\,|\,\mu_j\rangle}, \end{cases} \tag{B.19}$$

where $Z(0) = (Z_1(0), Z_2(0), \cdots)$ and $Z^*(0) = (Z_1^*(0), Z_2^*(0), \cdots)$. This implies that $|\,\overline{Z_k(t)}\rangle$ is Gâteaux differentiable and $i(\tilde{Y}_k) = |\,\overline{\tilde{Y}_k}\rangle$ is the Gâteaux derivative. Let us show that $i(\tilde{Y}_k) = |\,\overline{\tilde{Y}_k}\rangle$ is the Fréchet derivative of $|\,\overline{Z_k(t)}\rangle$ with respect to the initial condition. Dividing by κ and taking the limit $\kappa \to 0$ in Eq.(B.12) yields

$$\tilde{y}(t, Z(0), Z^*(0)) = \int_0^t \nabla f_1(x, \eta, \alpha)(\tilde{y}, \tilde{\xi}, \tilde{\gamma})^T ds, \tag{B.20}$$

and

$$\begin{aligned} &\tilde{Y}_k(t, Z(0), Z^*(0)) \\ = \;& \int_0^{2\pi} e^{\sqrt{-1}kx} \sqrt{-1}\, k\tilde{y}(t, Z(0), Z^*(0)) h(\theta, \omega) d\theta + \int_0^{2\pi} e^{\sqrt{-1}kx} h^*(\theta, \omega) d\theta \\ & + \frac{1}{2\pi} \int_0^{2\pi} e^{\sqrt{-1}kx} \sqrt{-1}\, k\tilde{y}(t, Z(0), Z^*(0)) \int_0^t \exp\Bigl[\int_s^0 f_2(x, \eta, \alpha) d\tau\Bigr] f_3(x, \eta, \alpha) ds d\theta \\ & + \frac{1}{2\pi} \int_0^{2\pi} e^{\sqrt{-1}kx} \int_0^t \exp\Bigl[\int_s^0 f_2(x, \eta, \alpha) d\tau\Bigr] f_3(x, \eta, \alpha) \int_s^0 \nabla f_2(x, \eta, \alpha)(\tilde{y}, \tilde{\xi}, \tilde{\gamma})^T d\tau ds d\theta \\ & + \frac{1}{2\pi} \int_0^{2\pi} e^{\sqrt{-1}kx} \int_0^t \exp\Bigl[\int_s^0 f_2(x, \eta, \alpha) d\tau\Bigr] \cdot \nabla f_3(x, \eta, \alpha)(\tilde{y}, \tilde{\xi}, \tilde{\gamma})^T ds d\theta, \end{aligned} \tag{B.21}$$

where

$$\nabla f_i(x, \eta, \alpha)(\tilde{y}, \tilde{\xi}, \tilde{\gamma})^T = \frac{\partial f_i}{\partial x}(x, \eta, \alpha)\tilde{y} + \frac{\partial f_i}{\partial \eta}(x, \eta, \alpha)\tilde{\xi} + \frac{\partial f_i}{\partial \alpha}(x, \eta, \alpha)\tilde{\gamma}.$$

Integral equations for $\tilde{\xi}$ and $\tilde{\gamma}_j$ are derived in a similar manner. Then, they define a system of linear integral equations of $\tilde{y}, \tilde{Y}_k, \tilde{\xi}, \tilde{\gamma}_j$, and initial conditions of them are linear in $Z^*(0) = (Z_1^*(0), Z_2^*(0), \cdots)$. This implies that solutions of these integral equations are linear with respect to $Z^*(0)$. Therefore, we can write

$$\tilde{Y}_k(t, Z(0), Z^*(0)) = dY_k(t, Z(0)) Z^*(0), \tag{B.22}$$

and

$$|\overline{\tilde{Y}_k(t, Z(0), Z^*(0))}\rangle = dY_k(t, Z(0))^\times |\overline{Z^*(0)}\rangle, \tag{B.23}$$

Then, $(dY_1(t, Z(0)), dY_2(t, Z(0)), \cdots)$ defines a linear operator from $\mathcal{F}$ into $\mathcal{F}$, and the dual operator $(dY_1(t, Z(0))^\times, dY_2(t, Z(0))^\times, \cdots)$ defines a linear operator from $i(\mathcal{F})$ into $i(\mathcal{F})$. For the integral equations of $\tilde{y}, \tilde{Y}_k, \tilde{\xi}, \tilde{\gamma}_j$ above, we again use the iteration method to prove that $\tilde{Y}_k(t, Z(0), Z^*(0))$ is continuous in a parameter $Z(0)$ and an initial condition $Z^*(0)$ with respect to the weak dual topology uniformly in k (the detail is the same as the proof of Prop.B.3). Therefore, the mapping $(dY_1(t, \cdot)^\times, dY_2(t, \cdot)^\times, \cdots) : i(\mathcal{F}) \times i(\mathcal{F}) \to i(\mathcal{F})$ is continuous with respect to the metric d_∞, which means that solutions of Eq.(7.30) are C^1 in initial conditions and $(dY_1(t, Z(0))^\times, dY_2(t, Z(0))^\times, \cdots)$ gives the Fréchet derivative at $|\overline{Z(0)}\rangle = (|\overline{Z_1(0)}\rangle, |\overline{Z_2(0)}\rangle, \cdots)$. ■

Acknowledgements.

This work was supported by Grant-in-Aid for Young Scientists (B), No.22740069 from MEXT Japan.

References

[1] J. A. Acebrón, L. L. Bonilla, C. J. P. Vicente, F. Ritort, R. Spigler, The Kuramoto model: A simple paradigm for synchronization phenomena, Rev. Mod. Phys., Vol. 77 (2005), pp. 137-185

[2] J. A. Acebrón, L. L. Bonilla, Asymptotic description of transients and synchronized states of globally coupled oscillators, Phys. D 114 (1998), no. 3-4, 296–314

[3] L. V. Ahlfors, Complex analysis. An introduction to the theory of analytic functions of one complex variable, McGraw-Hill Book Co., New York, 1978

[4] N. J. Balmforth, R. Sassi, A shocking display of synchrony, Phys. D 143 (2000), no. 1-4, 21–55

[5] P. W. Bates, C. K. R. T Jones, Invariant manifolds for semilinear partial differential equations, Dynamics reported, Vol. 2, 1–38, 1989

[6] W. Bergweiler, A. Eremenko, On the singularities of the inverse to a meromorphic function of finite order, Rev. Mat. Iberoamericana 11 (1995), no. 2, 355–373

[7] L. L. Bonilla, John C. Neu, R. Spigler, Nonlinear stability of incoherence and collective synchronization in a population of coupled oscillators, J. Statist. Phys. 67 (1992), no. 1-2, 313–330

[8] L. L. Bonilla, C. J. Pérez Vicente, R. Spigler, Time-periodic phases in populations of nonlinearly coupled oscillators with bimodal frequency distributions, Phys. D 113 (1998), no. 1, 79–97

[9] X-Y. Chen, J. K. Hale, T. Bin, Invariant foliations for C^1 semigroups in Banach spaces, J. Differential Equations 139 (1997), no. 2, 283–318

[10] H. Chiba, Continuous limit and the moments system for the globally coupled phase oscillators, (submitted)

[11] H. Chiba, D. Pazó, Stability of an $[N/2]$-dimensional invariant torus in the Kuramoto model at small coupling, Physica D, Vol.238, 1068-1081 (2009)

[12] J. D. Crawford, Amplitude expansions for instabilities in populations of globally-coupled oscillators, J. Statist. Phys. 74 (1994), no. 5-6, 1047–1084

[13] J. D. Crawford, Scaling and Singularities in the Entrainment of Globally Coupled Oscillators, Phys. Rev. Lett. 74, 4341 (1995)

[14] J. D. Crawford, K. T. R. Davies, Synchronization of globally coupled phase oscillators: singularities and scaling for general couplings, Phys. D 125 (1999), no. 1-2, 1–46

[15] H. Daido, Onset of cooperative entrainment in limit-cycle oscillators with uniform all-to-all interactions: bifurcation of the order function, Phys. D 91 (1996), no. 1-2, 24–66

[16] A. E. Eremenko, The set of asymptotic values of a finite order meromorphic function, (Russian) Mat. Zametki 24 (1978), no. 6, 779–783

[17] I. M. Gelfand, G. E. Shilov, Generalized functions. Vol. 2. Spaces of fundamental and generalized functions, Academic Press, New York-London, 1968

[18] I. M. Gelfand, N. Ya. Vilenkin, Generalized functions. Vol. 4. Applications of harmonic analysis, Academic Press, New York-London, 1964

[19] W. Gross, Eine ganze Funktion, fur die jede komplexe Zahl Konvergenzwert ist, Math. Ann. 79 (1918), no. 1-2, 201–208

[20] A. Grothendieck, Topological vector spaces, Gordon and Breach Science Publishers, New York-London-Paris, 1973

[21] E. Hille, R. S. Phillips, Functional analysis and semigroups, American Mathematical Society, 1957

[22] T. Kato, Perturbation theory for linear operators, Springer-Verlag, Berlin, 1995

[23] H. Komatsu, Projective and injective limits of weakly compact sequences of locally convex spaces, J. Math. Soc. Japan, 19, 1967 366–383

[24] T. Krisztin, Invariance and noninvariance of center manifolds of time-t maps with respect to the semiflow, SIAM J. Math. Anal. 36 (2004/05), no. 3, 717–739

[25] Y. Kuramoto, Self-entrainment of a population of coupled non-linear oscillators, International Symposium on Mathematical Problems in Theoretical Physics, pp. 420–422. Lecture Notes in Phys., 39. Springer, Berlin, 1975

[26] Y. Kuramoto, Chemical oscillations, waves, and turbulence, Springer Series in Synergetics, 19. Springer-Verlag, Berlin, 1984

[27] Y. Maistrenko, O. Popovych, O. Burylko, P. A. Tass, Mechanism of desynchronization in the finite-dimensional Kuramoto model, Phys. Rev. Lett. 93 (2004) 084102

[28] Y. L. Maistrenko, O. V. Popovych, P. A. Tass, Chaotic attractor in the Kuramoto model, Int. J. of Bif. and Chaos 15 (2005) 3457–3466

[29] K. Maurin, General eigenfunction expansions and unitary representations of topological groups, Polish Scientific Publishers, Warsaw, 1968

[30] E. A. Martens, E. Barreto, S. H. Strogatz, E. Ott ,P. So, T. M. Antonsen, Exact results for the Kuramoto model with a bimodal frequency distribution, Phys. Rev. E 79, 026204 (2009)

[31] S. A. Marvel, R. E. Mirollo, S. H. Strogatz, Identical phase oscillators with global sinusoidal coupling evolve by Mobius group action, Chaos 19, 043104 (2009)

[32] R. E. Mirollo, S. H. Strogatz, Amplitude death in an array of limit-cycle oscillators, J. Statist. Phys. 60 (1990), no. 1-2, 245–262

[33] R. Mirollo, S. H. Strogatz, The spectrum of the partially locked state for the Kuramoto model, J. Nonlinear Sci. 17 (2007), no. 4, 309–347

[34] D. J. Newman, A simple proof of Wiener's $1/f$ theorem. Proc. Amer. Math. Soc. 48 (1975), 264–265

[35] E. Ott, T. M. Antonsen, Low dimensional behavior of large systems of globally coupled oscillators, Chaos 18 (2008), no. 3, 037113

[36] E. Ott, T. M. Antonsen, Long time evolution of phase oscillator systems, Chaos 19 (2009), no. 2, 023117

[37] A. Pikovsky, M. Rosenblum, J. Kurths, Synchronization: A Universal Concept in Nonlinear Sciences, Cambridge University Press, Cambridge, 2001

[38] M. Reed, B. Simon, Methods of modern mathematical physics IV. Analysis of operators, Academic Press, New York-London, 1978

[39] J. A. Sanders, F. Verhulst, Averaging methods in nonlinear dynamical systems, Springer-Verlag, New York, 1985

[40] J. A. Shohat, J. D. Tamarkin, The Problem of Moments, American Mathematical Society, New York, 1943

[41] E. M. Stein, G. Weiss, Introduction to Fourier analysis on Euclidean spaces, Princeton University Press, Princeton, 1971

[42] S. H. Strogatz, From Kuramoto to Crawford: exploring the onset of synchronization in populations of coupled oscillators, Phys. D 143 (2000), no. 1-4, 1–20

[43] S. H. Strogatz, R. E. Mirollo, Stability of incoherence in a population of coupled oscillators, J. Statist. Phys. 63 (1991), no. 3-4, 613–635

[44] S. H. Strogatz, R. E. Mirollo, P. C. Matthews, Coupled nonlinear oscillators below the synchronization threshold: relaxation by generalized Landau damping, Phys. Rev. Lett. 68 (1992), no. 18, 2730–2733

[45] E. C. Titchmarsh, Introduction to the theory of Fourier integrals, Chelsea Publishing Co., New York, 1986

[46] F. Tréves, Topological vector spaces, distributions and kernels, Academic Press, New York-London, 1967

[47] A. Vanderbauwhede, G. Iooss, Center manifold theory in infinite dimensions, Dynam. Report. Expositions Dynam. Systems, 1, Springer, Berlin, 1992

[48] N. J. Vilenkin, Special functions and the theory of group representations, American Mathematical Society, 1968

[49] K. Yosida, Functional analysis, Springer-Verlag, Berlin, 1995

[50] G. H. Zhang, Theory of entire and meromorphic functions, Deficient and asymptotic values and singular directions, American Mathematical Society, 1993